


SHORT REPORT

Open Access



The mechanism of (+) taxifolin's protective antioxidant effect for $\bullet\text{OH}$ -treated bone marrow-derived mesenchymal stem cells

Xican Li^{1,2*} , Hong Xie^{1,2†}, Qian Jiang¹, Gang Wei^{1,2}, Lishan Lin¹, Changying Li¹, Xingmei Ou¹, Lichan Yang¹, Yulu Xie^{1,2}, Zhen Fu^{3,4}, Yamei Liu^{3,4} and Dongfeng Chen^{4*}

* Correspondence: lixican@126.com;
lixc@gzucm.edu.cn;
CDF27212@21cn.com;
chen888@gzucm.edu.cn

[†]Equal contributors

¹School of Chinese Herbal Medicine, Guangzhou University of Chinese Medicine, Waihuang East Road No. 232, Guangzhou Higher Education Mega Center, Guangzhou 510006, China

⁴The Research Center of Integrative Medicine, Guangzhou University of Chinese Medicine, Guangzhou 510006, China

Full list of author information is available at the end of the article

Abstract

The natural dihydroflavonol (+) taxifolin was investigated for its protective effect on Fenton reagent-treated bone marrow-derived mesenchymal stem cells (bmMSCs). Various antioxidant assays were used to determine the possible mechanism. These included $\bullet\text{OH}$ -scavenging, 2-phenyl-4, 4, 5, 5-tetramethylimidazole-1-oxyl-3-oxide radical-scavenging (PTIO-scavenging), 1, 1-diphenyl-2-picryl-hydrazyl radical-scavenging (DPPH-scavenging), 2, 2'-azino-bis (3-ethylbenzo-thiazoline-6-sulfonic acid) radical-scavenging (ABTS⁺-scavenging), Fe³⁺-reducing, and Cu²⁺-reducing assays. The Fe²⁺-binding reaction was also investigated using UV-Vis spectra. The results revealed that cell viability was fully restored, even increasing to $142.9 \pm 9.3\%$ after treatment with (+) taxifolin. In the antioxidant assays, (+) taxifolin was observed to efficiently scavenge $\bullet\text{OH}$, DPPH \bullet and ABTS⁺ radicals, and to increase the relative Cu²⁺- and Fe³⁺-reducing levels. In the PTIO-scavenging assay, its IC₅₀ values varied with pH. In the Fe²⁺-binding reaction, (+) taxifolin was found to yield a green solution with two UV-Vis absorbance peaks: $\lambda_{\text{max}} = 433 \text{ nm}$ ($\epsilon = 5.2 \times 10^2 \text{ L mol}^{-1} \text{ cm}^{-1}$) and $\lambda_{\text{max}} = 721 \text{ nm}$ ($\epsilon = 5.1 \times 10^2 \text{ L mol}^{-1} \text{ cm}^{-1}$). These results indicate that (+) taxifolin can act as an effective $\bullet\text{OH}$ -scavenger, protecting bmMSCs from $\bullet\text{OH}$ -induced damage. Its $\bullet\text{OH}$ -scavenging action consists of direct and indirect antioxidant effects. Direct antioxidation occurs via multiple pathways, including ET, PCET or HAT. Indirect antioxidation involves binding to Fe²⁺.

Keywords: (+) Taxifolin; bmMSCs, $\bullet\text{OH}$ damage, Antioxidant mechanism, Electron transfer, Fe²⁺ binding

Background

Antioxidant supplementation has been suggested as a means to reduce the DNA damage and relieve oxidative stress during the expansion and proliferation of bone marrow-derived mesenchymal stem cells (bmMSCs) [1]. This oxidative stress is a result of the imbalance between ROS production and diminished endogenous antioxidant protection. Accumulative ROS (especially $\bullet\text{OH}$ with a half-life of 10^{-9} s) not only have the potential to damage all types of biomolecules (such as DNA, proteins, lipids and carbohydrates), but can also inhibit MSC immunomodulation, thus increasing senescence and reducing ex vivo expansion, which is critical for clinical application off the cells [2]. Effective antioxidants that could protect MSCs from oxidative stress are a desirable focus of research.

From the perspective of free radical biology, plants also encounter serious oxidative stress from strong UV-Vis light, atmospheric ROS, temperature changes, and the processes of oxygen consumption for photosynthesis. Notably, some plants, such as pine, have what could be considered a strong vital force and a long history of survival. They have successfully resisted oxidation from complicated ecological environments and may serve as a library of efficient phenolic antioxidants [3].

Pine grows on the Sharon Plain in Israel and in mountains and highlands around the world. Notable species and varieties are *Pinus pinaster* (French maritime pine) [4], *Pseudotsuga amenziesii* [5], *Pinus massoniana* Lamb [6], *Pinus sylvestris* var. *mongovica* Litvin [7] and *Larix olgensis* Henry var. *Koreana* Nakai [8]. Pine has survived for approximately 1.9 hundred million years, suggesting that it possesses strong defenses, probably including a strong antioxidant defense with numerous antioxidant components. In fact, extract from the bark of French maritime pine has been developed as an antioxidant supplement known commercially as Pycnogenol, which has a bioactive component named (+) taxifolin (2*R*,3*R*-dihydroquercetin, Fig. 1) [4, 9].

As shown in Fig. 1a, (+) taxifolin is actually a dihydroflavonol that exists in the aforementioned pine types. It was reported to inhibit free radical formation at key stages of apoptosis in cellular mitochondria [10] and to correct cerebral ischemia-reperfusion injury [11]. Recently, (+) taxifolin was found to exhibit anticancer and neuroprotective effects [12–14].


This indicates that (+) taxifolin has potential as an antioxidant for protecting MSCs against oxidative stress damage. However, no study has reported on the protective effects of (+) taxifolin towards $\bullet\text{OH}$ -treated bmMSCs.

Here, we applied the methyl thiazolyl tetrazolium (MTT) assay to assess the protective effects of (+) taxifolin on $\bullet\text{OH}$ -treated bmMSCs. We then explored the possible mechanisms for this effect.

Methods

Chemicals and animals

The chemicals (+) taxifolin (CAS number: 480–18-2, 98%), dihydromyricetin (CAS number: 27,200–12-0, 98%), and 4'-*O*-methyltaxifolin (CAS number: 70,411–27-7, 98%) were obtained from Chengdu Biopurify Phytochemicals Ltd. Catechol (CAS



number: 120–80-9, 99.5%) and DNA sodium salt (fish sperm) were purchased from Aladdin Chemistry Co. The 1, 1-diphenyl-2-picryl-hydrazyl radical (DPPH•), (±)-6-hydroxyl-2, 5, 7, 8-tetramethylchromane-2-carboxylic acid (trolox), 2, 9-dimethyl-1, 10-phenanthroline (neocuproine), 3-(2-pyridyl)-5, 6-bis (4-phenylsulfonicacid)-1, 2, 4-triazine (ferrozine), 2, 4, 6-tripyridyltriazine (TPTZ), 2-phenyl-4, 4, 5, 5-tetramethylimidazoline-1-oxyl-3-oxide radical (PTIO•), and methyl thiazolyl tetrazolium (MTT) were purchased from Sigma-Aldrich Shanghai Trading Co. The CCK-8 (BB-4221-2) kits were from Best-Bio Inc. (NH₄)₂ABTS [2, 2'-azino-bis (3-ethylbenzo-thiazoline-6-sulfonic acid diammonium salt)] was obtained from Amresco Chemical Co. Dulbecco's modified Eagle's medium (DMEM), fetal bovine serum (FBS) and trypsin were purchased from Gibco. All other reagents were of analytical grade.

Sprague-Dawley (SD) rats (4 weeks old) were obtained from the Animal Center of Guangzhou University of Chinese Medicine. The protocol was performed under the supervision of the Institutional Animal Ethics Committee at the Guangzhou University of Chinese Medicine.

MTT assay to assess the protective effect against •OH-induced damage

The bmMSCs were cultured according to our previous report [15] with slight modifications. In brief, bone marrow was obtained from the femur and tibia of the rats. Marrow samples were diluted with low-glucose DMEM containing 10% FBS. MSCs were prepared by gradient centrifugation at 900×g for 30 min on 1.073 g/ml Percoll. The prepared cells were detached by treatment with 0.25% trypsin and passaged in culture flasks at 1 × 10⁴/cm². At passage 3, bmMSCs were evaluated for cell homogeneity using CD44 detection via flow cytometry. These cells were used for the subsequent experiments.

The protective effect of (+) taxifolin against •OH-induced bmMSC damage was investigated based on the method described in [16, 17] with slight modifications. Briefly, bmMSCs were seeded at 5000 cells per well into 96-well plates. After adherence for 24 h, bmMSCs were divided into control, model and sample [(+) taxifolin] groups.

In the control group, bmMSCs were incubated for 24 h in DMEM. In the model and sample groups, bmMSCs were incubated in the presence of FeCl₂ (100 μM) followed by H₂O₂ (50 μM). After incubation for 20 min, the mixture of FeCl₂ and H₂O₂ was removed. The bmMSCs in the model group were incubated for 24 h in DMEM, while bmMSCs in the sample group were incubated for 24 h in DMEM with the indicated (+) taxifolin concentrations.

After incubation, 20 μl MTT (5 mg/ml) was added, and the culture was incubated for an additional 3 h. The culture medium was discarded and replaced with 150 μl DMSO. Absorbance was measured at 490 nm on a Bio-Kinetics reader (PE-1420; Bio-Kinetics Corporation). Culture medium containing serum was used for the control group and each sample test was repeated in five independent wells.

Hydroxyl-scavenging assay based on DNA

The hydroxyl-scavenging effect of (+) taxifolin was estimated using a method developed by our laboratory [18]. Briefly, methanol sample solutions (1.2 mg/ml, 20–100 μl) were separately aliquoted into mini tubes. After completely evaporating the methanol solvent

in each tube to dryness, the sample residue was treated with 300 μ l of phosphate buffer (0.2 M, pH 7.4), followed by 50 μ l of DNA sodium (10 mg/ml), 75 μ l of H₂O₂ (33.6 mM), 50 μ l of FeCl₃ (3.2 mM), 100 μ l of Na₂EDTA (0.5 mM) and 75 μ l of ascorbic acid (12 mM). After incubation at 50 °C for 20 min, 250 μ l of trichloroacetic acid (10%, w/v) was added to the tube. After heating the mixture at 105 °C for 15 min with 150 μ l of 2-thiobarbituric acid (TBA, 5% in 1.25% NaOH aqueous solution), the absorbance was measured using a Unico Spectrophotometer UV 2100 against the buffer (blank). The protective effect is expressed as follows:

$$\text{Protective effect}\% = \frac{A_0 - A}{A_0} \times 100\%,$$

where A₀ indicates the absorbance of the blank and A indicates the absorbance of the sample (+) taxifolin.

PTIO•-scavenging assay

The PTIO•-scavenging assay was conducted based on our method [19]. In brief, 80 μ l of an aqueous PTIO• solution (0.1 mM) was mixed with 20 μ l of phosphate buffer at pH 5.0, 6.0, 7.0, 7.4, 8.0 and 9.0 containing 1 mg/ml of sample at the indicated concentrations. The mixture was maintained at 37 °C for 30 min, and the absorbance was measured at 560 nm on a microplate reader (Multiskan FC, Thermo Scientific). The PTIO• inhibition percentage was calculated as follows:

$$\text{Scavenging}\% = \frac{A_0 - A}{A_0} \times 100\%,$$

where A₀ indicates the absorbance of the blank and A indicates the absorbance of the sample, (+) taxifolin.

DPPH•-scavenging assay and ABTS⁺•-scavenging assay

DPPH• radical-scavenging activity was determined as previously described [20]. Briefly, 1 ml of DPPH• solution (0.1 M) was mixed with the indicated concentrations of sample (0.15 mg/ml, 14–70 μ l) dissolved in methanol. The mixture was maintained at room temperature for 30 min, and the absorbance was measured at 519 nm on a Unico Spectrophotometer 2100.

ABTS⁺•-scavenging activity was evaluated according to a previously described method [21]. ABTS⁺• was produced by mixing 0.2 ml of ABTS diammonium salt (7.4 mM) with 0.35 ml of potassium persulfate (2.6 mM). The mixture was maintained in the dark at room temperature for 12 h to allow completion of radical generation and then diluted with 95% ethanol. To determine the scavenging activity, the test sample (x = 15–75 μ l, 0.03 mg/ml) was added to (200- x) μ l of 95% ethanol followed by 800 μ l of ABTS⁺• reagent, and the absorbance was measured at 734 nm on a Unico Spectrophotometer 2100 6 min after the initial mixing using 95% ethanol as the blank.

The percentage of DPPH•-scavenging (or ABTS⁺•-scavenging) activity was calculated based on the formula given in the **PTIO•-scavenging assay** section.

Cu²⁺-reducing assay

The reducing power capacity of cupric ions (Cu²⁺) was measured according to a previously described method [22] with a slight modification. Briefly, 125 µl of CuSO₄ aqueous solution (10 mM), 125 µl of neocuproine ethanolic solution (7.5 mM) and 750 µl of CH₃COONH₄ buffer solution (0.1 M, pH 7.5) were added to test tubes with different volumes of sample (0.15 mg/ml, 15–75 µl). The total volume was adjusted to 1 ml with buffer and mixed vigorously. The absorbance against a buffer blank was measured at 450 nm after 30 min. An increase in the absorbance of the reaction mixture indicates an increase in reduction capability. The relative reducing power of the sample relative to the maximum absorbance was calculated using the following formula:

$$\text{Relative reducing effect}\% = \frac{A - A_{\min}}{A_{\max} - A_{\min}} \times 100\%,$$

where A_{min} is the absorbance of the control without sample, A is the absorbance of the reaction mixture with sample, and A_{max} is the maximum absorbance of the reaction mixture with sample.

Ferric-reducing antioxidant power (FRAP) assay

The FRAP assay was adapted from Benzie and Strain [23]. Briefly, FRAP reagent was freshly prepared by mixing 10 mM TPTZ, 20 mM FeCl₃ and 0.25 M acetate buffer at 1:1:10 (pH 3.6). The test sample (x = 20–100 µl, 0.5 mg/ml) was added to (100 - x) µl of 95% ethanol followed by 400 µl of FRAP reagent. The absorbance was measured at 593 nm after a 30-min incubation at ambient temperature using distilled water as the blank. The relative reducing power was calculated using the formula given in the **Cu²⁺-reducing assay** section.

UV-vis spectra and color reaction of Fe²⁺-binding

The (+) taxifolin–Fe²⁺ complex was evaluated using UV-Vis spectroscopy. For these experiments, 300 µl of a methanolic solution of (+) taxifolin and 100 µl of an aqueous solution of FeCl₂·4H₂O were added to 600 µl of an aqueous mixture of distilled water and methanol (1:1). The solution was then mixed vigorously and continuously scanned using a UV-Vis spectrophotometer (Unico 2600A) from 200 to 900 nm after 0, 10, 20, 30, and 60 min.


The above experiment was repeated using 4'-O-methyltaxifolin.

Statistical analysis

Each experiment was performed in triplicate and data were recorded as the means ± SD (standard deviation). Dose response curves were plotted using Origin 6.0 software (OriginLab). IC₅₀ was defined as the final concentration of 50% radical inhibition (relative reducing power or binding effect). Statistical comparisons were made using one-way ANOVA to detect significant differences using SPSS 13.0 (SPSS Inc.) for Windows. *p* < 0.05 was considered statistically significant.

Results and discussion

As shown in Fig. 2, in the model group, the bmMSCs damaged by •OH presented only 33.1 ± 4.4% viability. However, following treatment with (+) taxifolin (3.3–164.3 µM),



cell viability was restored or even increased. This result suggests that (+) taxifolin effectively protects bmMSCs from \bullet OH-mediated damage. This is consistent with the recent report that (+) taxifolin could reduce cholesterol oxidation product-induced neuronal apoptosis [24]. At higher concentrations ($>50 \mu\text{g/ml}$, $164.3 \mu\text{M}$), (+) taxifolin could even further promote the viability of bmMSCs, reaching $142.9 \pm 9.3\%$ viability.

To test the possible toxicity to MSCs, the effect of (+) taxifolin towards normal MSCs was measured using the CCK-8 assay (an updated version of the MTT assay). The results indicated that (+) taxifolin ($3.3\text{--}328.7 \mu\text{M}$) had no effect on proliferation and no toxic effect on normal MSCs without \bullet OH-treatment (Additional file 1: Figure S1). These results align with the previous findings that (+) taxifolin could be an apparent exception that could efficiently inhibit the Fenton reaction and superoxide radical formation [25, 26] while being completely nonphototoxic, unlike its analogue quercetin [13, 27]. These results are inconsistent with another previous study that showed taxifolin was toxic to oocytes at higher concentration ($50 \mu\text{g/ml}$, $164.3 \mu\text{M}$) [28].

It is assumed that when (+) taxifolin was mixed with Fenton reagents, some reaction products may be generated to bring about the beneficial (especially protective) effect. In fact, a similar situation is observed with salvianolic acid B, which can increase cell viability to 175.1% [29]. In the case of the salvianolic acid B molecule, some characteristic chemical structures, such as catechol or lactone moieties [29], have been suggested to be partly responsible for the protective effect. The moiety of fused rings (A/B) is also hypothesized to play a role in the process. Some antioxidants comprising 8-hydroxyquinol have been demonstrated to induce MSC proliferation [30, 31]. However, the detailed mechanisms should be investigated further.

Such protective effects from \bullet OH damage have been reported to be related to \bullet OH scavenging [32]. In this study, (+) taxifolin was found to exhibit \bullet OH-scavenging ability in a dose-dependent manner (Additional file 1: Figure S2). The $\text{IC}_{50 \text{ Trolox}}/\text{IC}_{50 (+) \text{ taxifolin}}$ value (1.67; Table 1) suggests that (+) taxifolin is a better \bullet OH scavenger than trolox, which is a standard antioxidant.

Table 1 The IC₅₀ values of (+) taxifolin and trolox in various assays (μM)

Assays	(+) taxifolin	Trolox	Ratio value	
•OH-scavenging	259.2 ± 4.4 ^a	411.4 ± 17.0 ^b	1.59	Average 1.67
PTIO•-scavenging*	663.9 ± 34.4 ^a	736.8 ± 29.9 ^b	1.11	
DPPH•-scavenging	16.0 ± 0.2 ^a	18.5 ± 0.4 ^b	1.16	
ABTS ^{•+} -scavenging	4.6 ± 0.2 ^a	11.4 ± 0.2 ^b	2.48	
Cu ²⁺ -reducing	22.4 ± 0.5 ^a	40.4 ± 1.9 ^b	1.80	
FRAP	33.7 ± 1.0 ^a	62.8 ± 1.0 ^b	1.86	

The IC₅₀ value was defined as the final concentration of 50% radical inhibition (relative reducing power). It was calculated by linear regression analysis, and expressed as the mean ± SD (n = 3). The linear regression was analyzed using Origin 6.0. Mean values with different superscripts (a or b) in the same row are significantly different (p < 0.05). *The assay was conducted at pH 7.4. The ratio value is defined as IC_{50 Trolox}/IC_{50 (+) taxifolin}. The dose–response curves are shown in Additional file 1: Figures S2-S7

•OH scavenging comprises two pathways: direct and indirect. The direct antioxidant pathway directly scavenges the •OH free radical that has been generated via the Fenton reaction. However, •OH is a very transient species so it is impossible to verify whether •OH is directly scavenged. Therefore, we used a stable oxygen-centered radical, PTIO•, for the investigation. As seen in Additional file 1: Figure S3A, (+) taxifolin scavenged the PTIO• radical at various pH values in a dose-dependent manner.

Correspondingly, the IC₅₀ values varied with various pH values: 2.6 ± 0.5, 1.6 ± 0.2, 0.7 ± 0.03, 0.6 ± 0.04, 0.5 ± 0.04 and 0.4 ± 0.02 mM respectively for pH 5.0, 6.0, 7.0, 7.4, 8.0 and 9.0 (Additional file 1: Table S1). This indicates the involvement of the direct antioxidant pathway in •OH scavenging by (+) taxifolin. When the IC₅₀ values were plotted against pH values, a first-order decay curve was observed (Additional file 1: Figure S3B), suggesting that a high level of H⁺ (low pH value) considerably suppressed the PTIO•-scavenging ability of (+) taxifolin. Thus, the radical-scavenging ability of (+) taxifolin is hypothesized to be involved in H⁺ transfer, consistent with the cyclic voltammetry-based evidence [33].

It has been documented that at a pH ≤ 5.0, PTIO• can be scavenged via electron transfer (ET) [33]. Our assay suggests that (+) taxifolin may also scavenge PTIO• at pH 5.0, indicating the involvement of ET in its antioxidant action. This is further supported by its ABTS^{•+}-scavenging, Cu²⁺-reducing and Fe³⁺-reducing (i.e., FRAP) abilities (Additional file 1: Figures S4-S6). ABTS^{•+}-scavenging is considered to be an ET-based pathway [34]. The ABTS^{•+}-scavenging ability of (+) taxifolin indicates the involvement of ET in the antioxidant process. Furthermore, (+) taxifolin increased the relative Cu²⁺-reducing and FRAP-reducing abilities in a concentration-dependent manner. The FRAP (at pH 3.6) and Cu²⁺-reducing activities have been demonstrated to be an ET reaction [35]. It should be noted that the Fe³⁺-reducing potential of flavonoids may also reduce Fe³⁺ into Fe²⁺ to cause pro-antioxidation [36]. It remains unknown whether the pro-antioxidation is linked to (+) taxifolin cytotoxicity to oocytes at higher concentration [28].


In this study, (+) taxifolin efficiently scavenged the DPPH• radical (Additional file 1: Figure S7). DPPH• scavenging is regarded as a hydrogen atom transfer-based multi-pathway [32]. Successful DPPH• scavenging by (+) taxifolin indicated that hydrogen atom transfer may occur in its direct antioxidative process. Moreover, it was recently reported that these direct antioxidative pathways are not exclusive but are rather competitive based on various reaction conditions [34].


Because Fe^{2+} can catalyze the Fenton reaction, where H_2O_2 yields $\bullet\text{OH}$ radicals, an attenuation of Fe^{2+} levels via a binding reaction is considered an indirect antioxidant mechanism to scavenge $\bullet\text{OH}$ radicals [37]. In the indirect antioxidant assay, (+) taxifolin bound to Fe^{2+} to yield a green solution and two Vis absorbance peaks: $\lambda_{\text{max}} = 433 \text{ nm}$ ($\epsilon = 5.2 \times 10^2 \text{ L mol}^{-1} \text{ cm}^{-1}$) and $\lambda_{\text{max}} = 721 \text{ nm}$ ($\epsilon = 5.1 \times 10^2 \text{ L mol}^{-1} \text{ cm}^{-1}$). In the UV spectra, Fe^{2+} binding enhanced the peak strength around 290 nm (Fig. 3). These results strongly indicate a binding reaction between Fe^{2+} and (+) taxifolin and that Fe^{2+} binding may act as one indirect pathway in the antioxidative process of (+) taxifolin.

As reported previously [36], adjacent keto or hydroxyl groups are potential targets of Fe^{2+} binding, while isolated keto-group (or hydroxyl-group) cannot bind iron. Nevertheless, the 3, 4-hydroxyl-keto moiety cannot give a planar conformation (Fig. 1b), and can barely form the planar five-membered Fe^{2+} -complex. As a dihydroflavonol, (+) taxifolin contains only two Fe^{2+} -binding sites: the 3', 4'-catechol moiety and the 4, 5-hydroxyl-keto moiety (Fig. 4) [38].

Despite several reports on the metal-binding of flavonoids [38–40] and descriptions of Na^+ interacting with flavonoids [41], studies focusing on UV-Vis spectral analyses (especially peak assignment) are lacking. To confirm the assignment of the UV-Vis peaks in Fig. 3, we investigated the Fe^{2+} -binding of catechol and dihydromyricetin (reference compounds), because in (+) taxifolin and dihydromyricetin, the possible π - π conjugation is blocked by a single 2, 3 carbon-carbon bond, and the B ring and A/C fused rings are independent of each other. Thus, the whole (+) taxifolin molecule can be divided into two spectroscopic systems: the benzoyl skeleton and the catechol moiety (Additional file 1: Figure S8). Catechol contains a similar chemical structure to the B ring of (+) taxifolin, while dihydromyricetin bears a similar chemical structure to the A/C fused rings (benzoyl skeleton) of (+) taxifolin.

Catechol- Fe^{2+} gave two similar absorbance peaks (at approximately λ_{max} 433 and 721 nm) in the Vis spectra to those of (+) taxifolin- Fe^{2+} and yielded a green solution (Additional file 1: Figure S9). By contrast, the dihydromyricetin molecule bearing a





pyrogallol moiety in the *B* ring presented a strong absorbance peak at λ_{max} 589 nm [42], and 4'-*O*-methyltaxifolin without catechol moiety only gave a UV absorbance peak at λ_{max} 289 nm (Additional file 1: Figure S10). Thus, it can be deduced that the peaks in the Vis spectra of the (+) taxifolin- Fe^{2+} complex are from the Fe^{2+} -binding reaction with catechol in the *B* ring.

With respect to the UV spectra, an enhanced strength of the UV peaks was observed in the Fe^{2+} -binding reaction with (+) taxifolin (Fig. 3a), similar to the dihydromyricetin- Fe^{2+} complex and 4'-*O*-methyltaxifolin- Fe^{2+} complex (Additional file 1: Figures S8 & S11). Dihydromyricetin and 4'-*O*-methyltaxifolin share a similar benzoyl skeleton with (+) taxifolin. Thus, the enhancement of peak around 290 nm can be attributed to the Fe^{2+} -binding reaction of the 4-hydroxyl-5-keto moiety. This assumption is further supported by the different colors between the (+) taxifolin- Fe^{2+} complex and the 4'-*O*-methyltaxifolin- Fe^{2+} complex.

Conclusion

As an effective $\bullet\text{OH}$ -scavenger, (+) taxifolin can protect bmMSCs from $\bullet\text{OH}$ -induced damage. Its $\bullet\text{OH}$ -scavenging action consists of direct and indirect antioxidant effects. The direct antioxidation occurs via multiple pathways, including ET, PCET and HAT. The indirect antioxidation involved Fe^{2+} binding. Upon binding to Fe^{2+} , the 3',4'-catechol moiety in the *B* ring gives rise to two peaks (λ_{max} 433 nm and 721 nm), and the 4-hydroxyl-5-keto of the benzoyl skeleton causes an enhanced peak intensity around 290 nm.

Additional file

Additional file 1: Figure S1. The CCK-8 assay for normal bmMSCs exposed to (+) taxifolin. **Figure S2.** Dose-response curves for (+) taxifolin $\bullet\text{OH}$ -scavenging assay based on DNA. **Figure S3.** Dose-response curves for (+) taxifolin in PTIO \bullet radical-scavenging assay and its IC_{50} values at various pH values. **Figure S4.** Dose-response curves for (+) taxifolin in the $\text{ABTS}^{\bullet+}$ radical-scavenging assay. **Figure S5.** Dose-response curves for (+) taxifolin in the Cu^{2+} -reducing assay. **Figure S6.** Dose-response curves for (+) taxifolin in the FRAP assay. **Figure S7.** Dose-response curves for (+) taxifolin in the DPPH \bullet -radical-scavenging assay. **Figure S8.** The UV-visible spectra for the 4'-*O*-methyltaxifolin- Fe^{2+} complex. **Figure S9.** The UV absorption bands of flavonoid. **Figure S10.** The UV-Vis spectra and solution colors for (+) taxifolin- Fe^{2+} and catechol- Fe^{2+} . **Figure S11.** The UV-Vis spectra for (+) taxifolin- Fe^{2+} and dihydromyricetin- Fe^{2+} . **Table S1.** The IC_{50} values listed in different units. (DOCX 789 kb)

Abbreviations

ABTS: 2,2'-azino-bis (3-ethylbenzo-thiazoline-6-sulfonic acid); bmMSCs: bone marrow-derived mesenchymal stem cells; CCK-8: Cell counting kit-8; DMEM: Dulbecco's modified Eagle's medium; ET: Electron transfer; FBS: Fetal bovine serum; Ferrozine: 3-(2-pyridyl)-5,6-bis (4-phenylsulfonic acid)-1,2,4-triazine; FRAP: Ferric ion-reducing antioxidant power; HAT: Hydrogen atom transfer; MTT: Methyl thiazolyl tetrazolium; Neocuproine: 2, 9-dimethyl-1, 10-phenanthroline; PCET: Proton-coupled electron transfer; PTIO: 2-phenyl-4,4,5, 5-tetramethylimidazole-1-oxyl-3-oxide; ROS: Reactive oxygen species; TPTZ: 2,4,6-tripyridyl triazine; Trolox: (\pm)-6-hydroxyl-2,5,7,8-tetramethylchromane-2-carboxylic acid

Acknowledgements

None.

Funding

This research was supported by the National Natural Science Foundation of China (81,503,593, 81,573,558), the Natural Science Foundation of Guangdong Province (2017A030312009, 2016A030313649), and the Guangdong Science and Technology Project (2017A050506043).

Availability of data and materials

Data are all contained within the article.

Authors' contributions

XL, GW, and DC conceived and designed the experiments; HX and ZF performed the cellular experiments and drew the Figures; LL, CL, XO, and LY conducted the DPPH-scavenging assay and ABTS⁺-scavenging assay; QJ, YX, and YL analyzed the data; XL wrote the paper. All authors read and approved the final manuscript.

Ethics approval

The institution Animal Ethics Committee in Guangzhou University of Chinese Medicine (Guangzhou, China) approved the protocols used in this study.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹School of Chinese Herbal Medicine, Guangzhou University of Chinese Medicine, Waihuang East Road No. 232, Guangzhou Higher Education Mega Center, Guangzhou 510006, China. ²Innovative Research & Development Laboratory of TCM, Guangzhou University of Chinese Medicine, Guangzhou 510006, China. ³School of Basic Medical Science, Guangzhou University of Chinese Medicine, Guangzhou 510006, China. ⁴The Research Center of Integrative Medicine, Guangzhou University of Chinese Medicine, Guangzhou 510006, China.

Received: 15 August 2017 Accepted: 15 December 2017

Published online: 27 December 2017

References

1. Alves H, Mentink A, Le B, Van Blitterswijk CA, De Boer J. Effect of antioxidant supplementation on the total yield, oxidative stress levels, and multipotency of bone marrow-derived human mesenchymal stromal cells. *Tissue Eng Part A*. 2013;19:928–37.
2. Ryan A, Denu, Peiman H. Effects of oxidative stress on mesenchymal stem cell biology. *Oxidative Med Cell Longev*. 2016;1:1–9.
3. Fang YZ, Zheng RL. Theory and application of free radical biology. Science Press, Beijing, China. 2002; pp. 541–594.
4. Mülle M, Seefried L, Genest F, Högger P. Distribution of constituents and metabolites of maritime pine bark extract (Pycnogenol®) into serum, blood cells, and synovial fluid of patients with severe osteoarthritis: a randomized controlled trial. *Nutrients*. 2017;9(5):443.
5. Stafford HA, Lester HH. Flavan-3-ol biosynthesis: the conversion of (+)-dihydromyricetin to its flavan-3,4-diol (leucodelphinidin) and to (+)-galocatechin by reductases extracted from tissue cultures of ginkgo biloba and pseudotsugamenziesii. *Plant Physiol*. 1985;78(4):791–4.
6. Dong Y, Ma XF, RJ W, Yang MY, Shou D, Li HY. Simultaneous determination of taxifolin rutin in *Pinus massoniana* lamb. By high performance liquid chromatography with diode array detection. *Chin arch of Tradit. Chin Med*. 2012;30(11):2517–9.
7. Fan TB, Liu HY, Tang Q, Liu SM. Determination the content of dihydroquercetin in pine needles by HPLC. *China Pharm*. 2009;12(8):1046–8.
8. Zhang WP, Liu W, JH F, Chai J, Liu WC, Zhen YN. Structural identification and quantitative analysis of taxifolin in *Larix olgensis* Henry Var. *Koreana Nakai*. *J Food Sci*. 2013;34(16):293–6.
9. Bayomy NA, Abdelaziz EZ, Said MA, Badawi MS, El-Bakary RH. Effect of pycnogenol and spirulina on vancomycin-induced renal cortical oxidative stress, apoptosis, and autophagy in adult male albino rat. *Can J Physiol Pharmacol*. 2016;94(8):838–48.
10. Vladimirov YA, Proskurnina EV, Demin EM, Matveeva NS, Lubitskiy OB, Novikov AA, Izmailov DY, Osipov AN, Tikhonov VP, Kagan VE. Dihydroquercetin (taxifolin) and other flavonoids as inhibitors of free radical formation at key stages of apoptosis. *Biochemistry (Mosc)*. 2009;74:301–7.
11. Maksimovich NY, Dremza IK, Troian EI, Maksimovich YN, Borodinski AN. The correcting effects of dihydroquercetin in cerebral ischemia-reperfusion injury. *Biomed Khim*. 2014;60(6):643–50.
12. Manig and an K, Manimaran D, Jayaraj RL, Elangovan N, Dhivya V, Kaphle A. Taxifolin curbs NF- κ B-mediated Wnt/ β -catenin signaling via up-regulating Nrf2 pathway in experimental colon carcinogenesis. *Biochimie*. 2015; 119: 103–112.

13. Zhang ZR, Al Zaharna M, Wong MM, Chiu SK, Cheung HY. Taxifolin enhances and rographolide-induced mitotic arrest and apoptosis in human prostate cancer cells via spindle assembly checkpoint activation. *PLoS One*. 2013; 8(1):54577.
14. Dok-Go H, Lee KH, Kim HJ, Lee EH, Lee J, Song YS, Lee YH, Jin C, Lee YS, Cho J. Neuroprotective effects of antioxidative flavonoids, quercetin, (+)-dihydroquercetin and quercetin 3-methyl ether, isolated from *Opuntia ficus-indica* var. *saboten*. *Brain Res*. 2003;965(1–2):130–6.
15. Chen DF, Li X, Xu Z, Liu X, SH D, Li H, Zhou JH, Zeng HP, Hua ZC. Hexadecanoic acid from Buzhong Yiqi decoction induces proliferation of bone marrow mesenchymal stem cells. *J Med Food*. 2010;13:967–70.
16. Li XC, Hu Q, Jiang S, Li F, Lin J, Han L, Hong Y, Lu W, Gao Y, Chen D. *Flos Chrysanthemi indicis* protects against hydroxyl-induced damages to DNA and MSCs via antioxidant mechanism: a chemistry study. *J Saudi Chem Soc*. 2015;19:454–60.
17. Wang T, Zeng G, Li X, Zeng H. *Vitro* studies on the antioxidant and protective effect of 2-substituted-8-hydroxyquinoline derivatives against H₂O₂-induced oxidative stress in BMSCs. *Chem Biol Drug Des*. 2010;75(2): 214–22.
18. Li XC, Mai W, Wang L, Han W. A hydroxyl-scavenging assay based on DNA damage *in vitro*. *Anal Biochem*. 2013; 438:29–31.
19. Li XC. 2-Phenyl-4,4,5,5-tetramethylimidazole-1-oxyl 3-oxide (PTIO•) radical-scavenging: a new and simple antioxidant assay *in vitro*. *J Agric Food Chem*. 2017;65:6288–97.
20. Anna F, Dae-Ok K, Chung SJ, Sung I, Koo, Ock KC. Comparison of ABTS/DPPH assays to measure antioxidant capacity in popular antioxidant-rich US foods. *J Food Compos Anal*. 2011;24:1043–8.
21. Li X, Chen D, Mai Y, Wen B, Wang X. Concordance between antioxidant activities *in vitro* and chemical components of radix *Astragali* (Huangqi). *Nat Prod Res*. 2012;26:1050–3.
22. Li XC, Han W, Mai W. Antioxidant activity and mechanism of tetrahydroamentoflavone *in vitro*. *Nat Prod Commun*. 2013;8:787–9.
23. Benzie IF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: the FRAP assay. *Anal Biochem*. 1996;239:70–6.
24. Kim A, Nam YJ, Lee CS. Taxifolin reduces the cholesterol oxidation product-induced neuronal apoptosis by suppressing the Akt and NF- κ B activation-mediated cell death. *Brain Res Bull*. 2017;134:63–71.
25. Macáková K, Mladěnka P, Filipický T, Říha M, Jahodář L, Trejtnar F, Bovicelli P, Proietti Silvestri I, Hrdina R, Saso L. Iron reduction potentiates hydroxyl radical formation only in flavonols. *Food Chem*. 2012;135(4):2584–92.
26. Moridani MY, Pourahmad J, Bui H, Siraki A, O'Brien PJ. Dietary flavonoid iron complexes as cytoprotective superoxide radical scavengers. *Free Radic Biol Med*. 2003;34(2):243–53.
27. Rajnochová Svobodová A, Ryšavá A, Psotová M, Kosina P, Zálešák B, Ulrichová J, Vostálová J. The phototoxic potential of the flavonoids, taxifolin and quercetin. *Photochem Photobiol*. 2017;93(5):1240–7.
28. Kang JT, Moon JH, Choi JY, Park SJ, Kim SJ, Saadeldin IM, Lee BC. Effect of antioxidant flavonoids (quercetin and taxifolin) on *in vitro* maturation of porcine oocytes. *Asian-Australas J Anim Sci*. 2016;29(3):352–8.
29. Wang T, Li X, Wu J, Huang Y, Wei G, Chen D. Mechanistic chemistry of extraordinary capacity of salivianolic acid B on oxidatively-damaged mesenchymal stem cells. *J Chin Chem Soc*. 2016;63(11):924–9.
30. Wang G, Li X, Zeng H. Synthesis antioxidant activity of (E)-9-p-tolyl-3-[2-(8-hydroxy-quinol-2-yl)vinyl]-carbazole and (E)-9-(p-Anisyl)-3-[2-(8-hydroxy-quinol-2-yl)vinyl]-carbazole and their induction proliferation of mesenchymal stem cells. *Acta Chimica Sinica*. 2009;67(9):974–82.
31. Li X, Wei G, Wang X, Liu DH, Deng RD, Li H, Zhou JH, Li YW, Zeng HP, Chen DF. Targeting of the sonic hedgehog pathway by Atractylenolides promotes Chondrogenic differentiation of mesenchymal stem cells. *Biol Pharm Bull*. 2012;35:1328–35.
32. Lin J, Li X, Chen L, Lu W, Chen X, Han L, Chen D. Protective effect against hydroxyl radical-induced DNA damage and antioxidant mechanism of [6]-gingerol: a chemical study. *B Korean. Chem Soc*. 2014;35(6):1633–8.
33. Goldstein S, Russo A, Samuni A. Reactions of PTIO• and carboxy-PTIO• with •NO, •NO₂, and O₂. *J Biol Chem*. 2003; 278:50949–55.
34. Apak R, Özyürek M, Güçlü K, Çapanoğlu E. Antioxidant activity/capacity measurement. 2. Hydrogen atom transfer (HAT)-based, mixed-mode (electron transfer (ET)/HAT), and lipid peroxidation assays. *J Agric Food Chem*. 2016;64: 1028–45.
35. Gülçin İ. Antioxidant activity of food constituents: an overview. *Arch Toxicol*. 2012;86:345–91.
36. Sugihara N, Arakawa T, Ohnishi M, Furuno K. Anti- and pro-oxidative effects of flavonoids on metal-induced lipid hydroperoxide-dependent lipid peroxidation in cultured hepatocytes loaded with α -linolenic acid. *Free Radic Biol Med*. 1999;27(11–12):1313–23.
37. Jiang Q, Li X, Tian Y, Lin Q, Xie H, Lu W, Chi Y, Chen D. Lyophilized aqueous extracts of *Mori Fructus* and *Mori Ramulus* protect mesenchymal stem cells from •OH-treated damage: bioassay and antioxidant mechanism. *BMC Complem Altern M*. 2017;17:242.
38. Mladěnka P, Macáková K, Filipický T, Zatloukalová L, Jahodář L, Bovicelli P, Silvestri IP, Hrdina R, Saso L. *Vitro* analysis of iron binding activity of flavonoids. *J Inorg Biochem*. 2011;105:693–701.
39. Mira L, Fernandez MT, Santos M, Rocha R, Florêncio MH, Jennings KR. Interactions of flavonoids with iron and copper ions: a mechanism for their antioxidant activity. *Free Radic Res*. 2002;36:1199–208.
40. Li X, Gao Y, Li F, Liang A, Xu Z, Bai Y, Mai W, Han L, Chen D. Maclurin protects against hydroxyl radical-induced damages to mesenchymal stem cells: antioxidant evaluation and mechanistic insight. *Chem Biol Interact*. 2014; 219:221–8.
41. Xiao CH. *Chemistry of Chinese Materia Medica*. 3rd ed. Shanghai, China: Shanghai Scientific & Technical Publishers; 1997. p. 289–99.
42. Li XC, Liu JJ, Lin J, Wang TT, Huang JY, Lin YQ, Chen DF. Protective effects of dihydromyricetin against •OH-induced mesenchymal stem cells damage and mechanistic chemistry. *Molecules*. 2016;21:604.