



## NOTE

Immunology

# The enhancement of Th1 immune response by anti-PD-L1 antibody in cattle infected with *Mycobacterium avium* subsp. *paratuberculosis*

Yamato SAJIKI<sup>1)</sup>, Satoru KONNAI<sup>1,2)\*</sup>, Reiko NAGATA<sup>3)</sup>, Satoko KAWAJI<sup>3)</sup>, Hayato NAKAMURA<sup>1)</sup>, Sotaro FUJISAWA<sup>1)</sup>, Tomohiro OKAGAWA<sup>2)</sup>, Naoya MAEKAWA<sup>2)</sup>, Yukinari KATO<sup>4,5)</sup>, Yasuhiko SUZUKI<sup>2,6)</sup>, Shiro MURATA<sup>1,2)</sup>, Yasuyuki MORI<sup>3)</sup> and Kazuhiko OHASHI<sup>1,2)</sup>

<sup>1)</sup>Department of Disease Control, Faculty of Veterinary Medicine, Hokkaido University, Sapporo, Hokkaido 060-0818, Japan

<sup>2)</sup>Department of Advanced Pharmaceutics, Faculty of Veterinary Medicine, Hokkaido University, Sapporo, Hokkaido 060-0818, Japan

<sup>3)</sup>Bacterial and Parasitic Disease Research Division, National Institute of Animal Health, Tsukuba, Ibaraki 305-0856, Japan

<sup>4)</sup>Department of Antibody Drug Development, Tohoku University Graduate School of Medicine, 2-1 Seiryomachi, Aoba-ku, Sendai, Miyagi 980-8575, Japan

<sup>5)</sup>New Industry Creation Hatchery Center, Tohoku University, 2-1 Seiryomachi, Aoba-ku, Sendai, Miyagi 980-8575, Japan

<sup>6)</sup>Division of Bioresources, Research Center for Zoonosis Control, Hokkaido University, Sapporo, Hokkaido 001-0019, Japan

*J. Vet. Med. Sci.*

83(2): 162–166, 2021

doi: 10.1292/jvms.20-0590

Received: 6 October 2020

Accepted: 24 November 2020

Advanced Epub:

7 December 2020

**ABSTRACT.** Johne's disease, caused by *Mycobacterium avium* subsp. *paratuberculosis* (MAP), is a chronic enteritis of ruminants. Previous studies have shown that programmed death-ligand 1 (PD-L1) is associated with the disease progression, and PD-L1 blockade activates MAP-specific Th1 responses *in vitro*. Here, we performed anti-PD-L1 antibody administration using 2 MAP-infected cattle at the late subclinical stage of infection. After administration, bacterial shedding was reduced or maintained at a low level. Additionally, MAP-specific Th1 cytokine production was upregulated, and *CD69* expression was increased in T cells. Collectively, the treatment has a potential as a novel control method against Johne's disease.

**KEY WORDS:** cattle, immunotherapy, Johne's disease, programmed death-ligand 1, Th1 response

Johne's disease (paratuberculosis) is a chronic enteritis of ruminants, and is caused by the bacteria *Mycobacterium avium* subsp. *paratuberculosis* (MAP). The clinical signs of Johne's disease include chronic diarrhea, severe weight loss, reduced milk production, and mortality [17]. Johne's disease is endemic in many countries, including Japan [13]. In an early stage of infection, MAP induces strong Th1 responses that cause the activation of macrophages to kill intracellular mycobacteria [4, 21, 22]. Th1 cytokines, such as interferon (IFN)- $\gamma$  and tumor necrosis factor (TNF)- $\alpha$ , are important for enhancing bactericidal activity of macrophages with the production of reactive oxygen and nitrogen species, T cell activation, and dendritic cell maturation [8, 12]. During the late subclinical stage, the Th1 response declines, which allows bacterial growth and progression to clinical disease [1, 2, 24]. Therefore, the Th1 response is essential for the prevention of the disease progression.

Programmed death (PD)-1 is one of the immunoinhibitory receptors expressed on T cells, and induces immunosuppression by binding to PD-ligand 1 (PD-L1) [11]. In chronic infections, the upregulation of PD-1 and PD-L1 expression is involved in the exhaustion of antigen-specific T cells which contributes to the disease progression [11, 25]. During human tuberculosis that is caused by *Mycobacterium tuberculosis*, the PD-1/PD-L1 pathway inhibits effector function of T cells [9], and PD-1 expression on *M. tuberculosis*-specific CD4<sup>+</sup> T cells is associated with bacterial loads [5]. Previous studies have shown that the PD-1/PD-L1 pathway is involved in the suppression of Th1 responses in cattle infected with MAP [16]. The blockade of PD-L1 using a specific antibody (Ab) increases MAP-specific Th1 immune responses *in vitro* [18]. Thus, the PD-1/PD-L1 pathway is considered to have

\*Correspondence to: Konnai, S.: [konnai@vetmed.hokudai.ac.jp](mailto:konnai@vetmed.hokudai.ac.jp)

©2021 The Japanese Society of Veterinary Science



This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License. (CC-BY-NC-ND 4.0: <https://creativecommons.org/licenses/by-nc-nd/4.0/>)

a therapeutic potential for Johne's disease. In addition, previous studies have demonstrated that anti-PD-L1 rat-bovine chimeric antibody (chAb) has therapeutic effects against other chronic bovine infections, such as bovine leukemia virus (BLV) infection and *Mycoplasma bovis* infection [7, 15, 19]. However, there is no report which evaluates the function of PD-L1 blockade in MAP-infected animals. Therefore, in this study, we performed the administration of anti-PD-L1 chAb using 2 MAP experimentally-infected cattle to examine the responses to the antibody administration by immunological and bacteriological analyses.


For the experimental infection of MAP, 2 male Holstein calves, animals #80 (3 weeks of age) and #99 (a week of age), were orally inoculated with intestinal tissue homogenate from an infected cow containing MAP (#80:  $1.36 \times 10^8$  CFU; #99:  $2.50 \times 10^8$  CFU) which was measured by using Middlebrook 7H10 agar-based slants as described in a previous paper [10]. Both animals were sourced from farms with no history of paratuberculosis and confirmed negative by fecal quantitative polymerase chain reaction (qPCR) targeting MAP-specific gene *IS900* as described previously [10] and by Pourquier ELISA (Institut Pourquier, Montpellier, France) prior to inoculation with MAP. Animals #80 (770 kg, 212 weeks post-infection) and #99 (320 kg, 47 weeks post-infection) were intravenously administered with 2 mg/kg of the purified anti-PD-L1 chAb (Boch4G12) [15] a time and three times at 2 week-intervals, respectively. Both animals were kept in a biosafety level 2 animal facility at the National Institute of Animal Health, Tsukuba, Japan. All experiments using these animals were approved by the National Institute of Animal Health Ethics Committee (approval No. 17-077-2 and 18-077). After the experimental infection, we collected blood and fecal samples at intervals of 2–4 weeks, and monitored IFN- $\gamma$  production responded to Johnin purified protein derivative (J-PPD) by whole-blood cultures as described previously [16], the serum levels of Abs against MAP by Pourquier ELISA, and fecal shedding of MAP by qPCR. To examine the effects of anti-PD-L1 Ab in MAP-infected cattle, blood samples were collected from animal #80 on the day of administration (day 0), and on several points after administration (days 1, 3, 8, 15, 29, 43, 57, and 85). Blood samples on day 0 were obtained before administration. Blood samples were collected from animal #99 on days 0, 7, 14, 21, 28, 42, 56, 70, 84, 98, and 112. Blood samples on days 0, 14, and 28 were obtained before administration. Peripheral blood mononuclear cells (PBMCs) were purified from blood samples using density gradient centrifugation on Percoll (GE Healthcare, Little Chalfont, UK), and cultured with 2  $\mu\text{g}/\text{ml}$  of J-PPD or 20  $\mu\text{g}/\text{ml}$  of concanavalin A (Con A; Sigma-Aldrich, St. Louis, MO, USA). Phosphate buffered saline (PBS) and PPD from *Mycobacterium bovis* BCG strain (B-PPD) were used as a negative control and a control antigen, respectively. After 6 days, collected culture supernatants were assayed for IFN- $\gamma$  and TNF- $\alpha$  by Bovine IFN- $\gamma$  ELISA Development Kit (Mabtech, Nacka Strand, Sweden) and Bovine TNF alpha Do-It Yourself ELISA (Kingfisher Biotech, St. Paul, MN, USA), respectively. Additionally, the isolation of T cells ( $\text{CD}3^+$  cells) from PBMCs was performed according to a previous report [15], and the purity of  $\text{CD}3^+$  cells (>95%) was confirmed using FACS Verse (BD Biosciences, San Jose, CA, USA). To quantitate gene expression levels of *IFN- $\gamma$* , *TNF- $\alpha$* , and *CD69*, an activation marker of lymphocytes, qPCR was conducted as described previously [18]. Primers for *IFN- $\gamma$* , *TNF- $\alpha$* , and *ACTB* were shown in a previous report [18]. Primers for *CD69* were 5'-ATA GCT CTC GTT GCT CTA TCA GTG-3' and 5'-CCT TGT GTC CAA TCC AAT CA-3'.

In animal #80, J-PPD-specific IFN- $\gamma$  response peaked during the acute phase of infection (Fig. 1A). This response was gradually suppressed from approximately 60 weeks post-inoculation, and was maintained at a low level until the time of administration (Fig. 1A). In contrast, the serum levels of Abs against MAP were increased from 160 weeks post-infection (Fig. 1B). The tentative shedding of MAP was observed in feces of animal #80 from 4 to 16 weeks post-infection (Fig. 1C). Fecal shedding was not detected between 18–78 weeks, and then the MAP DNA quantity gradually increased from 80 weeks post-infection (Fig. 1C). In animal #99, J-PPD-specific IFN- $\gamma$  response peaked at 23 weeks after the experimental infection, and was maintained at a high level (Fig. 1A). The serum levels of Abs against MAP were increased and then turned positive at 46 weeks post-infection, a week before the administration (Fig. 1B). We observed bacterial shedding intermittently until the time of administration (Fig. 1C). Although both animals #80 and #99 showed the bacterial shedding, these animals did not show clinical symptoms such as diarrhea. The serum levels of Abs against MAP was increasing before anti-PD-L1 Ab administration. A previous report has described that detectable levels of Abs against MAP return in mid- to late-stage subclinical infections [3]. From these data, we concluded that animals #80 and #99 were both in the late subclinical stage at the time of administration.

To evaluate anti-bacterial effects of Boch4G12 in MAP-infected cattle, fecal shedding of MAP was monitored by qPCR for 60 weeks for animal #80 and 34 weeks for #99 after administration. The bacterial loads in feces from #80 and #99 were maintained at a low level after administration (Fig. 1C). Remarkably, in animal #99, MAP DNA was not detected in feces at 30 and 34 weeks after Boch4G12 administration (correspondence to 77 and 81 weeks-post infection, respectively) (Fig. 1C). A previous report has shown that persistent shedding patterns related on ELISA-positive samples rarely reverse to negativity [14]. In addition, cattle detected more than  $1.0 \times 10^{-2}$  pg MAP DNA in feces are associated with progressing to severe disease [23]. These data suggest that anti-PD-L1 Ab treatment has a possibility to regulate the bacterial shedding in MAP-infected cattle.

We then examined the effects of Boch4G12 on Th1 responses *in vitro*. IFN- $\gamma$  and TNF- $\alpha$  production from PBMCs was significantly increased in the presence of J-PPD and Con A (Fig. 2A and 2B). TNF- $\alpha$  was not detected from culture supernatants of groups stimulated with PBS and B-PPD. In addition, the expression levels of *IFN- $\gamma$* , *TNF- $\alpha$* , and *CD69* in T cells of #80 were significantly upregulated after Boch4G12 administration (Fig. 2C). *CD69* upregulation in T cells was also observed in animal #99 (Fig. 2C). Collectively, these results demonstrated that treatment with anti-PD-L1 chAb activated Th1 responses in both animals, suggesting a therapeutic potential for Johne's disease.


In this study, the effects of anti-PD-L1 Ab treatment on bacterial shedding were different between #80 and #99. Animal #80 was administrated with anti-PD-L1 Ab a time, whereas animal #99 was administrated three times. Before anti-PD-L1 Ab administration, IFN- $\gamma$  production responded to J-PPD in animal #80 had been suppressed for more than 100 weeks, whereas IFN- $\gamma$  production in animal #99 had not been suppressed. These differences might be factors to determine the effect of anti-PD-L1 Ab treatment on fecal



**Fig. 1.** Interferon (IFN)- $\gamma$  responses in blood, antibody (Ab) levels in sera, and fecal shedding of *Mycobacterium avium* subsp. *paratuberculosis* (MAP). (A) Whole-blood cells were cultured with Johnin purified protein derivative for 24 hr. IFN- $\gamma$  concentrations in culture supernatants were determined by ELISA. (B) The serum levels of Abs against MAP were examined by ELISA. Results were indicated as percentage S/P, and the dotted lines indicated cutoff as recommended by the manufacturer. (C) MAP DNA quantity in feces detected by qPCR. (A–C) Arrowheads show the points of anti-programmed death-ligand 1 Ab administration.

shedding. Further clinical studies using a number of MAP-infected animals are required to examine the influence of these factors, such as the experimental conditions (administration time, interval, and dosage) and examined animals (disease stage, response to J-PPD, and age), on the clinical efficacy of anti-PD-L1 Ab.

Th1 responses, especially IFN- $\gamma$  production, are considered to be essential for the prevention of the disease progression. Our previous and present studies have shown that treatment with anti-PD-L1 Ab enhanced MAP-specific Th1 responses both *in vitro* and *in vivo* [18]. Therefore, the observed effect of anti-PD-L1 Ab on the bacterial shedding is presumably due to the activation of Th1 responses. Additional experiments are required to elucidate the underlying mechanism of the anti-bacterial effect of anti-PD-L1 Ab on MAP-infected cattle. Further, recent studies have described that the combined treatment of anti-PD-L1 Ab with other medicines has a potential to enhance therapeutic effects of PD-1/PD-L1 blockade in several chronic bovine infections [6, 7, 18–20]. A previous study on Johne's disease revealed that the dual blockade of the PD-1/PD-L1 pathway and prostaglandin E<sub>2</sub> production enhanced MAP-specific Th1 responses *in vitro* [18]. Hence, the combination with other medicines could be a strategy to enhance the anti-bacterial effect of anti-PD-L1 Ab in cattle infected with MAP.



**Fig. 2.** Activation of *Mycobacterium avium* subsp. *paratuberculosis* (MAP) specific Th1 responses by anti-programmed death-ligand 1 chimeric antibody. (A and B) PBMCs were cultured with Phosphate buffered saline (PBS), purified protein derivative from *Mycobacterium bovis* BCG strain (B-PPD), Johnin-PPD (J-PPD), or concanavalin A (Con A) for 6 days. Interferon (IFN)- $\gamma$  (A) and tumor necrosis factor (TNF)- $\alpha$  (B) concentrations in culture supernatants were determined by ELISA. (A and B) The symbols represent the means of three independent culture wells, and white symbols show  $P < 0.05$  compared with the value on day 0. (C) Gene expression levels of *IFN- $\gamma$* , *TNF- $\alpha$* , and *CD69* in  $CD3^+$  cells were quantitated by qPCR. The symbols represent the means of three independent experiments, and white symbols show  $P < 0.05$  compared with the value on day 0. (A–C) Statistical significance was determined by the Dunnett's test.

In conclusion, we showed a potential of anti-PD-L1 Ab treatment for the regulation of bacterial shedding. Additionally, anti-PD-L1 Ab treatment also activated MAP-specific Th1 cytokine production in MAP-infected cattle. To our best knowledge, this is the first study which shows immune activating effects of the PD-L1 blockade in cattle infected with MAP. Although the number of experimentally-infected cattle used in this study was limited, the observations in the present study could play an important role to establish a novel therapeutic strategy against Johne's disease.

**CONFLICT OF INTEREST.** The authors declare that they have no competing interests.

**ACKNOWLEDGMENTS.** This work was supported by JSPS KAKENHI grant number 19KK0172 [to S.K.], grants from the Project of the NARO, Bio-oriented Technology Research Advancement Institution (Research Program on Development of Innovative Technology 26058 BC [to S.K.] and Special Scheme Project on Regional Developing Strategy, Grant 16817557 [to S.K.]), regulatory



research projects for food safety, animal health and plant protection (JPJ008617.17935709) funded by the Ministry of Agriculture, Forestry and Fisheries of Japan, and AMED under grant number JP20am0101078 [to Y.K.]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. We are grateful to Dr. Hideyuki Takahashi and Dr. Tomio Ibayashi for valuable advice and discussions.

## REFERENCES

- Bassey, E. O. and Collins, M. T. 1997. Study of T-lymphocyte subsets of healthy and *Mycobacterium avium* subsp. *paratuberculosis*-infected cattle. *Infect. Immun.* **65**: 4869–4872. [Medline] [CrossRef]
- Burrells, C., Clarke, C. J., Colston, A., Kay, J. M., Porter, J., Little, D. and Sharp, J. M. 1999. Interferon-gamma and interleukin-2 release by lymphocytes derived from the blood, mesenteric lymph nodes and intestines of normal sheep and those affected with paratuberculosis (Johne's disease). *Vet. Immunol. Immunopathol.* **68**: 139–148. [Medline] [CrossRef]
- Coussens, P. M. 2001. *Mycobacterium paratuberculosis* and the bovine immune system. *Anim. Health Res. Rev.* **2**: 141–161. [Medline] [CrossRef]
- Coussens, P. M. 2004. Model for immune responses to *Mycobacterium avium* subspecies *paratuberculosis* in cattle. *Infect. Immun.* **72**: 3089–3096. [Medline] [CrossRef]
- Day, C. L., Abrahams, D. A., Bunjun, R., Stone, L., de Kock, M., Walzl, G., Wilkinson, R. J., Burgers, W. A. and Hanekom, W. A. 2018. PD-1 expression on *Mycobacterium tuberculosis*-specific CD4 T cells is associated with bacterial load in human tuberculosis. *Front. Immunol.* **9**: 1995. [Medline] [CrossRef]
- Goto, S., Konnai, S., Hirano, Y., Kohara, J., Okagawa, T., Maekawa, N., Sajiki, Y., Watari, K., Minato, E., Kobayashi, A., Gondaira, S., Higuchi, H., Koiwa, M., Tajima, M., Taguchi, E., Uemura, R., Yamada, S., Kaneko, M. K., Kato, Y., Yamamoto, K., Toda, M., Suzuki, Y., Murata, S. and Ohashi, K. 2020. Upregulation of PD-L1 expression by prostaglandin E<sub>2</sub> and the enhancement of IFN- $\gamma$  by anti-PD-L1 antibody combined with a COX-2 inhibitor in *Mycoplasma bovis* infection. *Front. Vet. Sci.* **7**: 12. [Medline] [CrossRef]
- Goto, S., Konnai, S., Hirano, Y., Kohara, J., Okagawa, T., Maekawa, N., Sajiki, Y., Watari, K., Minato, E., Kobayashi, A., Gondaira, S., Higuchi, H., Koiwa, M., Tajima, M., Taguchi, E., Uemura, R., Yamada, S., Kaneko, M. K., Kato, Y., Yamamoto, K., Toda, M., Suzuki, Y., Murata, S. and Ohashi, K. 2020. Clinical efficacy of the combined treatment of anti-PD-L1 rat-bovine chimeric antibody with a COX-2 inhibitor in calves infected with *Mycoplasma bovis*. *Jpn. J. Vet. Res.* **68**: 77–90.
- Hussain, T., Shah, S. Z., Zhao, D., Sreevatsan, S. and Zhou, X. 2016. The role of IL-10 in *Mycobacterium avium* subsp. *paratuberculosis* infection. *Cell Commun. Signal.* **14**: 29. [Medline] [CrossRef]
- Jurado, J. O., Alvarez, I. B., Pasquinelli, V., Martínez, G. J., Quiroga, M. F., Abbate, E., Musella, R. M., Chuluyan, H. E. and García, V. E. 2008. Programmed death (PD)-1:PD-ligand 1/PD-ligand 2 pathway inhibits T cell effector functions during human tuberculosis. *J. Immunol.* **181**: 116–125. [Medline] [CrossRef]
- Kawaji, S., Nagata, R. and Mori, Y. 2014. Detection and confirmation of *Mycobacterium avium* subsp. *paratuberculosis* in direct quantitative PCR positive fecal samples by the manual fluorescent MGIT culture system. *J. Vet. Med. Sci.* **76**: 65–72. [Medline] [CrossRef]
- Keir, M. E., Butte, M. J., Freeman, G. J. and Sharpe, A. H. 2008. PD-1 and its ligands in tolerance and immunity. *Annu. Rev. Immunol.* **26**: 677–704. [Medline] [CrossRef]
- Khalifeh, M. S., Al-Majali, A. M. and Stabel, J. R. 2009. Role of nitric oxide production in dairy cows naturally infected with *Mycobacterium avium* subsp. *paratuberculosis*. *Vet. Immunol. Immunopathol.* **131**: 97–104. [Medline] [CrossRef]
- Li, L., Katani, R., Schilling, M. and Kapur, V. 2016. Molecular Epidemiology of *Mycobacterium avium* subsp. *paratuberculosis* on Dairy Farms. *Annu. Rev. Anim. Biosci.* **4**: 155–176. [Medline] [CrossRef]
- Navarro-Gonzalez, N., Fourichon, C., Blanquefort, P., Delafosse, A., Joly, A., Ngwa-Mbot, D., Biet, F., Boichard, D., Schibler, L., Journaux, L., Meens, E. and Guatteo, R. 2019. Longitudinal study of *Mycobacterium avium* ssp. *paratuberculosis* fecal shedding patterns and concurrent serological patterns in naturally infected dairy cattle. *J. Dairy Sci.* **102**: 9117–9137. [Medline] [CrossRef]
- Nishimori, A., Konnai, S., Okagawa, T., Maekawa, N., Ikebuchi, R., Goto, S., Sajiki, Y., Suzuki, Y., Kohara, J., Ogasawara, S., Kato, Y., Murata, S. and Ohashi, K. 2017. *In vitro* and *in vivo* antiviral activity of an anti-programmed death-ligand 1 (PD-L1) rat-bovine chimeric antibody against bovine leukemia virus infection. *PLoS One* **12**: e0174916. [Medline] [CrossRef]
- Okagawa, T., Konnai, S., Nishimori, A., Ikebuchi, R., Mizorogi, S., Nagata, R., Kawaji, S., Tanaka, S., Kagawa, Y., Murata, S., Mori, Y. and Ohashi, K. 2015. Bovine immunoinhibitory receptors contribute to suppression of *Mycobacterium avium* subsp. *paratuberculosis*-specific T-cell responses. *Infect. Immun.* **84**: 77–89. [Medline] [CrossRef]
- Rathnaiah, G., Zinniel, D. K., Bannantine, J. P., Stabel, J. R., Gröhn, Y. T., Collins, M. T. and Barletta, R. G. 2017. Pathogenesis, molecular genetics, and genomics of *Mycobacterium avium* subsp. *paratuberculosis*, the etiologic agent of Johne's disease. *Front. Vet. Sci.* **4**: 187. [Medline] [CrossRef]
- Sajiki, Y., Konnai, S., Okagawa, T., Nishimori, A., Maekawa, N., Goto, S., Ikebuchi, R., Nagata, R., Kawaji, S., Kagawa, Y., Yamada, S., Kato, Y., Nakajima, C., Suzuki, Y., Murata, S., Mori, Y. and Ohashi, K. 2018. Prostaglandin E<sub>2</sub> induction suppresses the Th1 immune responses in cattle with Johne's disease. *Infect. Immun.* **86**: e00910–e00917. [Medline] [CrossRef]
- Sajiki, Y., Konnai, S., Okagawa, T., Nishimori, A., Maekawa, N., Goto, S., Watari, K., Minato, E., Kobayashi, A., Kohara, J., Yamada, S., Kaneko, M. K., Kato, Y., Takahashi, H., Terasaki, N., Takeda, A., Yamamoto, K., Toda, M., Suzuki, Y., Murata, S. and Ohashi, K. 2019. Prostaglandin E<sub>2</sub>-induced immune exhaustion and enhancement of antiviral effects by anti-PD-L1 antibody combined with COX-2 inhibitor in bovine leukemia virus infection. *J. Immunol.* **203**: 1313–1324. [Medline] [CrossRef]
- Sajiki, Y., Konnai, S., Okagawa, T., Maekawa, N., Nakamura, H., Kato, Y., Suzuki, Y., Murata, S. and Ohashi, K. 2021. A TLR7 agonist activates bovine Th1 response and exerts antiviral activity against bovine leukemia virus. *Dev. Comp. Immunol.* **114**: 103847. [Medline] [CrossRef]
- Sohal, J. S., Singh, S. V., Tyagi, P., Subhodh, S., Singh, P. K., Singh, A. V., Narayanasamy, K., Sheoran, N. and Singh Sandhu, K. 2008. Immunology of mycobacterial infections: with special reference to *Mycobacterium avium* subspecies *paratuberculosis*. *Immunobiology* **213**: 585–598. [Medline] [CrossRef]
- Stabel, J. R. 2006. Host responses to *Mycobacterium avium* subsp. *paratuberculosis*: a complex arsenal. *Anim. Health Res. Rev.* **7**: 61–70. [Medline] [CrossRef]
- Taniguchi, Y., Sakakibara, S. I., Fujihara, M., Yagi, A. and Fujiyoshi, S. 2020. The association between detection of *Mycobacterium avium* subsp. *paratuberculosis* DNA in feces and histopathological classification. *J. Vet. Med. Sci.* **82**: 541–545. [Medline] [CrossRef]
- Weiss, D. J., Evanson, O. A. and Souza, C. D. 2006. Mucosal immune response in cattle with subclinical Johne's disease. *Vet. Pathol.* **43**: 127–135. [Medline] [CrossRef]
- Wherry, E. J. 2011. T cell exhaustion. *Nat. Immunol.* **12**: 492–499. [Medline] [CrossRef]