



POSTER PRESENTATION

Open Access

Familial Mediterranean Fever in older children

Semanur Özdel¹, Zeynep Birsin Özçakar^{1*}, Seda Sahin², Mesiha Ekim¹, Atilla Elhan³, Fatos Yalcinkaya¹

From 21st European Pediatric Rheumatology (PReS) Congress
Belgrade, Serbia. 17-21 September 2014

Introduction

Familial Mediterranean Fever (FMF) is an autosomal recessive disease, characterised by recurrent, self limited attacks of fever with serositis.

Objectives

The aim of this study was to compare the demographic, clinical and genetic features of FMF patients who had late onset disease to those with earlier onset during childhood period.

Methods

Files of patients who had been seen in our department (during routine follow-up visits) between January 2013 and January 2014 were retrospectively evaluated. Patients were divided into two groups according to age of disease onset (Group I: ≤ 8 years of age; Group II: > 8 years of age).

Results

The study group comprised 317 FMF patients (170 females, 147 males) with a mean age of 12.2 ± 5.7 years. There were 267 patients (84.3%) in group I and 50 patients (15.7%) in Group II. Median attack frequency before colchicine therapy was 24/year in Group I and 12/year in Group II ($p < 0.05$). Although the frequency of majority of the clinical features did not differ between the groups, fever was seen less frequently in Group II patients ($p = 0.003$). M694V homozygosity was also less frequently detected in group II patients ($p = 0.022$). Median disease severity scores and final colchicine dosages were lower in Group II ($p < 0.001$; $p = 0.003$). Median delay in diagnosis was 24 months in Group I and 12 months in Group II ($p = 0.002$).

Conclusion

Only a small number of FMF patients had disease onset at older ages in childhood period. It seems that FMF patients

with late onset disease have milder illness. However, more readily expression of their clinical findings in older ages yields earlier diagnosis in this group.

Disclosure of interest

None declared.

Authors' details

¹Pediatric Rheumatology, Ankara University, Ankara, Turkey. ²Pediatrics, Ankara University, Ankara, Turkey. ³Biostatistics, Ankara University, Ankara, Turkey.

Published: 17 September 2014

doi:10.1186/1546-0096-12-S1-P254

Cite this article as: Özdel et al.: Familial Mediterranean Fever in older children. *Pediatric Rheumatology* 2014 **12**(Suppl 1):P254.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Pediatric Rheumatology, Ankara University, Ankara, Turkey
Full list of author information is available at the end of the article