

POSTER PRESENTATION

Open Access

Humoral immunogenicity of ChAd63_MVA ME-TRAP vaccination in African infants and children

Georgina Bowyer^{1*}, Muhammed O Afolabi², Carly M Bliss¹, Alfred B Tiono³, Abdoulie Drammeh², Issa Nébié³, Ya Jankey Jagne², Jean Baptiste Yaro³, Egeruan B Imoukhuede⁴, Katie L Flanagan², Beate Kampman², Nicola Viebig⁵, Sodiomon B Sirima³, Kalifa Bojang², Adrian VS Hill^{1,4}, Katie J Ewer¹

From Challenges in malaria research: Core science and innovation
Oxford, UK. 22-24 September 2014

Background

The only malaria vaccine to have been tested in Phase III clinical vaccine trials [1] induces protection which has been associated with high titres of antibodies against sporozoites. Efficacy against clinical malaria in infants of 5-17 months and 6-12 weeks was 55.8% and 31.3% respectively. A successful malaria vaccine will likely need to induce both cellular and humoral immunity in order to achieve a deployable level of efficacy in the target age groups. Prime-boost immunisation with viral vectors ChAd63 and MVA, both encoding ME-TRAP, has been shown to induce high T cell responses and modest antibody responses to the pre-erythrocytic malaria antigen TRAP. A Phase II study with this vaccine regime in malaria-naïve adults showed significant efficacy, which correlated with frequency of monofunctional CD8⁺ T cells secreting IFN γ . TRAP antibody titres were modest and did not correlate with protection [2]. In contrast to this, high titres of vaccine-induced anti-TRAP antibodies are measured in infants in malaria-endemic settings.

Materials and methods

Antibody titres were measured in 138 malaria-exposed children vaccinated with ChAd63 MVA ME-TRAP in three Phase I studies in The Gambia and Burkina Faso. Age groups at first immunisation were 2-6 years, 5-12 months and 10 weeks in The Gambia and 5-17 months in Burkina Faso. Avidity and isotype profiles were also analyzed.

Results

Antibody responses to TRAP were significantly higher in 10 week old and 5-12 month old infants in The Gambia

and 5-17 month old infants in Burkina Faso compared to 2-6 year old children and adults in The Gambia and malaria-naïve UK adults. IgG isotype responses were predominantly IgG1 and IgG3 and we also detected IgA and IgM. TRAP-specific IgG avidity was significantly higher in Burkinabe infants aged 5-17 months and Gambian infants aged 5-12 months compared to Gambian adults and 2-6 year old children. TRAP-specific IgG1 avidity significantly correlated with age at vaccination in 5-17 month old Burkinabe infants and was significantly higher than in 10 week old Gambian infants. Functional activity of anti-TRAP antibodies will be analysed *in vitro* using a sporozoite invasion inhibition assay.

Conclusions

We demonstrate excellent humoral immunogenicity in key target populations vaccinated with a pre-erythrocytic malaria vaccination regime, exceeding that seen in UK malaria-naïve adults.

Authors' details

¹Jenner Institute Laboratories, University of Oxford, Oxford, UK. ²Medical Research Council Unit, Fajara, Gambia. ³Centre National de Recherche et de Formation sur le Paludisme, Ouagadougou, Burkina Faso. ⁴Centre for Clinical Vaccinology and Tropical Medicine, Oxford, UK. ⁵European Vaccine Initiative, Heidelberg, Germany.

Published: 22 September 2014

References

1. Vansadia P, et al: A Phase 3 Trial of RTS,S/AS01 Malaria Vaccine in African Infants. *N Engl J Med* 2012, **367**:2284-2295.
2. Ewer K, O'Hara G: Protective CD8⁺ T-cell immunity to human malaria induced by chimpanzee adenovirus-MVA immunisation. *Nat Commun* 2013, **4**.

doi:10.1186/1475-2875-13-S1-P16

Cite this article as: Bowyer et al: Humoral immunogenicity of ChAd63_MVA ME-TRAP vaccination in African infants and children. *Malaria Journal* 2014 **13**(Suppl 1):P16.

¹Jenner Institute Laboratories, University of Oxford, Oxford, UK
Full list of author information is available at the end of the article