

347. SARS-CoV-2 and Acute Otitis Media in Children: A Case Series

Holly M. Frost, MD¹; Theresia Sebastian, MD, MPH²; Amy Keith, MPH²; Melanie Kurtz, BS²; Samuel R. Dominguez, MD, PhD³; Samuel R. Dominguez, MD, PhD³; Sarah Parker, MD⁴; Timothy C. Jenkins, MD⁵; ¹Denver Health and Hospital Authority, University of Colorado School of Medicine, Denver, Colorado; ²Denver Health and Hospital Authority, Denver, Colorado; ³University of Colorado, School of Medicine, Aurora, CO; ⁴Children's Hospital Colorado, Aurora, CO; ⁵Denver Health Medical Center, University of Colorado School of Medicine, Denver, Colorado

Session: P-14. COVID-19 Complications, Co-infections, and Clinical Outcomes

Background. Reports in adults with COVID-19 and acute otitis media (AOM) show that severe symptoms and hearing loss may be more common than with the clinical presentation of typical AOM. However, the association of SARS-CoV-2 with AOM in children is poorly understood.

Methods. Cases were identified as a subpopulation enrolled in the NOTEARS prospective AOM study in Denver, CO from March-December 2020. Children enrolled were 6-35 months of age with uncomplicated AOM and prescribed amoxicillin. Children diagnosed with AOM and SARS-CoV-2, detected by polymerase chain reaction assay, were included in the case series. Data was obtained from electronic medical records and research case report forms. Patients completed surveys at enrollment and 5, 14 and 30 days after enrollment that included the Acute Otitis Media Severity of Symptoms (AOM-SOS[®]) scale. All patients had nasopharyngeal otopathogen testing completed.

Results. A total of 108 patients had been enrolled through December 2020 (all of whom were subsequently tested for SARS CoV-2). During the study period for this case series, 16 patients were enrolled, and 7 (43.6%) were identified with AOM/SARS-CoV-2 co-infection. Among these 7 patients, fever was present in 3 children (29%). Four children (57%) attended daycare. Only 2 children (29%) had testing for SARS CoV-2 as part of their clinical workup. Mean AOM-SOS[®] scores were similar among the SARS CoV-2 positive and negative patients with no statistical significance noted with two-sided t-tests: 13.6 (± 4.5) vs 14.2 (± 4.9) at enrollment, 1.4 (± 1.8) vs 4.2 (±4.9) on Day 5, and 0.6 (± 0.9) vs. 2.5 (±6.1) on Day 14 (Table 1). Among the 7 patients, no child had an AOM treatment failure or recurrence. Of the 6 patients in whom bacterial and viral testing have been completed, a bacterial otopathogen was identified in 6 (100%), and a viral pathogen in 3 (50%) children (Table 2).

Table 1. Clinical features of children with concurrent SARS-CoV-2 and AOM

Category	Characteristic	Case 1	Case 2	Case 3	Case 4	Case 5	Case 6	Case 7
Demographics	Age (months)	30	29	16	8	11	7	24
	Gender-Male	+	+	+	+	+	+	+
	Race other than white	++				+		
	Hispanic	+		+	+	+	+	+
Medical History	Public Insurance	+		+	+	+	+	+
	Past medical problems					±	±	
	2+ doses PCV	+	+		+	+	+	+
	Smoke exposure	+			+			
Clinical Presentation	Breastfeeding	+	+	+	+	+	+	+
	Daycare attendance	+	+	+	+	+	+	+
	Ear pain/tugging	+	+	+	+	+	+	+
	Nasal congestion	+	+	+	+	+	+	+
Outcomes	Fussiness/irritability	+	+	+	+	+	+	+
	Cough	+	+	+	+	+	+	+
	Reduced oral intake							
	Eye redness/pain/discharge							
	Fever, subjective or over 100.4							
	Vomiting							
	Diarrhea							
	Reduced urine output							
	Bilateral infection							
	AOM-SOS [®] a							
Outcomes	Diagnosis	10	6	10	18	15	18	18
	Day 5	0	0	0	4	4	0	2
	Day 14	0	0	2	0	0	0	0
	Treatment Failure	No	No	No	No	No	No	No
	Recurrence	No	No	No	No	No	No	No

¹Unknown/not reported ²Black ³Prenatal methadone exposure, wheezing ⁴Prematurity ⁵Acute otitis media severity of symptoms score (UPMC, Pittsburgh, PA); Minimum significant clinical different[®] 20%; scores of 4 and lower are considered equivalent to a normal state of health.

Table 2. Laboratory findings of children with concurrent SARS-CoV-2 and AOM.

Laboratory Test ^a	Specific Pathogen	Case 1	Case 2	Case 3	Case 4	Case 5	Case 6	Case 7
SARS-CoV-2 PCR	Clinical SARS-CoV-2 PCR (Abbott)							
	Research SARS-CoV-2 PCR (Quidel)	+	+	+	+	+	+	NYT ^b
Respiratory Viral PCR	Adenovirus							NYT
	Influenza A							NYT
	Influenza B							NYT
	Parainfluenza 1							NYT
	Parainfluenza 2							NYT
	Parainfluenza 3							NYT
	RSV							NYT
	Human Metapneumovirus					+		NYT
	Rhinovirus					+		NYT
	Coronavirus spp. (not SARS-CoV-2 or MERS)							NYT
	MERS							NYT
	Enterovirus							NYT
	Bacterial PCR	<i>S. pneumoniae</i>	+	+	+	+	+	+
<i>H. influenzae</i>		+	+	+	+	+	+	NYT
<i>M. catarrhalis</i>		+	+	+	+	+	+	NYT
<i>S. aureus</i>								NYT
Culture	<i>S. pneumoniae</i>		+	+	+	+	+	±
	<i>H. influenzae</i>		+	+	+	+	+	±
	<i>M. catarrhalis</i>		+	+	+	+	+	±
	<i>S. aureus</i>							±

¹All testing completed using NP swabs collected at enrollment ²NYT- not yet tested ³Inconclusive ⁴Culture completed after receiving 1 dose of antibiotics

Conclusion. SARS-CoV-2 can occur in children with AOM. It is important that providers maintain a high index of suspicion for COVID-19 even in patients with clinical evidence of AOM, particularly to ensure families are appropriately advised on isolation and quarantine requirements. AOM with SARS-CoV-2 does not appear to be more severe than AOM without SARS-CoV-2.

Disclosures. Samuel R. Dominguez, MD, PhD, BioFire Diagnostics (Consultant, Research Grant or Support)DiaSorin Molecular (Consultant)Pfizer (Grant/Research Support) Samuel R. Dominguez, MD, PhD, BioFire (Individual(s) Involved: Self): Consultant, Research Grant or Support; DiaSorin Molecular (Individual(s) Involved: Self): Consultant; Pfizer (Individual(s) Involved: Self): Grant/Research Support

348. Characteristics and Outcomes in Hospitalized Patients with Covid-19 Complicated by Fungemia: A Single Center Retrospective Study

Aleena Zahra, MD¹; Bettina C. Fries, MD/FIDSA²; Bettina C. Fries, MD/FIDSA²; Sutthichai Sae-Tia, MD¹; ¹SUNY Stony Brook University Hospital, Stony Brook, New York ²Stony Brook University Hospital, Stony Brook, NY

Session: P-14. COVID-19 Complications, Co-infections, and Clinical Outcomes

Background. Covid19 caused by SARS-CoV2 can lead to significant morbidity and mortality. Fungemia is a rare hospital-associated infection and there are limited data regarding its association with Covid19. We reviewed all cases of fungemia in our Covid19 cohort at Stony Brook University Hospital (SBUH).

Methods. We conducted a retrospective medical record review of patients admitted with Covid19 in a 3-month interval. We reviewed positive blood cultures for fungi and recorded co-morbidities, co-infections, length of stay, treatments, and outcomes (survival vs death). There were 60 positive blood cultures for fungi in 25 unique patients (Table 1); in prior years < 30 per year reported at SBUH.

Table 1. Fungal Blood Cultures

Fungal Species Name	Number of Unique Patients with Positive Fungal Blood Cultures
CANDIDA ALBICANS	8
CANDIDA PARAPSILOSIS	7
CANDIDA TROPICALIS	3
CANDIDA LUSITANIAE	3
CANDIDA GLABRATA	2
CANDIDA DUBLINIENSIS	1
CRYPTOCOCCUS NEOFORMANS	1
Total Positive Cultures	25

Collation of each unique identified fungal species from fungal blood cultures in patients hospitalized with Covid-19

Results. During a 3 month interval at the local peak of the pandemic 1398 patients hospitalized with Covid19 at SBUH, 25 cases of fungemia were detected; *C. albicans* (CA) n=8,32%, non *C. albicans* species (nCA) n=16,64%, and *C. neoformans* n=1,4%, 17/25 (68%) also with bacteremia during same hospitalization. In same 3 months there were 264 cases of bacteremia and Covid19 co-infection. Demographics and medical co-morbidities of fungemic patients are in Table 2. Majority were men (76%). No difference between fungemic (FC) and total cohort (TC) in median age (62 vs 62), DM p=0.31, HTN p=1.0, COPD p=0.12. Within FC, DM was higher in nCA group (58.8%) vs CA group (37%). Mortality was 40% in FC vs 15% in TC, p< 0.001. Within FC mortality was 56% in nCA and 25% in CA group. *C. parapsilosis* was the most common nCA species isolated with 43% mortality. FC more likely to require ICU and mechanical ventilation (88% vs 15%, p< 0.0001) and had longer median length of stay 42 days vs 22 days. The median time from admission to fungemia was 21d, from central line placement 19d, Table 3. Of FC 21 (84%) were treated with steroids/Tocilizumab concurrently. Of note, no mortality was recorded in the 4 patients that did not receive steroids/Tocilizumab.

PCT and WBC were significantly higher at time of fungemia as compared to admission, Table 3.

Table 2. Patient co-morbidities and hospitalization stay characteristics

	Total Cohort, N=1325	Fungemic Cohort, N=25	p-value
Hypertension	464	15	1.0
Diabetes mellitus (DM)	324	13	0.31
Coronary Artery Disease (CAD)	254	8	0.12
Chronic Kidney Disease (CKD)	194	1	0.24
COPD/Asthma	256	6	0.12
Malignancy	188	2	0.56
Intensive Care Unit (ICU) admission	198	22	<0.0001
Mechanical Ventilation	198	22	<0.0001
Age	62 (49-75)	62 (53-75)	1.0
Male	755	19	0.06
Female	570	6	0.06
Caucasian	739	8	0.02
Hispanic	343	11	0.06
Black or African American	92	2	0.69
Asian	47	4	0.01

Co-morbidities and requirement for ICU stay, mechanical ventilation for total cohort Covid-19 and fungemic cohort

Table 3. Patient Characteristics and Laboratory Parameters

Patient Characteristics	Median	IQR	Laboratory Parameters	Mean Difference Day Fungemia-Day Hospitalization	SEM, p value
BMI kg/m ²	26.12	22.9–32.73	Procalcitonin ng/mL	1.81	+/- 1.05 p<0.05
SARS-CoV2 PCR positive from hospitalization (day)	1	-1 – 1	C-Reactive Protein mg/dL	-7.68	+/- 2.09 p<0.0006
Temperature (C) on day of fungemia	38.1	37.3-38.6	WBC K/uL	6.58	+/- 1.75 P<0.0005
Time Fungemia from hospitalization (day)	21	14-37			
Time Fungemia from central line placement (day)	19	8.5-25			
Total Number Antibiotics (1+dose/antibiotic)	6	5-7			