

RESEARCH ARTICLE

Open Access

The renal urate transporter SLC17A1 locus: confirmation of association with gout

Jade E Hollis-Moffatt^{1†}, Amanda J Phipps-Green^{1†}, Brett Chapman², Gregory T Jones³, Andre van Rij³, Peter J Gow⁴, Andrew A Harrison⁵, John Highton³, Peter B Jones⁶, Grant W Montgomery², Lisa K Stamp⁷, Nicola Dalbeth⁶ and Tony R Merriman^{1*}

Abstract

Introduction: Two major gout-causing genes have been identified, the urate transport genes *SLC2A9* and *ABCG2*. Variation within the *SLC17A1* locus, which encodes sodium-dependent phosphate transporter 1, a renal transporter of uric acid, has also been associated with serum urate concentration. However, evidence for association with gout is equivocal. We investigated the association of the *SLC17A1* locus with gout in New Zealand sample sets.

Methods: Five variants (*rs1165196*, *rs1183201*, *rs9358890*, *rs3799344*, *rs12664474*) were genotyped across a New Zealand sample set totaling 971 cases and 1,742 controls. Cases were ascertained according to American Rheumatism Association criteria. Two population groups were studied: Caucasian and Polynesian.

Results: At *rs1183201* (*SLC17A1*), evidence for association with gout was observed in both the Caucasian (odds ratio (OR) = 0.67, $P = 3.0 \times 10^{-6}$) and Polynesian (OR = 0.74, $P = 3.0 \times 10^{-3}$) groups. Meta-analysis confirmed association of *rs1183201* with gout at a genome-wide level of significance (OR = 0.70, $P = 3.0 \times 10^{-8}$). Haplotype analysis suggested the presence of a common protective haplotype.

Conclusion: We confirm the *SLC17A1* locus as the third associated with gout at a genome-wide level of significance.

Introduction

Regulation of serum urate concentration is central to the development of gout, with renal uric acid excretion a critical checkpoint [1]. Genome-wide association scans examining the genetic control of serum urate concentrations have identified two renal urate transporters - *SLC2A9* and *ABCG2* - that have a strong effect on gout risk in multiple ethnic groups [2]. Whilst other loci (*SLC22A11*, *GCKR*, *INHBC*, *SLC17A1*, *RREB1*, *PDZK1*, *SLC16A9*, *LRRC16A*) have been associated with serum urate concentrations at a genome-wide level of significance in genome-wide association scans [3,4], only some of them (*SLC22A11*, *GCKR*, *INHBC*, *SLC17A1*) were associated with gout at a nominal level of significance ($P < 0.05$) in 1,100 cases nested within a large genome-wide association scan population-based cohort [4]. To understand why some loci do not associate

with gout, and to assess the weakly associated loci in clinical gout, it will be necessary to minimize heterogeneity owing to the type of gout (primary or secondary to other causes such as diuretic use) and to test for association in clinically proven cases.

The solute carrier family 17 member 1 (encoded by *SLC17A1*), also known as sodium phosphate transport protein 1 (NPT1), is expressed on the apical membrane of renal tubular cells and mediates sodium and inorganic phosphate co-transport [5]. Sodium-dependent transporter 1 has also been identified as a urate transport protein [6,7], probably secretory [7] with the gout-protective allele of I269T [8] leading to increased sodium-dependent transporter 1 activity [6] and, presumably, increased secretion of uric acid. Genome-wide association scans have shown that genetic variants associate with serum urate concentration in a Caucasian sample [3,4]. *SLC17A1* has been associated with gout in a Japanese sample set (I269T (*rs1165196*), odds ratio (OR) = 0.55, $P = 0.005$) [8] but with conflicting results in Caucasian sample sets. Marker

* Correspondence: tony.merriman@otago.ac.nz

† Contributed equally

¹Department of Biochemistry, University of Otago, 710 Cumberland Street, Dunedin 9054, New Zealand

Full list of author information is available at the end of the article

rs1165205 in *SLC17A3* was first associated with gout (OR = 0.85, $P = 0.002$) [9]. A later study incorporating the same clinical material with additional cases and controls, however, reported reduced combined evidence for association with gout using a strongly correlated marker within *SLC17A1* (*rs1165196*, $r^2 = 0.96$; OR = 0.89, $P = 0.013$) [4] - in this study the markers most strongly associated with serum urate were within *SLC17A1* (*rs1165196* and other tightly correlated markers), suggesting that this gene was more likely than *SLC17A3* to harbor an etiological variant. A separate study reported no evidence in Caucasian for association with gout (*rs1183201*, r^2 with *rs1165196* = 0.87, OR = 0.97, $P = 0.68$) [10]. This equivocal evidence for association with gout in a Caucasian population is notable given the genome-wide evidence for association with serum urate concentration [4]. Both studies had adequate power to detect association of a moderate effect size, but neither study used clinical criteria to define gout.

Here, we aimed to test the *SLC17A1* locus for association with gout, in multiple ancestral groups, using cases defined as a diagnosis of gout by the 1977 American College of Rheumatology (ARA) clinical criteria. The variants tested were *rs1183201*, demonstrated to influence serum urate concentration in Caucasian populations [3], the maximally gout-associated SNP (*rs1165196* (I269T)) in Japanese [8], and three other SNPs predicted to tag major variation in Polynesian populations.

Materials and methods

Study participants

There were a total of four New Zealand (NZ) case-control sample sets, one of Caucasian ancestry and three of different Polynesian ancestries (see Supplemental Table S1 in Additional file 1). The sample sets were Eastern Polynesian (EP; NZ Māori and Cook Islands, 284 cases and 349 controls), Western Polynesian (WP; Samoa, Tonga, Niue and Tokelau, 251 cases and 144 controls), combined Eastern and Western Polynesian (EP/WP; 15 cases and 21 controls) and Caucasian (421 cases and 1,228 controls; of the controls, 590 had been SNP typed genome wide [11,12]). The EP samples were further subdivided into two groups to remove effects of stratification, as described in more detail below, based on the estimated proportion of EP ancestry (EP/N, 236 cases and 192 controls; and EP/Z, 48 cases and 157 controls). All gout cases recruited had a diagnosis of gout confirmed according to the ARA preliminary diagnostic criteria [13]. Controls self-reported as having no history of gout. Recruitment of cases was approved by the NZ Multi-Region Ethics Committee (MEC/05/10/130), and recruitment of controls by the Lower South Ethics Committee. All patients provided written informed consent for the collection of samples and subsequent analysis.

Analysis of genome-wide microsatellite data indicates a difference in population structure between Samoa and NZ Māori [14] - with a Māori sample set estimated to be ~85% Polynesian and ~15% Caucasian ancestry, and a Samoan sample set estimated at ~70% Polynesian, ~15% Asian and smaller components of Melanesian and Caucasian ancestry [14]. Analysis of genome-wide SNP data by principal component analysis also shows a difference in the first component between Samoan and Cook Island genomes [15]. Given data also showing heterogeneity in association of *ABCG2* with gout in EP and WP sample sets [16], the analysis groups here were EP and WP. People of combined EP and WP ancestry were included as a separate group.


Power

The individual sample sets were inadequately powered to detect an effect size in gout equivalent to that reported previously (OR = 1.12) [4], with the largest dataset (Caucasian) estimated to have 29% power at $\alpha = 0.05$. However, at larger effect sizes the Caucasian sample set was better powered (63% power at OR = 1.2, 90% power at OR = 1.3 and 99% power at OR = 1.4). The smaller individual Polynesian sample sets had less power (for OR = 1.4: EP = 67% and WP = 45% using a Han Chinese Beijing (CHB) estimate of minor allele frequency = 0.18), although the combined Polynesian sample set was adequately powered with 88% power at OR = 1.4.

SNP selection and determination of genotypes

Using CHB HapMap data as the most closely related and available population to Polynesia, Haploview software (Broad Institute, Cambridge, MA, USA) was used to define four haplotype blocks using the Gabriel confidence interval method covering *SLC17A1* (defined by *rs4712972* (25.772 Mb) to *rs12192635* (25.881 Mb)). Variants tagging major haplotypes were selected: *rs9358890* in block 1, *rs3799344* in block 2, *rs1183201* in block 3 (previously associated with control of serum urate concentration) [3] and *rs12664474* in block 4. The haplotype blocks extended into flanking genes (*SLC17A4* and *SLC17A3*). In Centre d'Etude du Polymorphisme Humain (CEU) Caucasian population and the CHB population, *rs1183201* and *rs3799344* exhibited some intermarker linkage disequilibrium (LD) ($r^2 = 0.77$ and 0.50 , respectively) and *rs9358890* and *rs12664474* also exhibited LD ($r^2 = 0.35$ and 0.62 , respectively). SNP *rs9358890* is in *SLC17A4*, and *rs12664474* is in *SLC17A3*. SNP *rs1165196* (*SLC17A1*) was also selected, in strong LD with *rs1183201* ($r^2 = 0.87$ in CEU and 0.91 in CHB) (Figure 1).

Genotyping was done by TaqMan[®] SNP genotyping assays (Applied Biosystems, Foster City, CA, USA) using a Lightcycler[®] 480 Real-Time PCR System (Roche,



Indianapolis, IN, USA) for four SNPs: *rs1183201* (assay ID: C_1911034_10), *rs9358890* (assay ID: C_25595118_10), *rs3799344* (assay ID: C_194536_10) and *rs12664474* (assay ID: C_11189653_10). SNP *rs1165196* was genotyped using Sequenom technology (Sequenom, Inc. San Diego, CA, USA). SNPs *rs9358890* and *rs12664474* had been genotyped over 590 of the Caucasian controls on the Affymetrix 6 SNP array (Affymetrix, Santa Clara, CA, USA) [12] - genotypes were imputed for *rs3799344* and *rs1183201* with IMPUTE2, using HapMap3 CEU (NCBI Build 36 (db126b)) as reference haplotypes.

Statistical analysis

ORs were calculated using PLINK software [17]. Because the case-control sample sets were not matched for sex, association analysis also included sex as a possible confounder. Analysis of association of haplotypes was also performed using PLINK. Meta-analysis was carried out using Rmeta software (within STATA 8.0, Stata, College Station, TX, USA) to calculate the combined Mantel-Haenszel OR using a fixed-effects model and the Breslow-Day test for heterogeneity.

The Māori population of NZ is admixed, primarily Caucasian. This leads to genetic stratification, which is a confounding factor for case-control genetic studies, especially when the prevalence of disease differs between the interbreeding populations. The prevalence of gout in NZ Māori is approximately double that in NZ Caucasian [18]. Given this, it is not surprising that the EP case sample set, which is predominantly NZ Māori (78% of cases, 93% of

controls), has a significantly greater proportion of self-reported EP grandparents than does the control sample set (average of 3.1 EP grandparents in cases vs. 2.5 in controls, $P = 3.1 \times 10^{-14}$ by t test). Sixty-seven biallelic genomic control markers (see Supplemental Table 2 in Additional file 1) were genotyped in the EP sample set, and STRUCTURE software [19] was used to estimate the individual proportion of EP ancestry. This estimation was performed using the following parameters: number of populations assumed to be two, 30,000 burn-in period, and 100,000 Markov chain Monte Carlo replications after burn-in. Caucasian control individuals genotyped for the 67 markers were included as representative of the ancestral Caucasian population to aid in population clustering, although we were unable to include EP ancestral representatives. Plots of self-reported ancestry versus STRUCTURE estimated ancestry are shown in Supplemental Figure 1 in Additional file 1. For association analysis we created two datasets matched for EP ancestry - EP/N with estimated EP ancestry > 0.65 (the geometric mean; 236 cases and 192 controls), and EP/Z with estimated EP ancestry ≤ 0.65 (48 cases and 157 controls). The estimated average proportion of EP ancestry in the EP/N sample set was 0.90 in cases and 0.88 in controls, and for the EP/Z group was 0.41 in cases and 0.40 in controls.

Results

Association with gout was observed in the NZ Caucasian sample set for *rs1165196*, *rs1183201*, *rs3799344* and *rs12664474* (OR = 0.71 (95% confidence interval

(CI) = 0.60 to 0.83), $P = 5.5 \times 10^{-5}$; OR = 0.67 (95% CI = 0.57 to 0.79), $P = 3.0 \times 10^{-6}$; OR = 0.69 (95% CI = 0.58 to 0.81), $P = 2.8 \times 10^{-5}$; and OR = 1.36 (95% CI = 1.12 to 1.66), $P = 1.3 \times 10^{-3}$, respectively), but not for *rs9358890* (OR = 1.31 (95% CI = 0.93 to 1.85), $P = 0.17$) (Table 1). Given the low LD between *rs12664474* and *rs1183201* in CEU ($r^2 = 0.16$), suggesting the possibility of an independent effect at *rs12664474*, we tested for association of *rs12664474* conditional on genotype at *rs1183201* in the NZ Caucasian samples; there was no evidence for a separate genetic effect on gout risk at *rs12664474* ($P = 0.37$). We also tested for conditional associations at *rs1183201* and *rs1165196* (r^2 in controls = 0.90) - there was association at *rs1183201* conditional on genotype at *rs1165196* ($P = 0.007$), but not at *rs1165196* when conditioned on genotype at *rs1183201* ($P = 0.14$).

The five variants were then tested for association in the Polynesian sample sets (Table 1), with the only evidence for association in individual sample sets coming from WP at *rs1183201* (OR = 0.70, $P = 0.03$) and *rs3799344* (OR = 0.67, $P = 0.02$). However, meta-analysis of the Polynesian sample sets - carried out to increase power - replicated the association observed in Caucasian at *rs1165196* (OR = 0.75 (95% CI = 0.60 to 0.94), $P = 0.013$, $P_{\text{Het}} = 0.33$), *rs1183201* (OR = 0.74 (95% CI = 0.61 to 0.91), $P = 0.003$, $P_{\text{Het}} = 0.57$) and *rs3799344* (OR = 0.74 (95% CI = 0.61 to 0.90), $P = 0.003$, $P_{\text{Het}} = 0.33$), but not at *rs9358890* (OR = 1.15 (95% CI = 0.95 to 1.40), $P = 0.16$, $P_{\text{Het}} = 0.28$) or *rs12664474* (OR = 1.16 (95% CI = 0.96 to 1.40), $P = 0.13$, $P_{\text{Het}} = 0.23$).

The Caucasian and Polynesian sample sets were combined in meta-analysis for *rs1165196* (OR = 0.72 (95% CI = 0.64 to 0.82), $P = 5.7 \times 10^{-7}$), *rs1183201* (OR = 0.70 (95% CI = 0.62 to 0.79), $P = 3.0 \times 10^{-8}$, $P_{\text{Het}} = 0.64$), *rs9358890* (OR = 1.19 (95% CI = 1.00 to 1.41), $P = 0.05$, $P_{\text{Het}} = 0.37$), *rs3799344* (OR = 0.71 (95% CI = 0.62 to 0.80), $P = 7.4 \times 10^{-8}$, $P_{\text{Het}} = 0.43$), and *rs12664474* (OR = 1.25 (95% CI = 1.09 to 1.43), $P = 2.0 \times 10^{-3}$, $P_{\text{Het}} = 0.23$). Of the five SNPs, *rs1183201* was the only one significant at a genome-wide level of significance ($P < 5 \times 10^{-8}$). None of the SNPs were significantly associated with serum urate in either the Caucasian controls (for whom there were serum urate data available; see Supplemental Table 1 in Additional file 1) or the less admixed combined WP and EP/N controls (all $P > 0.28$).

Because haplotypes are multi-allelic we analyzed association of haplotypes with gout, with the purpose of investigating the mechanism of effect - that is, whether risk and/or protective variants are present and comparing association pattern between populations. Analysis of four-marker haplotypes (*rs9358890-rs3799344-rs1183201-rs12664474*; Table 2) revealed the most consistent evidence for association to come from the A-T-A-A haplotype (OR = 0.30 to

0.84), with significant association in the Caucasian, WP and EP/WP sample sets ($P = 1.5 \times 10^{-6}$ to 0.035).

Discussion

Genetic regulators of serum urate concentration that have been previously associated with gout at a genome-wide level of significance ($P < 5 \times 10^{-8}$) in Caucasian samples are *SLC2A9* [4,9,20] and *ABCG2* [4,9,16]. Here, we provide strong evidence for a role of the *SLC17A1* locus in gout in a Caucasian population (*rs1183201*, OR = 0.67, $P = 3.0 \times 10^{-6}$; Table 1) that was replicated in Polynesian samples, with the minor allele of *rs1183201* also conferring a similar degree of risk (OR = 0.74, $P_{\text{meta-analysis}} = 3.0 \times 10^{-3}$). The haplotype data (Table 2) are consistent with the presence of at least one genetic variant influencing the risk of gout at the *SLC17A1* locus. We hypothesize that the variant is protective of gout and is contained on a common haplotype (27 to 43%; A-T-A-A), conferring significant protection in three out of the five sample sets (also with OR < 1 in both EP sample sets). There were no haplotypes consistently conferring risk. Combining the populations provided a genome-wide level of significance for association of *rs1183201* with gout (OR = 0.70, $P = 3.0 \times 10^{-8}$). This confirms the *SLC17A1* locus as the third associated with gout.

The role of *SLC17A1* has been previously evaluated in gout in a Japanese sample set [8], with the nonsynonymous variant I269T (*rs1165196*) having the strongest evidence for association (OR = 0.55, $P = 0.004$, minor allele (269T) protective). *rs1165196* is in strong LD with *rs1183201* - the maximally associated variant in our study - in Japanese (HapMap JPT) and Caucasian (HapMap CEU) samples ($r^2 = 0.92$ and $r^2 = 0.87$, respectively). Given that I269T has been shown to affect the function of SLC17A1, with the protective variant (269T, minor allele of *rs1165196*) leading to increased activity in *Xenopus* oocytes and, presumably, increased renal elimination of urate [6], it is therefore possible that *rs1165196* is an etiological variant. However, we found no evidence in the Caucasian sample set supporting association at *rs1165196* when conditioned on genotype at *rs1183201*, and association was weaker at *rs1165196* than *rs1183201* in combined Caucasian and Polynesian meta-analysis (OR = 0.72, $P = 5.7 \times 10^{-7}$ and OR = 0.70, $P = 3 \times 10^{-8}$, respectively) and in Polynesian alone (OR = 0.75, $P = 0.013$ and OR = 0.74, $P = 0.003$, respectively) (we did not conditionally analyze the small Polynesian sample sets). Ostensibly this observation argues that *rs1183201* (or a variant in strong LD) is more likely than *rs1165196* to be an etiological variant within *SLC17A1*. Given that *rs1165196* has a stronger effect in serum urate levels in Caucasian ([4] $\beta = 6.205$ vs. 6.050 for *rs1183201*) populations, however, this interpretation

Table 1 Association analysis in New Zealand case-control sample sets

SNP ^a	Cases			MAF	Controls			MAF	<i>P</i> _{unadjusted}	<i>P</i> _{adjusted} ^b	OR (95% CI)	HWE case	HWE control
<i>rs1165196</i>	TT	CT	CC	C	TT	CT	CC	C					
Caucasian	167 (0.400)	201 (0.482)	49 (0.118)	0.359	389 (0.318)	590 (0.482)	246 (0.200)	0.442	1.0 × 10 ⁻⁴	5.5 × 10 ⁻⁵	0.71 (0.60 to 0.83)	0.33	0.41
EP/N	130 (0.570)	86 (0.377)	12 (0.053)	0.241	98 (0.516)	77 (0.405)	15 (0.079)	0.282	0.19	0.26	0.81 (0.60 to 1.11)	0.65	0.98
EP/Z	16 (0.364)	22 (0.500)	6 (0.136)	0.386	53 (0.358)	64 (0.432)	31 (0.209)	0.426	0.51	0.88	0.85 (0.52 to 1.38)	0.72	0.16
WP	138 (0.556)	99 (0.399)	11 (0.044)	0.244	69 (0.496)	54 (0.388)	16 (0.115)	0.309	0.05	0.06	0.72 (0.52 to 1.00)	0.20	0.28
EP/WP	10 (0.667)	5 (0.333)	0 (0.000)	0.167	6 (0.316)	10 (0.526)	3 (0.158)	0.421	0.02	0.07	0.28 (0.09 to 0.87)	0.33	0.73
Combined ^c									5.7 × 10 ⁻⁷		0.72 (0.64 to 0.82)		
<i>rs1183201</i> ^d	TT	TA	AA	A	TT	TA	AA	A					
Caucasian	158 (0.384)	205 (0.499)	48 (0.117)	0.366	356 (0.291)	608 (0.496)	261 (0.213)	0.461	2.0 × 10 ⁻⁶	3.0 × 10 ⁻⁶	0.67 (0.57 to 0.79)	0.13	0.89
EP/N	130 (0.570)	86 (0.377)	12 (0.053)	0.241	96 (0.505)	78 (0.411)	16 (0.084)	0.289	0.11	0.14	0.78 (0.57 to 1.06)	0.65	0.98
EP/Z	15 (0.349)	21 (0.488)	7 (0.163)	0.407	52 (0.349)	61 (0.409)	36 (0.242)	0.446	0.49	0.95	0.85 (0.52 to 1.39)	0.94	0.04
WP	132 (0.543)	101 (0.416)	10 (0.041)	0.249	68 (0.479)	57 (0.401)	17 (0.120)	0.320	0.03	0.03	0.70 (0.51 to 0.97)	0.08	0.35
EP/WP	9 (0.600)	6 (0.400)	0 (0.000)	0.200	7 (0.350)	10 (0.500)	3 (0.150)	0.400	0.07	0.11	0.38 (0.13 to 1.12)	0.25	0.99
Combined ^c									3.0 × 10 ⁻⁸		0.70 (0.62 to 0.79)		
<i>rs9358890</i>	AA	AG	GG	G	AA	AG	GG	G					
Caucasian	366 (0.884)	47 (0.114)	1 (0.002)	0.059	1114 (0.912)	104 (0.085)	4 (0.003)	0.046	0.12	0.17	1.31 (0.93 to 1.85)	0.69	0.35
EP/N	97 (0.418)	99 (0.427)	36 (0.155)	0.369	80 (0.419)	78 (0.408)	33 (0.173)	0.377	0.80	0.71	0.96 (0.73 to 1.28)	0.21	0.07
EP/Z	34 (0.753)	11 (0.244)	0 (0.000)	0.122	129 (0.827)	26 (0.167)	1 (0.006)	0.090	0.44	0.42	1.41 (0.67 to 2.96)	0.35	0.80
WP	110 (0.440)	101 (0.404)	39 (0.156)	0.358	73 (0.507)	61 (0.424)	10 (0.069)	0.281	0.03	0.06	1.43 (1.04 to 1.95)	0.06	0.57
EP/WP	8 (0.533)	7 (0.467)	0 (0.000)	0.233	12 (0.571)	7 (0.333)	2 (0.095)	0.262	0.78	0.87	0.86 (0.29 to 2.55)	0.24	0.53
Combined ^c									0.05		1.19 (1.00 to 1.41)		
<i>rs3799344</i>	CC	CT	TT	T	CC	CT	TT	T					
Caucasian	165 (0.404)	193 (0.473)	50 (0.123)	0.359	379 (0.309)	592 (0.483)	255 (0.207)	0.450	5.9 × 10 ⁻⁶	2.8 × 10 ⁻⁵	0.69 (0.58 to 0.81)	0.58	0.38
EP/N	128 (0.561)	88 (0.386)	12 (0.053)	0.246	97 (0.524)	73 (0.395)	15 (0.081)	0.278	0.46	0.37	0.84 (0.62 to 1.15)	0.53	0.80
EP/Z	17 (0.378)	22 (0.489)	6 (0.133)	0.378	53 (0.353)	64 (0.427)	33 (0.220)	0.433	0.35	0.72	0.79 (0.49 to 1.29)	0.79	0.49
WP	136 (0.551)	98 (0.397)	13 (0.053)	0.251	64 (0.457)	59 (0.421)	17 (0.121)	0.332	0.02	0.02	0.67 (0.49 to 0.93)	0.39	0.55
EP/WP	10 (0.667)	5 (0.333)	0 (0.000)	0.167	7 (0.350)	10 (0.500)	3 (0.150)	0.400	0.04	0.08	0.30 (0.10 to 0.95)	0.44	0.85
Combined ^c									7.4 × 10 ⁻⁸		0.71 (0.62 to 0.80)		
<i>rs12664474</i>	AA	AG	GG	G	AA	AG	GG	G					
Caucasian	264 (0.632)	134 (0.321)	20 (0.048)	0.208	867 (0.707)	321 (0.262)	38 (0.031)	0.162	0.01	1.2 × 10 ⁻³	1.36 (1.12 to 1.66)	0.57	0.22
EP/N	90 (0.383)	104 (0.443)	41 (0.174)	0.396	74 (0.392)	77 (0.407)	38 (0.201)	0.405	0.79	0.17	0.96 (0.73 to 1.27)	0.25	0.03
EP/Z	25 (0.543)	18 (0.391)	3 (0.065)	0.261	107 (0.704)	39 (0.257)	6 (0.039)	0.168	0.05	0.13	1.75 (1.00 to 3.05)	0.92	0.32
WP	102 (0.411)	106 (0.427)	40 (0.161)	0.375	63 (0.444)	67 (0.472)	12 (0.085)	0.320	0.13	0.10	1.27 (0.93 to 1.73)	0.16	0.32
EP/WP	7 (0.467)	6 (0.400)	2 (0.133)	0.333	12 (0.571)	7 (0.333)	2 (0.095)	0.262	0.51	0.38	1.41 (0.51 to 3.92)	0.70	0.53
Combined ^c									2.0 × 10 ⁻³		1.25 (1.09 to 1.43)		

CI, confidence interval; HWE, {AU Query: Define}; MAF, {AU Query: Define}; OR, odds ratio. ^aGenotyping success rate: *rs1165196* Caucasian, Eastern Polynesian (EP), Western Polynesian (WP) and EP/WP - cases 99.0%, 95.8%, 98.8% and 100%, respectively, controls 99.8%, 96.8%, 96.5% and 90.5%, respectively; *rs1183201* Caucasian, EP, WP and EP/WP - cases 97.6%, 95.4%, 96.8% and 100%, respectively, controls 99.7%, 97.1%, 98.6% and 95.2%, respectively; *rs9358890* Caucasian, EP, WP and EP/WP - cases 98.3%, 97.5%, 99.6% and 100%, respectively, controls 99.2%, 99.4%, 100% and 90.5%, respectively; *rs3799344* Caucasian, EP, WP, EP/WP - cases 96.9%, 96.1%, 98.4% and 100%, respectively, controls 99.8%, 96.0%, 97.2% and 95.2%, respectively; *rs12664474* Caucasian, EP, WP, EP/WP - cases 99.3%, 98.9%, 98.8% and 100%, respectively, controls 99.8%, 97.7%, 98.6% and 100%, respectively. ^bAdjusted for sex. ^cMeta-analysis by the Mantel-Haenszel method. Breslow-Day test for heterogeneity: *rs1165196*, *P* = 0.45; *rs1183201*, *P* = 0.75; *rs9358890*, *P* = 0.77; *rs3799344*, *P* = 0.58; *rs12664474*, *P* = 0.97. ^dPopulation attributable risk (the reduction in gout incidence if the risk variant was absent) for positivity of the risk (T) allele of *rs1183201* is 38.2% for Caucasian, 21.4% for EP/N, 27.3% for EP/N and 41.3% for WP.

Table 2 Association of four-marker *rs9358890-rs3799344-rs1183201-rs12664474* haplotypes with gout

Haplotype ^a	Frequency		OR (95% CI)	P value
	Case	Control		
Caucasian				
A-T-A-A	270 (0.339)	1044 (0.428)	0.66 (0.56 to 0.78)	1.5 × 10 ⁻⁶
A-C-T-A	327 (0.410)	876 (0.360)	1.22 (1.03 to 1.44)	0.014
A-C-T-G	121 (0.152)	284 (0.117)	1.34 (1.06 to 1.69)	0.015
G-C-T-G	47 (0.059)	104 (0.043)	1.39 (0.98 to 1.98)	0.067
EP/N				
A-T-A-A	107 (0.239)	101 (0.274)	0.84 (0.61 to 1.15)	0.27
A-C-T-A	160 (0.356)	109 (0.297)	1.33 (0.99 to 1.78)	0.06
G-C-T-G	161 (0.360)	140 (0.380)	0.93 (0.70 to 1.23)	0.60
EP/Z				
A-T-A-A	32 (0.372)	120 (0.414)	0.84 (0.51 to 1.38)	0.49
A-C-T-A	32 (0.372)	111 (0.384)	0.95 (0.58 to 1.56)	0.84
G-C-T-G	9 (0.105)	26 (0.090)	1.19 (0.53 to 2.64)	0.68
A-C-T-G	10 (0.116)	22 (0.075)	1.63 (0.74 to 3.59)	0.23
WP				
A-C-T-A	171 (0.357)	96 (0.343)	1.07 (0.78 to 1.45)	0.70
A-T-A-A	116 (0.241)	89 (0.318)	0.68 (0.49 to 0.94)	0.021
G-C-T-G	171 (0.356)	77 (0.275)	1.46 (1.06 to 2.02)	0.021
A-C-T-G	12 (0.025)	12 (0.043)	0.57 (0.25 to 1.29)	0.17
EP/WP				
A-T-A-A	5 (0.167)	16 (0.400)	0.30 (0.10 to 0.95)	0.035
A-C-T-A	15 (0.500)	13 (0.324)	2.08 (0.78 to 5.50)	0.14
G-C-T-G	6 (0.200)	11 (0.275)	0.66 (0.21 to 2.05)	0.47

CI, confidence interval; EP, Eastern Polynesian; OR, odds ratio; WP, Western Polynesian. ^aHaplotypes are listed in descending order of control frequency; those with frequency < 0.03 were excluded.

should await further testing in larger gout and serum urate sample sets.

In the Caucasian analysis, *rs1183201* was strongly associated with gout (OR = 0.67 (95% CI = 0.57 to 0.79)). This SNP, or SNPs in strong LD, has been studied for association with gout in two previous studies: Yang and colleagues [4], with OR = 0.89 (95% CI = 0.82 to 0.98); and Stark and colleagues [10], with OR = 0.97 (95% CI = 0.86 to 1.11). The strength of effect in our study is considerably greater than the previous studies, with a 95% CI that does not overlap with either study. Given that the control allele frequencies were similar between our study and those of Yang and colleagues [4] and Stark and colleagues [10] (0.461 (*rs1183201*), 0.46 (*rs1165196*), and 0.487 (*rs1183201*), respectively), the differences in effect size are therefore caused by differences in allele frequency in case sample sets. Differences in ascertainment of cases are a possible reason for this effect. Here, cases were clinically ascertained by ARA criteria with exclusion of patients suspected of having diuretic-induced gout. In Yang and colleagues' study, cases were drawn from five population-based cohorts and were ascertained by: self-report or allopurinol

treatment (AGES Reykjavik Study); self-report (Atherosclerosis Risk in Communities Study); receiving gout medication (allopurinol, colchicine, probenecid; Cardiovascular Health Study); self-report (Framingham Heart Study); and receiving gout medication (allopurinol, colchicine, probenecid, benzbromarone; Rotterdam Study) [4]. In Stark and colleagues' study, cases were ascertained by self-report and review of medical history [10].

In the study by Yang and colleagues no details were included about the inclusion, or otherwise, of diuretic-induced cases [4]; and in the study by Stark and colleagues 36.1% of cases were taking diuretic medication [10]. The use of self-reported gout probably results in participants without clinical evidence for gout being included in case sample sets; for example, only 69% of men who self-reported as new cases of gout met the ARA classification criteria for gout [21], and reliability and sensitivity for self-reported gout have been estimated at 63 to 73% and 84%, respectively [22]. Although the reliability of use of medications such as allopurinol, colchicine, probenecid and benzbromarone has not been extensively investigated for gout classification, the use of allopurinol prescription gives a positive predictive value of 39% for probable/definite gout [23]. Certainly, gout case sample sets ascertained using such indirect criteria had lower effect sizes reported at *SLC2A9*, compared with sets using ARA criteria [20]. The method of ascertainment in the previous studies [4,10] would thus reduce power to detect association at *SLC17A1* owing to inclusion of nongout participants in the case sample sets. The use of diuretic medications is well established as a gout risk factor [24], perhaps by inhibition of urate excretion mediated by human organic anion transporter 4 [25]. In Stark and colleagues' study [10], this could reduce power to detect association by studying cases with secondary gout, since the inhibitory effect of diuretic medication on organic anion transporter 4-mediated renal urate excretion would predominate over the genetic effect on gout risk mediated by the *SLC17A1* locus. It is also conceivable that diuretics directly influence the function of urate transporters encoded in the locus. The loop diuretic bumetanide has recently been shown to be a transport substrate for sodium-dependent transporter 4 (encoded by *SLC17A3*), and functional polymorphic variants are likely to influence transport ability [26]. Given the likelihood that gene-diuretic interactions exist, one would be prudent to exclude gout cases taking diuretic medication as a potential confounding factor in order to evaluate the direct effect of genetic variation in the *SLC17A1* locus on primary gout.

Conclusion

We provide, for the first time, a genome-wide level of evidence supporting a role for genetic variation in the

SLC17A1 locus in the etiology of gout. This is the third urate transport locus associated with gout with this robust level of evidence, and our results further emphasize the importance of urate transport in gout.

Additional material

Additional file 1: Supplemental Table 1 presenting participant demographic and clinical details, Supplemental Table 2 presenting genomic control SNPs, and Supplemental Figure 1 showing the correlation of self-reported number of EP grandparents with estimated EP ancestry using 67 genomic control markers.

Abbreviations

ABC: ATP-binding cassette; ARA: American Rheumatism Association; CEU: Centre d'Etude du Polymorphisme Humain; CHB: Han Chinese Beijing; CI: confidence interval; EP: Eastern Polynesian; LD: linkage disequilibrium; NZ: New Zealand; OR: odds ratio; SLC: solute carrier family; SNP: single nucleotide polymorphism; WP: Western Polynesian.

Acknowledgements

This work is supported by the Health Research Council of New Zealand. JE-H-M was supported by a New Zealand National Heart Foundation Research Fellowship. Jill Drake, Roddi Laurence, Karen Lindsay, Maria Lobo, Karen Pui and Gabrielle Sexton are thanked for assistance in recruitment, Morgan Smith for genotyping, and Marilyn Merriman and Ruth Topless for technical assistance.

Author details

¹Department of Biochemistry, University of Otago, 710 Cumberland Street, Dunedin 9054, New Zealand. ²Queensland Institute of Medical Research, 300 Herston Road, 4006 Brisbane, Queensland, Australia. ³Department of Medicine, University of Otago, 210 Great King Street, Dunedin 9016, New Zealand. ⁴Department of Rheumatology, Middlemore Hospital, 100 Hospital Road, Auckland 2025, New Zealand. ⁵Department of Medicine, University of Otago, 23A Mein Street, Wellington 6242, New Zealand. ⁶Department of Medicine, University of Auckland, 2 Park Road, Auckland 1023, New Zealand. ⁷Department of Medicine, University of Otago, 2 Riccarton Avenue, Christchurch 8140, New Zealand.

Authors' contributions

JE-H-M, AJP-G and TRM helped to design the study, oversee its execution, and prepare the manuscript. GTJ, AvR, PJG, AAH, JH, PBJ, LKS and ND helped to provide clinical recruitment and prepare the manuscript. BC and GWM helped to collect data and prepare the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 5 December 2011 Revised: 9 April 2012

Accepted: 27 April 2012 Published: 27 April 2012

References

1. Simmonds HA, McBride MB, Hatfield PJ, Graham R, McCaskey J, Jackson M: **Polynesian women are also at risk for hyperuricaemia and gout because of a genetic defect in renal urate handling.** *Br J Rheumatol* 1994, **33**:932-937.
2. Merriman TR, Dalbeth N: **The genetic basis of hyperuricaemia and gout.** *Joint Bone Spine* 2011, **38**:35-40.
3. Kolz M, Johnson T, Sanna S, Teumer A, Vitart V, Perola M, Mangino M, Albrecht E, Wallace C, Farrall M, Johansson A, Nyholt DR, Aulchenko Y, Beckmann JS, Bergmann S, Bochud M, Brown M, Campbell H, EUROSPAN Consortium, Connell J, Dominiczak A, Homuth G, Lamina C, McCarthy MI, ENGAGE Consortium, Meisinger T, Mooser V, Munroe P, Nauck M, Peden J, et al: **Meta-analysis of 28,141 individuals identifies common variants within five new loci that influence uric acid concentrations.** *PLoS Genet* 2009, **5**:e1000504.
4. Yang Q, Kottgen A, Dehghan A, Smith AV, Glazer NL, Chen MH, Chasman DI, Aspelund T, Eiriksdottir G, Harris TB, Launer L, Nalls M, Hernandez D, Arking DE, Boerwinkle E, Grove ML, Li M, Linda Kao WH, Chonchol M, Haritunians T, Li G, Lumley T, Psaty BM, Shlipak M, Hwang SJ, Larson MG, O'Donnell CJ, Upadhyay A, van Duijn CM, et al: **Multiple genetic loci influence serum urate and their relationship with gout and cardiovascular disease risk factors.** *Circ Cardiovasc Genet* 2010, **3**:523-530.
5. Busch AE, Schuster A, Waldegger S, Wagner CA, Zempel G, Broer S, Biber J, Murer H, Lang F: **Expression of a renal type I sodium/phosphate transporter (NaPi-1) induces a conductance in Xenopus oocytes permeable for organic and inorganic anions.** *Proc Natl Acad Sci USA* 1996, **93**:5347-5351.
6. Iharada M, Miyaji T, Fujimoto T, Hiasa M, Anzai N, Omote H, Moriyama Y: **Type 1 sodium-dependent phosphate transporter (SLC17A1 protein) is a Cl(-)-dependent urate exporter.** *J Biol Chem* 2010, **285**:26107-26113.
7. Uchino H, Tamai I, Yamashita K, Minemoto Y, Sai Y, Yabuuchi H, Miyamoto K, Takeda E, Tsuji A: **p-aminohippuric acid transport at renal apical membrane mediated by human inorganic phosphate transporter NPT1.** *Biochem Biophys Res Commun* 2000, **270**:254-259.
8. Urano W, Taniguchi A, Anzai N, Inoue E, Kanai Y, Yamanaka M, Endou H, Kamatani N, Yamanaka H: **Sodium-dependent phosphate cotransporter type 1 sequence polymorphisms in male patients with gout.** *Ann Rheum Dis* 2010, **69**:1232-1234.
9. Dehghan A, Köttgen A, Yang Q, Hwang SJ, Kao WL, Rivadeneira F, Boerwinkle E, Levy D, Hofman A, Astor BC, Benjamin EJ, van Duijn CM, Witteman JC, Coresh J, Fox CS: **Association of three genetic loci with uric acid concentration and risk of gout: a genome-wide association study.** *Lancet* 2008, **372**:1953-1961.
10. Stark K, Reinhard W, Grassl M, Erdmann J, Schunkert H, Illig T, Hengstenberg C: **Common polymorphisms influencing serum uric acid levels contribute to susceptibility to gout, but not to coronary artery disease.** *PLoS One* 2009, **4**:e7729.
11. Roberts RL, Van Rij AM, Phillips LV, Young S, McCormick SP, Merriman TR, Jones GT: **Interaction of the inflammasome genes CARD8 and NLRP3 in abdominal aortic aneurysms.** *Atherosclerosis* 2011, **218**:123-126.
12. Gretarsdottir S, Baas AF, Thorleifsson G, Holm H, den Heijer M, de Vries JP, Kranendonk SE, Zeebregts CJ, van Sterkenburg SM, Geelkerken RH, van Rij AM, Williams MJ, Boll AP, Kostic JP, Jonasdottir A, Jonasdottir A, Walters GB, Masson G, Sulem P, Saemundsdottir J, Mouy M, Magnusson KP, Tromp G, Elmore JR, Sakalihan N, Limet R, Defraigne JO, Ferrell RE, Ronkainen A, Ruigrok YM: **Genome-wide association study identifies a sequence variant within the DAB2IP gene conferring susceptibility to abdominal aortic aneurysm.** *Nat Genet* 2010, **42**:692-697.
13. Wallace SL, Robinson H, Masi AT, Decker JL, McCarty DJ, Yu TF: **Preliminary criteria for the classification of the acute arthritis of primary gout.** *Arthritis Rheum* 1977, **20**:895-900.
14. Friedlaender JS, Friedlaender FR, Reed FA, Kidd KK, Kidd JR, Chambers GK, Lea RA, Loo JH, Koki G, Hodgson JA, Merriwether DA, Weber JL: **The genetic structure of Pacific Islanders.** *PLoS Genet* 2008, **4**:e19.
15. Wollstein A, Lao O, Becker C, Brauer S, Trent RJ, Nürnberg P, Stoneking M, Kayser M: **Demographic history of Oceania inferred from genome-wide data.** *Curr Biol* 2010, **20**:1983-1992.
16. Phipps-Green AJ, Hollis-Moffatt JE, Dalbeth N, Merriman ME, Topless R, Gow PJ, Harrison AA, Highton J, Jones PB, Stamp LK, Merriman TR: **A strong role for the ABCG2 gene in susceptibility to gout in New Zealand Pacific Island and Caucasian, but not Maori, case and control sample sets.** *Hum Mol Genet* 2010, **19**:4813-4819.
17. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC: **PLINK: a tool set for whole-genome association and population-based linkage analyses.** *Am J Hum Genet* 2007, **81**:559-575.
18. Winnard D, Wright C, Taylor WJ, Jackson G, Te Karu L, Gow PJ, Arroll B, Thornley S, Gribben B, Dalbeth D: **National prevalence of gout derived from administrative health data in Aotearoa New Zealand.** *Rheumatology* 2012, **51**:901-909.
19. Pritchard JK, Stephens M, Donnelly P: **Inference of population structure using multilocus genotype data.** *Genetics* 2000, **155**:945-959.
20. Hollis-Moffatt JE, Xu X, Dalbeth N, Merriman ME, Topless R, Waddell C, Gow PJ, Harrison AA, Highton J, Jones PB, Stamp LK, Merriman TR: **Role of**

the urate transporter SLC2A9 gene in susceptibility to gout in New Zealand Maori, Pacific Island, and Caucasian case-control sample sets. *Arthritis Rheum* 2009, **60**:3485-3492.

21. Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G: **Purine-rich foods, dairy and protein intake, and the risk of gout in men.** *N Engl J Med* 2004, **350**:1093-1103.
22. McAdams MA, Maynard JW, Baer AN, Köttgen A, Clipp S, Coresh J, Gelber AC: **Reliability and sensitivity of the self-report of physician-diagnosed gout in the campaign against cancer and heart disease and the atherosclerosis risk in the community cohorts.** *J Rheumatol* 2011, **38**:135-141.
23. Harrold LR, Saag KG, Yood RA, Mikuls TR, Andrade SE, Fouayzi H, Davis J, Chan KA, Raebel MA, Von Worley A, Platt R: **Validity of gout diagnoses in administrative data.** *Arthritis Rheum* 2007, **57**:103-108.
24. Roddy E, Doherty M: **Epidemiology of gout.** *Arthritis Res Ther* 2010, **12**:223.
25. Hagos Y, Stein D, Ugele B, Burckhardt G, Bahn A: **Human renal organic anion transporter 4 operates as an asymmetric urate transporter.** *J Am Soc Nephrol* 2007, **18**:430-439.
26. Jutabha P, Anzai N, Kitamura K, Taniguchi A, Kaneko S, Yan K, Yamada H, Shimada H, Kimura T, Katada T, Fukutomi T, Tomita K, Urano W, Yamanaka H, Seki G, Fujita T, Moriyama Y, Yamada A, Uchida S, Wempe MF, Endou H, Sakurai H: **Human sodium phosphate transporter 4 (hNPT4/SLC17A3) as a common renal secretory pathway for drugs and urate.** *J Biol Chem* 2010, **285**:35123-35123.

doi:10.1186/ar3816

Cite this article as: Hollis-Moffatt *et al.*: The renal urate transporter SLC17A1 locus: confirmation of association with gout. *Arthritis Research & Therapy* 2012, **14**:R92.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

