



ORAL PRESENTATION

Open Access

Identification of novel genetic markers associated with the clinical phenotypes of systemic sclerosis through a genome wide association strategy

O Gorlova^{1†}, J M Martin^{2†}, B Rueda^{2†}, BPC Koeleman^{3†}, J Ying¹, M Teruel², L M Diaz-Gallo², J C Broen⁴, M C Vonk⁴, C P Simeon⁵, B Z Alizadeh⁶, MJH Coenen⁷, A E Voskuyl⁸, A J Schuerwegh⁹, PLCM van Riel⁴, M Vanthuyne¹⁰, R van 't Slot³, A Italiaander³, R A Ophoff³, N Hunzelmann¹¹, V Fonollosa⁵, N Ortego-Centeno¹², M A González-Gay¹³, F J García-Hernández¹⁴, M F González-Escribano¹⁵, P Airo¹⁶, J van Laar¹⁷, J Worthington¹⁸, R Hesselstrand¹⁹, V Smith²⁰, F De Keyser²⁰, F Houssiau¹⁰, M M Chee²¹, R Madhok²¹, P Shiels²¹, R Westhovens²², A Kreuter²³, E de Baere²⁴, T Witte²⁵, L Padyukov²⁶, A Nordin²⁶, R Scorza²⁷, C Lunardi²⁸, B A Lie²⁹, A M Hoffmann-Vold³⁰, P García de la Peña³¹, P Carreira^{13,32}, J Varga³³, M Hinchcliff³³, A T Lee³⁴, P Gourh³⁵, C I Amos¹, G Riemekasten³⁶, A Herrick¹⁸, L Beretta²⁷, C Fonseca³⁷, C P Denton³⁷, P K Gregersen³⁴, S Agarwal³⁵, S Assassi³⁵, F K Tan³⁵, F C Arnett^{35†}, TRDJ Radstake^{4†}, M D Mayes^{35†}, J Martin^{2*†}

From 5th European Workshop on Immune-Mediated Inflammatory Diseases Sitges-Barcelona, Spain. 1-3 December 2010

Purpose

The aim of this study was to determine the genetic components contributing to the different systemic sclerosis (SSc) clinical sub-phenotypes of limited (lcSSc) and diffuse (dcSSc) cutaneous involvement, and with the most common SSc-specific autoantibodies, anti-centromere (ACA) and anti-topoisomerase I (ATA) through a genome-wide association study (GWAS) and a large replication study stratified for these disease features.

Methods

In the discovery phase 2,296 SSc patients and 5,171 healthy controls were analyzed for genetic associations in lcSSc, dcSSc, ACA positive and ATA positive subgroups. The non-HLA SNPs associated with each subphenotype with a genomic control corrected *P* value lower than 1×10^{-5} , not previously associated with SSc, were selected for replication in 9 independent cohorts from the US and Europe comprising an additional 3,175 SSc patients and 4,971 controls. In addition, meta-analyses including all patients and healthy controls were conducted for each subphenotype.

Results

Three out of the 18 non-HLA SNPs selected for replication showed evidence of association. Meta-analysis including GWAS and replication cohorts showed a strong association of *IRF8* rs11642873 polymorphism ($P = 2.32 \times 10^{-12}$, OR = 0.75) and a suggestive but consistent association among populations of *GRB10* rs12540874 polymorphism ($P = 1.27 \times 10^{-6}$, OR = 1.15) with the lcSSc subtype of the disease. Furthermore, significant association of *SOX5* rs11047102 polymorphism ($P = 1.39 \times 10^{-7}$, OR = 1.36) with the ACA positive patients was detected. In the HLA region, specific patterns of SNPs associated with the ACA and ATA subgroups were observed, reflected by highly associated haplotype in the *HLA-DQB1* locus with ACA ($P = 1.79 \times 10^{-61}$), and in the *HLA-DPA1/B1* loci allelic combination with ATA ($P = 4.57 \times 10^{-76}$).

Conclusions

We have identified three new genes (*IRF8*, *GRB10*, and *SOX5*) associated with clinical manifestations of SSc, emphasizing the differential genetic component of each subphenotype of this disease. Within the HLA region, we have observed that *HLA-DQB1* and *HLA-DPA1/B1* associations with SSc are likely confined to specific auto-antibody positive patients.

† Contributed equally

²Instituto de Parasitología y Biomedicina López-Neyra, Consejo Superior de Investigaciones Científicas, Granada, Spain

Full list of author information is available at the end of the article

Author details

¹Dept. of Epidemiology, M.D. Anderson Cancer Center, Houston, Texas, USA. ²Instituto de Parasitología y Biomedicina López-Neyra, Consejo Superior de Investigaciones Científicas, Granada, Spain. ³Dept. of Medical Genetics, University Medical Center Utrecht, Utrecht, The Netherlands. ⁴Dept. of Rheumatology, Radboud University Nijmegen Medical Center, Nijmegen, The Netherlands. ⁵Servicio de Medicina Interna, Hospital Valle de Hebrón, Barcelona, Spain. ⁶University Medical Centre Groningen, Dept. of Epidemiology, Groningen, The Netherlands. ⁷Dept. of Human Genetics, Radboud University Nijmegen Medical Center, Nijmegen, The Netherlands. ⁸Dept. of Rheumatology, Vrije Universiteit (VU) Medical Centre, Amsterdam, The Netherlands. ⁹Dept. of Rheumatology, University of Leiden, Nijmegen, The Netherlands. ¹⁰Cliniques Universitaires Saint-Luc, Université catholique de Louvain, Brussels, Belgium. ¹¹Dept. of Dermatology, University of Cologne, Cologne, Germany. ¹²Servicio de Medicina Interna, Hospital Clínico Universitario, Granada, Spain. ¹³Servicio de Reumatología, Hospital Marqués de Valdecilla, Santander, Spain. ¹⁴Servicio de Medicina Interna, Hospital Virgen del Rocío, Sevilla, Spain. ¹⁵Servicio de Inmunología, Hospital Virgen del Rocío, Sevilla, Spain. ¹⁶Rheumatology Unit and Chair, Spedali Civili, Università degli Studi, Brescia, Italy. ¹⁷Institute of Cellular Medicine, Newcastle University, Newcastle Upon Tyne, UK. ¹⁸Dept. of Rheumatology and Epidemiology, University of Manchester, Manchester Academic Health Science Centre, Manchester, UK. ¹⁹Dept. of Clinical Sciences, Division of Rheumatology, Lund University, Lund, Sweden. ²⁰Ghent University, Ghent, Belgium. ²¹Centre for Rheumatic Diseases, Glasgow Royal Infirmary Glasgow, UK. ²²Katholieke Universiteit Leuven, Leuven, Belgium. ²³Dept. of Dermatology, Josefs-Hospital, Ruhr University Bochum, Germany. ²⁴Center for Medical Genetics, Ghent University Hospital, Ghent, Belgium. ²⁵Hannover Medical School, Hannover, Germany. ²⁶Center for Molecular Medicine, Karolinska Institutet, Stockholm, Sweden. ²⁷Referral Center for Systemic Autoimmune Diseases, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico and University of Milan, Milan, Italy. ²⁸Dept. of Clinical and Experimental Medicine, University Hospital, University of Verona, Italy. ²⁹Institute of Immunology, Oslo University Hospital Rikshospitalet, Oslo, Norway. ³⁰Dept. of Rheumatology, Rikshospitalet, Oslo University Hospital, Oslo, Norway. ³¹Servicio de Reumatología, Hospital Ramón y Cajal, Madrid, Spain. ³²Hospital 12 de Octubre, Madrid, Spain. ³³Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA. ³⁴Feinstein Institute of Medical Research, Manhasset, New York, USA. ³⁵The University of Texas Health Science Center–Houston, Houston, Texas, USA. ³⁶Dept. of Rheumatology and Clinical Immunology, Charité University Hospital, Berlin, Germany. ³⁷Centre for Rheumatology, Royal Free and University College School, London, UK.

Published: 25 November 2010

doi:10.1186/1479-5876-8-S1-O1

Cite this article as: Gorlova *et al.*: Identification of novel genetic markers associated with the clinical phenotypes of systemic sclerosis through a genome wide association strategy. *Journal of Translational Medicine* 2010 **8**(Suppl 1):O1.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

