



# Complete Genome Sequence of Buffalopox Virus

Babak Afrough,<sup>a</sup> Afia Zafar,<sup>b</sup> Rumina Hasan,<sup>b</sup> Roger Hewson<sup>a</sup>

<sup>a</sup>National Infection Service Research, Public Health England, Porton Down, United Kingdom

<sup>b</sup>Department of Microbiology, Aga Khan University, Karachi, Pakistan

**ABSTRACT** The first complete genome assembly of buffalopox virus isolate Karachi 2005, with a length of 195,630 bp, is presented here. Phylogenetic analysis shows the virus to cluster within *Vaccinia* species, and the genome contains 177 protein-coding sequences.

**B**uffalopox virus (BPXV), family *Poxviridae*, genus *Orthopoxvirus* (OPV), is related to monkeypox, horsepox, variola, and cowpox viruses (1). BPXV causes mild to severe disease in buffaloes and cattle, presenting as pock lesions. BPXV is a highly transmissible zoonosis, and the disease in the Indian subcontinent has persisted for over 30 years following the cessation of human vaccination against smallpox virus. Clinical symptoms include high fever, unilateral axillary lymphadenopathy, and edema, often associated with the eruption of lesions which are normally confined to the hands, forehead, face, buttocks, and legs (2, 3). It is not normally transmitted between humans; however, during 5 months in 2004 to 2005, this virus caused a nosocomial outbreak in association with burn units in Karachi, Pakistan (4). BPXV continues to sporadically reemerge across central Asia and the Indian subcontinent (5–7), and a deeper understanding of the viral genome will contribute to improvements in surveillance and disease control.

BPXV isolate Karachi 2005 was passaged once in Vero cells (ECACC accession number 84113001), and the infectious titer of this passage was determined at  $1.6 \times 10^5$  PFU/ml by a viral plaque assay. Total DNA was isolated using the DNeasy kit (Qiagen), purified using Agencourt AMPure XP beads (Beckman Coulter), and quantified using a double-stranded DNA (dsDNA) broad-range (BR) assay on a Qubit 3.0 fluorometer (Life Technologies). Two hundred nanograms of DNA was used to prepare an SQK-LSK108-1D sequencing library (Oxford Nanopore Technologies) without shearing. MinION Nanopore sequencing (R9.4 chemistry) generated 248,398 base-called reads (Albacore version 2.1.10), with a median length of 1,844 bp. A draft assembly was obtained using Canu version 1.6 (8) as a single 193,093-bp supercontig. All other contigs and unassembled reads were screened against a locally curated OPV database to ensure that no orphan fragments were missed during the *de novo* assembly step. The draft genome was assessed using Quast version 4.6.3 (9) and refined using Nanopolish version 0.8.5 (10), where 1.56% of the Nanopore reads mapped to BPXV.

A Nextera XT paired-end library (Illumina) was prepared using 100 ng of the same BPXV culture supernatant and was sequenced on a HiSeq 2500 platform. This produced 6,378,742 reads, with 2.29% mapping to BPXV. The reads were processed with Trimmomatic version 0.36 (11), mapped to the draft 193,093-bp Nanopore genome using BWA (12, 13), and sorted using SAMtools version 1.7 (14). Correction and consensus calling with Illumina reads were performed with Pilon version 1.22 (15) on the draft genome, producing a final sequence of 195,630 bp. Annotation was performed with Glimmer version 3.02 (16), GeneMarkS version 4.28 (17), and protein BLAST searches.

Long-read Nanopore sequencing permitted the assembly of the genome as a single supercontig. This included intact characteristic OPV terminal inverted repeat (TIR)

**Received** 16 April 2018 **Accepted** 18 April 2018 **Published** 24 May 2018

**Citation** Afrough B, Zafar A, Hasan R, Hewson R. 2018. Complete genome sequence of buffalopox virus. *Genome Announcements* 6:e00444-18. <https://doi.org/10.1128/genomeA.00444-18>.

© Crown copyright 2018. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Babak Afrough, [Babak.Afrough@phe.gov.uk](mailto:Babak.Afrough@phe.gov.uk).

regions that flank the BPXV genomic termini. Nanopore sequencing alone made it difficult to identify and annotate many of the protein-coding genes with confidence; thus, Illumina data were used to supplement the Nanopore sequencing. The BPXV genome presents an average G+C content of 33.4% and contains at least 177 protein-coding genes. Whole-genome phylogenetic characterization using maximum likelihood approaches indicates chorioallantois vaccinia virus Ankara, the parental strain of modified vaccinia virus Ankara, to be its closest relative.

**Accession number(s).** The annotated genome sequence is deposited in GenBank under the accession number [MG599038](#).

## ACKNOWLEDGMENTS

This work was supported by a grant-in-aid for New and Emerging Vector Borne and Zoonotic Disease from Public Health England.

BPXV is publicly available from the National Collection of Pathogenic Viruses (NCPV).

The findings, opinions, and views expressed herein belong to the authors and do not reflect an official position of the UK Department of Health.

## REFERENCES

1. Essbauer S, Pfeffer M, Meyer H. 2010. Zoonotic poxviruses. *Vet Microbiol* 140:229–236. <https://doi.org/10.1016/j.vetmic.2009.08.026>.
2. Ramanan C, Ghorpade A, Kalra SK, Mann S. 1996. Buffalopox. *Int J Dermatol* 35:128–130. <https://doi.org/10.1111/j.1365-4362.1996.tb03279.x>.
3. Riyesh T, Karuppusamy S, Bera BC, Barua S, Virmani N, Yadav S, Vaid RK, Anand T, Bansal M, Malik P, Pahuja I, Singh RK. 2014. Laboratory-acquired buffalopox virus infection, India. *Emerg Infect Dis* 20:324–326. <https://doi.org/10.3201/eid2002.130358>.
4. Zafar A, Swanepoel R, Hewson R, Nizam M, Ahmed A, Husain A, Grobelaar A, Bewley K, Mioulet V, Dowsett B, Easterbrook L, Hasan R. 2007. Nosocomial buffalopoxvirus infection, Karachi, Pakistan. *Emerg Infect Dis* 13:902–904. <https://doi.org/10.3201/eid1306.061068>.
5. Goyal T, Varshney A, Bakshi SK, Barua S, Bera BC, Singh RK. 2013. Buffalo pox outbreak with atypical features: a word of caution and need for early intervention! *Int J Dermatol* 52:1224–1230. <https://doi.org/10.1111/ijd.12120>.
6. Singh RK, Balamurugan V, Bhanuprakash V, Venkatesan G, Hosamani M. 2012. Emergence and reemergence of vaccinia-like viruses: global scenario and perspectives. *Indian J Virol* 23:1–11. <https://doi.org/10.1007/s13337-012-0068-1>.
7. Gurav YK, Raut CG, Yadav PD, Tandale BV, Sivaram A, Pore MD, Basu A, Mourya DT, Mishra AC. 2011. Buffalopox outbreak in humans and animals in Western Maharashtra, India. *Prev Vet Med* 100:242–247. <https://doi.org/10.1016/j.prevetmed.2011.03.008>.
8. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive *k*-mer weighting and repeat separation. *Genome Res* 27:722–736. <https://doi.org/10.1101/gr.215087.116>.
9. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
10. Loman NJ, Quick J, Simpson JT. 2015. A complete bacterial genome assembled *de novo* using only nanopore sequencing data. *Nat Methods* 12:733–735. <https://doi.org/10.1038/nmeth.3444>.
11. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
12. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>.
13. Li H, Ruan J, Durbin R. 2008. Mapping short DNA sequencing reads and calling variants using mapping quality scores. *Genome Res* 18:1851–1858. <https://doi.org/10.1101/gr.078212.108>.
14. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25:2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>.
15. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
16. Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. 1999. Improved microbial gene identification with GLIMMER. *Nucleic Acids Res* 27:4636–4641.
17. Besemer J, Lomsadze A, Borodovsky M. 2001. GeneMarkS: a self-training method for prediction of gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. *Nucleic Acids Res* 29:2607–2618.