



Received 5 October 2016

Accepted 10 October 2016

Edited by H. Stoeckli-Evans, University of
Neuchâtel, Switzerland**Keywords:** crystal structure; benzothiazine; allyl;
C—H... π interactions.**CCDC reference:** 1508990**Supporting information:** this article has
supporting information at journals.iucr.org/e

Crystal structure of methyl 1-allyl-4-methyl-1*H*-benzo[*c*][1,2]thiazine-3-carboxylate 2,2-dioxide

Liliana Azotla-Cruz,^a Svitlana Shishkina,^{b*} Igor Ukrainets,^c Irina Lijanov^a and
Natalya Likhanov^d^aNational Polytechnic Institute, CIITEC, Cerrada Cecati S/N, Colonia Santa Catarina de Azcapotzalco, CP 02250, Mexico, DF, Mexico, ^bSSI 'Institute for Single Crystals', National Academy of Sciences of Ukraine, 60 Lenina Avenue, Kharkiv 61001, Ukraine, ^cNational University of Pharmacy, 4 Blyukhera St., Kharkiv 61168, Ukraine, and ^dMexican Institute of Petroleum, Eje Central Lazaro Cardenas Norte 152, Col. San Bartolo Atepehuacan, 07730 Mexico, DF, Mexico.

*Correspondence e-mail: sveta@xray.isc.kharkov.com

In the title compound, C₁₄H₁₅NO₄S, the dihydrothiazine ring adopts a distorted sofa conformation with the S atom displaced from the mean plane through the N and C ring atoms by 0.767 (1) Å. The allyl substituent (C—C=C) is inclined to this mean plane by 78.5 (7)° and the acetate group [C(=O)—O—C] by 66.5 (3)°. In the crystal, molecules are linked by C—H... π interactions forming chains propagating along the *a*-axis direction.

1. Chemical context

Alkylation of nitrogen heterocycles, particularly those containing reactive exocyclic groups, always attracts attention with its ambiguity and dependence on a variety of factors. For example, esters of 4-hydroxy-2-oxo-1,2-dihydroquinoline-3-carboxylic acids are primarily alkylated exclusively at the 4-OH group (Ukrainets *et al.*, 1996). However, methyl 4-hydroxy-2,2-dioxo-1*H*-2 λ ⁶,1-benzothiazine-3-carboxylates that are structurally close to them easily form mixtures of isomeric 3-*C*- and 4-*O*-alkylation products under the same conditions (Ukrainets *et al.*, 2015). Consequently, it is quite difficult to predict their behaviour in the alkylation reactions of the esters of 4-methyl-2,2-dioxo-1*H*-2 λ ⁶,1-benzothiazine-3-carboxylic acids, and the determination of the true structure is essential. It has been found that methyl 4-methyl-2,2-dioxo-1*H*-2 λ ⁶,1-benzothiazine-3-carboxylate **1** in the K₂CO₃/DMSO system is rapidly alkylated with allyl bromide **2** by the cyclic nitrogen atom, with formation of the main product of the reaction studied *viz.* compound **3** (see Fig. 1).

2. Structural commentary

The molecular structure of the title compound, **3**, is illustrated in Fig. 2. The dihydrothiazine ring adopts a distorted sofa conformation: the puckering parameters (Zefirov *et al.*, 1990)






Figure 1
The synthesis of the title compound, **3**.

are: $S = 0.67$, $\Theta = 57.1^\circ$, $\Psi = 19.0^\circ$. Atom S1 deviates from the mean plane of the remaining atoms (N1/C1/C6–C8) by 0.767 (1) Å. The allyl substituent (C–C=C) is inclined to this mean plane by $78.5 (7)^\circ$ and the acetate group (O=C–O–C) by $66.5 (3)^\circ$. Atom N1 has a planar configuration, the sum of the bond angles being 359.1° .



The strong steric repulsion between methyl group at the C7 atom and the aromatic ring [there are short intramolecular contacts $H5 \cdots C11 = 2.63$ and $H11A \cdots C5 = 2.47$ Å in this fragment [the sum of the van der Waals radii (Zefirov, 1997) is 2.87 Å]] causes a disturbance of the conjugation between the π -systems of the aromatic ring and the C7=C8 double bond; the C5–C6–C7–C8 torsion angle is $-164.7 (4)^\circ$. The ester substituent is twisted relatively to the C7=C8 endocyclic double bond [the C7–C8–C9–O1 torsion angle is $46.0 (7)^\circ$], leading to its elongation: the C7=C8 bond length is 1.348 (6) Å as compared to the mean value of 1.326 Å (Bürgi & Dunitz, 1994). The methyl group of the ester substituent is




Figure 2
The molecular structure of compound **3**, with atom labelling. Displacement ellipsoids are drawn at the 50% probability level.

Table 1
Hydrogen-bond geometry (Å, °).

C_g is the centroid of the C1–C6 ring.

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$C12-H12A \cdots C_g^i$	0.97	2.95	3.576 (5)	123

Symmetry code: (i) $-x - \frac{1}{2}, y + \frac{3}{2}, z + \frac{1}{2}$.

located in the *ap*-position to the C8–C9 bond [C8–C9–O2–C10 = $-171.5 (5)^\circ$]. The allyl group is orthogonal to the benzothiazine fragment plane while the terminal double bond is synperiplanar to the N1–C12 bond [torsion angles C1–N1–C12–C13 and N1–C12–C13–C14 are $97.2 (5)$ and $3.5 (8)^\circ$, respectively]. The steric repulsion between the allyl substituent and the aromatic cycle (short intramolecular contacts $H2 \cdots C12 = 2.77$ Å and $H12A \cdots C2 = 2.83$ Å) results in the elongation of the C1–N1 bond [1.411 (5) Å], compared with the mean value of 1.371 Å (Bürgi & Dunitz, 1994).

3. Supramolecular features

In the crystal, molecules are linked by C–H $\cdots \pi$ interactions, forming chains propagating along the *a*-axis direction. (Table 1 and Fig. 3). There are no other significant intermolecular interactions in the crystal structure, despite the presence of a number of potential donor and acceptor atoms.

4. Database survey

A search of the Cambridge Structural Database (Version 5.37, update May 2016; Groom *et al.*, 2016) for 1*H*-benzo[*c*][1,2]thiazine 2,2-dioxide yielded 15 hits. These include the 4-hydroxy analogue of the title compound, *viz.* methyl 1-allyl-4-hydroxy-1*H*-benzo[*c*][1,2]thiazine-3-carboxylate 2,2-dioxide (MINJAW; Shishkina *et al.*, 2013). This compound crystallized with two molecules in the asymmetric unit. The conformation




Figure 3
A view along the *c* axis of the crystal packing of compound **3**. The C–H $\cdots \pi$ interactions are represented by dashed lines (see Table 1) and, for clarity, only H atom H12A (grey ball) is included.




Figure 4
The structural overlap of the two independent molecules of the 4-hydroxy analogue (MINJAW; Shishkina *et al.*, 2013), and the title compound **3**, shown in blue.

of the dihydrothiazine ring in both molecules resembles that in the title compound, which has a distorted sofa conformation. A view of the structural overlap of the three molecules is shown in Fig. 4.

5. Synthesis and crystallization

The synthesis of the title compound, **3**, is illustrated in Fig. 1. To a solution of 2.53 g (0.01 mol) of methyl 4-methyl-2,2-dioxo-1*H*-2λ⁶,1-benzothiazine-3-carboxylate, **1**, in 20 ml DMSO were added 2.07 g (0.015 mol) of K₂CO₃ and the mixture was stirred for 30 min. Allyl bromide (1.81 g, 0.015 mol) was then added and the mixture was stirred for a further 30 min at 298 K. It was then diluted with cold water and acidified with dilute HCl to pH 4. It was extracted with CH₂Cl₂ (3 × 10 ml). The organic extracts were combined and the solvent removed by distillation (at reduced pressure at the end). The residue was dissolved in 20 ml of hot methanol and filtered over charcoal. The resulting solution was then placed in a freezer (253 K) for 24 h, after which crystals of the title compound were harvested (yield 2.55 g, 87%; m.p. 360–362 K).

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. H atoms could all be located in difference Fourier maps. During refinement they were included in calculated positions and treated as riding: C–H = 0.93–0.97 Å with $U_{\text{iso}} = 1.5U_{\text{eq}}(\text{C-methyl})$ and $1.2U_{\text{eq}}(\text{C})$ for other H atoms.

Table 2
Experimental details.

Crystal data	
Chemical formula	C ₁₄ H ₁₅ NO ₄ S
M_r	293.33
Crystal system, space group	Orthorhombic, <i>Pna</i> 2 ₁
Temperature (K)	293
a, b, c (Å)	10.1970 (7), 18.6174 (12), 7.5136 (5)
V (Å ³)	1426.39 (16)
Z	4
Radiation type	Mo $K\alpha$
μ (mm ⁻¹)	0.24
Crystal size (mm)	0.20 × 0.10 × 0.02
Data collection	
Diffractometer	Agilent Xcalibur Sapphire3
Absorption correction	Multi-scan (<i>CrysAlis RED</i> ; Agilent, 2012)
$T_{\text{min}}, T_{\text{max}}$	0.706, 1.000
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	9346, 2425, 2111
R_{int} ($\sin \theta/\lambda$) _{max} (Å ⁻¹)	0.056 0.595
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.044, 0.120, 1.05
No. of reflections	2425
No. of parameters	181
No. of restraints	1
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ⁻³)	0.19, -0.20
Absolute structure	Flack x determined using 785 quotients $[(I^+) - (I^-)] / [(I^+) + (I^-)]$ (Parsons <i>et al.</i> , 2013)
Absolute structure parameter	0.09 (8)

Computer programs: *CrysAlis CCD* and *CrysAlis RED* (Agilent, 2012), *SHELXS2014* (Sheldrick, 2008), *SHELXTL* (Sheldrick, 2008), *Mercury* (Macrae *et al.*, 2008), *SHELXL2014* (Sheldrick, 2015), *PLATON* (Spek, 2009) and *publCIF* (Westrip, 2010).

References

- Agilent (2012). *CrysAlis CCD* and *CrysAlis RED*. Agilent Technologies, Yarnton, Oxfordshire, England.
- Bürgi, H.-B. & Dunitz, J. D. (1994). *Structure Correlation*, Vol. 2, pp. 767–784. Weinheim: VCH.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Shishkina, S. V., Ukrainets, I. V. & Petrushova, L. A. (2013). *Acta Cryst.* **E69**, o1698.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Ukrainets, I. V., Petrushova, L. A., Shishkina, S. V. & Sim, G. (2015). *Chem. Heterocycl. Compds*, **50**, 1741–1747.
- Ukrainets, I. V., Taran, S. G., Gorokhova, O. V., Goralcheva, I. V., Bezuglyi, P. A. & Turov, A. V. (1996). *Chem. Heterocycl. Compd.* **32**, 952–959.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zefirov, Yu. V. (1997). *Kristallografiya*, **42**, 936–958.
- Zefirov, N. S., Palyulin, V. A. & Dashevskaya, E. E. (1990). *J. Phys. Org. Chem.* **3**, 147–158.

supporting information

Acta Cryst. (2016). E72, 1574-1576 [https://doi.org/10.1107/S2056989016015978]

Crystal structure of methyl 1-allyl-4-methyl-1*H*-benzo[*c*][1,2]thiazine-3-carboxylate 2,2-dioxide

Liliana Azotla-Cruz, Svitlana Shishkina, Igor Ukrainets, Irina Lijanova and Natalya Likhanova

Computing details

Data collection: *CrysAlis CCD* (Agilent, 2012); cell refinement: *CrysAlis CCD* (Agilent, 2012); data reduction: *CrysAlis RED* (Agilent, 2012); program(s) used to solve structure: *SHELXS2014* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *SHELXTL* (Sheldrick, 2008) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2015), *PLATON* (Spek, 2009) and *pubCIF* (Westrip, 2010).

Methyl 1-allyl-4-methyl-1*H*-benzo[*c*][1,2]thiazine-3-carboxylate 2,2-dioxide

Crystal data

C₁₄H₁₅NO₄S

M_r = 293.33

Orthorhombic, *Pna*2₁

a = 10.1970 (7) Å

b = 18.6174 (12) Å

c = 7.5136 (5) Å

V = 1426.39 (16) Å³

Z = 4

F(000) = 616

D_x = 1.366 Mg m⁻³

Mo *Kα* radiation, λ = 0.71073 Å

Cell parameters from 2163 reflections

θ = 3.8–25.1°

μ = 0.24 mm⁻¹

T = 293 K

Plate, colourless

0.20 × 0.10 × 0.02 mm

Data collection

Agilent Xcalibur Sapphire3
diffractometer

Radiation source: Enhance (Mo) X-ray Source

Detector resolution: 16.1827 pixels mm⁻¹

ω-scans

Absorption correction: multi-scan

(*CrysAlis RED*; Agilent, 2012)

T_{min} = 0.706, *T_{max}* = 1.000

9346 measured reflections

2425 independent reflections

2111 reflections with *I* > 2σ(*I*)

R_{int} = 0.056

θ_{max} = 25.0°, θ_{min} = 3.0°

h = -12→10

k = -22→22

l = -7→8

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.044

wR(*F*²) = 0.120

S = 1.05

2425 reflections

181 parameters

1 restraint

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F_o*²) + (0.0618*P*)² + 0.1291*P*]

where *P* = (*F_o*² + 2*F_c*²)/3

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.19 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$$

Absolute structure: Flack x determined using
785 quotients $[(I^-)-(I)]/[(I^+)+(I)]$ (Parsons *et al.*,
2013)

Absolute structure parameter: 0.09 (8)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.88243 (9)	0.65549 (5)	0.26457 (17)	0.0457 (3)
O1	0.6502 (4)	0.5240 (2)	0.4865 (7)	0.0975 (14)
O2	0.7649 (4)	0.51032 (18)	0.2347 (6)	0.0801 (11)
O3	0.7789 (3)	0.66144 (16)	0.1376 (5)	0.0645 (9)
O4	1.0109 (3)	0.63899 (17)	0.2033 (4)	0.0590 (9)
N1	0.8881 (3)	0.73110 (17)	0.3775 (5)	0.0457 (8)
C1	0.9692 (4)	0.7307 (2)	0.5295 (6)	0.0443 (9)
C2	1.0333 (4)	0.7933 (3)	0.5818 (7)	0.0559 (11)
H2	1.0205	0.8355	0.5180	0.067*
C3	1.1146 (4)	0.7932 (3)	0.7260 (7)	0.0652 (14)
H3	1.1557	0.8355	0.7611	0.078*
C4	1.1362 (5)	0.7308 (3)	0.8195 (7)	0.0722 (15)
H4	1.1956	0.7305	0.9135	0.087*
C5	1.0705 (5)	0.6690 (3)	0.7747 (8)	0.0605 (11)
H5	1.0837	0.6277	0.8419	0.073*
C6	0.9842 (4)	0.6670 (2)	0.6302 (6)	0.0461 (10)
C7	0.9017 (4)	0.6034 (2)	0.5943 (6)	0.0490 (10)
C8	0.8424 (4)	0.5951 (2)	0.4353 (6)	0.0497 (11)
C9	0.7425 (5)	0.5388 (2)	0.3921 (8)	0.0623 (13)
C10	0.6651 (7)	0.4617 (4)	0.1624 (12)	0.110 (3)
H10A	0.6927	0.4446	0.0478	0.165*
H10B	0.6537	0.4217	0.2415	0.165*
H10C	0.5835	0.4870	0.1506	0.165*
C11	0.8826 (5)	0.5494 (3)	0.7415 (9)	0.0715 (15)
H11A	0.9319	0.5640	0.8442	0.107*
H11B	0.7913	0.5467	0.7717	0.107*
H11C	0.9126	0.5031	0.7025	0.107*
C12	0.7970 (4)	0.7897 (2)	0.3411 (7)	0.0547 (11)
H12A	0.7753	0.8127	0.4530	0.066*
H12B	0.7168	0.7693	0.2936	0.066*
C13	0.8440 (5)	0.8458 (2)	0.2154 (8)	0.0661 (15)
H13	0.7888	0.8848	0.1961	0.079*
C14	0.9528 (6)	0.8455 (3)	0.1316 (9)	0.0829 (18)
H14A	1.0112	0.8076	0.1466	0.099*
H14B	0.9737	0.8832	0.0555	0.099*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0486 (5)	0.0440 (5)	0.0446 (5)	-0.0024 (4)	-0.0040 (5)	-0.0029 (5)
O1	0.087 (3)	0.089 (3)	0.116 (4)	-0.038 (2)	0.020 (3)	-0.007 (3)
O2	0.084 (2)	0.063 (2)	0.094 (3)	-0.0162 (18)	-0.005 (2)	-0.028 (2)
O3	0.070 (2)	0.062 (2)	0.062 (2)	-0.0037 (16)	-0.0255 (17)	-0.0033 (16)
O4	0.0587 (18)	0.0636 (19)	0.0546 (19)	-0.0032 (15)	0.0108 (14)	-0.0081 (14)
N1	0.0528 (19)	0.0364 (17)	0.048 (2)	-0.0005 (14)	-0.0052 (15)	-0.0030 (17)
C1	0.044 (2)	0.046 (2)	0.043 (2)	-0.0011 (18)	0.0033 (18)	-0.0031 (18)
C2	0.061 (3)	0.053 (3)	0.053 (3)	-0.008 (2)	0.004 (2)	-0.005 (2)
C3	0.070 (3)	0.075 (3)	0.051 (3)	-0.020 (3)	0.005 (2)	-0.015 (2)
C4	0.075 (3)	0.096 (4)	0.046 (3)	-0.009 (3)	-0.013 (2)	-0.008 (3)
C5	0.070 (3)	0.067 (3)	0.045 (2)	0.005 (2)	-0.009 (3)	0.001 (3)
C6	0.049 (2)	0.047 (2)	0.042 (2)	0.0024 (18)	0.0034 (18)	-0.0002 (18)
C7	0.055 (2)	0.042 (2)	0.050 (3)	0.0070 (18)	0.003 (2)	0.0025 (19)
C8	0.051 (2)	0.038 (2)	0.060 (3)	-0.0031 (18)	0.003 (2)	-0.0031 (19)
C9	0.062 (3)	0.046 (2)	0.079 (4)	-0.009 (2)	0.000 (3)	0.000 (2)
C10	0.112 (5)	0.070 (4)	0.149 (7)	-0.028 (4)	-0.040 (5)	-0.028 (4)
C11	0.093 (4)	0.053 (3)	0.068 (4)	-0.001 (2)	-0.002 (3)	0.014 (3)
C12	0.054 (2)	0.046 (2)	0.064 (3)	0.0025 (19)	-0.001 (2)	0.001 (2)
C13	0.067 (3)	0.051 (3)	0.081 (4)	0.003 (2)	0.000 (3)	0.011 (2)
C14	0.076 (4)	0.090 (4)	0.082 (4)	-0.001 (3)	-0.001 (3)	0.026 (3)

Geometric parameters (Å, °)

S1—O4	1.422 (3)	C5—H5	0.9300
S1—O3	1.427 (3)	C6—C7	1.477 (6)
S1—N1	1.645 (3)	C7—C8	1.348 (6)
S1—C8	1.754 (4)	C7—C11	1.508 (7)
O1—C9	1.210 (6)	C8—C9	1.497 (6)
O2—C9	1.316 (6)	C10—H10A	0.9600
O2—C10	1.466 (6)	C10—H10B	0.9600
N1—C1	1.411 (5)	C10—H10C	0.9600
N1—C12	1.458 (5)	C11—H11A	0.9600
C1—C2	1.392 (6)	C11—H11B	0.9600
C1—C6	1.415 (6)	C11—H11C	0.9600
C2—C3	1.364 (7)	C12—C13	1.487 (7)
C2—H2	0.9300	C12—H12A	0.9700
C3—C4	1.376 (8)	C12—H12B	0.9700
C3—H3	0.9300	C13—C14	1.276 (8)
C4—C5	1.373 (7)	C13—H13	0.9300
C4—H4	0.9300	C14—H14A	0.9300
C5—C6	1.398 (6)	C14—H14B	0.9300
O4—S1—O3	118.8 (2)	C7—C8—C9	125.3 (4)
O4—S1—N1	108.66 (18)	C7—C8—S1	118.0 (3)
O3—S1—N1	107.72 (19)	C9—C8—S1	116.6 (4)

O4—S1—C8	108.2 (2)	O1—C9—O2	124.8 (5)
O3—S1—C8	111.5 (2)	O1—C9—C8	124.1 (5)
N1—S1—C8	100.3 (2)	O2—C9—C8	111.0 (4)
C9—O2—C10	117.4 (5)	O2—C10—H10A	109.5
C1—N1—C12	122.0 (3)	O2—C10—H10B	109.5
C1—N1—S1	115.7 (3)	H10A—C10—H10B	109.5
C12—N1—S1	121.4 (3)	O2—C10—H10C	109.5
C2—C1—N1	120.0 (4)	H10A—C10—H10C	109.5
C2—C1—C6	120.0 (4)	H10B—C10—H10C	109.5
N1—C1—C6	120.0 (3)	C7—C11—H11A	109.5
C3—C2—C1	120.5 (5)	C7—C11—H11B	109.5
C3—C2—H2	119.7	H11A—C11—H11B	109.5
C1—C2—H2	119.7	C7—C11—H11C	109.5
C2—C3—C4	120.3 (5)	H11A—C11—H11C	109.5
C2—C3—H3	119.9	H11B—C11—H11C	109.5
C4—C3—H3	119.9	N1—C12—C13	116.1 (4)
C5—C4—C3	120.3 (5)	N1—C12—H12A	108.3
C5—C4—H4	119.9	C13—C12—H12A	108.3
C3—C4—H4	119.9	N1—C12—H12B	108.3
C4—C5—C6	121.4 (5)	C13—C12—H12B	108.3
C4—C5—H5	119.3	H12A—C12—H12B	107.4
C6—C5—H5	119.3	C14—C13—C12	126.2 (5)
C5—C6—C1	117.4 (4)	C14—C13—H13	116.9
C5—C6—C7	121.5 (4)	C12—C13—H13	116.9
C1—C6—C7	120.8 (4)	C13—C14—H14A	120.0
C8—C7—C6	120.7 (4)	C13—C14—H14B	120.0
C8—C7—C11	121.0 (4)	H14A—C14—H14B	120.0
C6—C7—C11	118.3 (4)		
O4—S1—N1—C1	-60.7 (3)	C1—C6—C7—C8	21.7 (6)
O3—S1—N1—C1	169.5 (3)	C5—C6—C7—C11	17.0 (6)
C8—S1—N1—C1	52.7 (3)	C1—C6—C7—C11	-156.7 (4)
O4—S1—N1—C12	130.1 (3)	C6—C7—C8—C9	-171.1 (4)
O3—S1—N1—C12	0.2 (4)	C11—C7—C8—C9	7.3 (7)
C8—S1—N1—C12	-116.5 (3)	C6—C7—C8—S1	7.8 (6)
C12—N1—C1—C2	-42.8 (5)	C11—C7—C8—S1	-173.9 (4)
S1—N1—C1—C2	147.9 (3)	O4—S1—C8—C7	72.8 (4)
C12—N1—C1—C6	136.3 (4)	O3—S1—C8—C7	-154.8 (3)
S1—N1—C1—C6	-32.9 (5)	N1—S1—C8—C7	-40.9 (4)
N1—C1—C2—C3	-178.4 (4)	O4—S1—C8—C9	-108.2 (4)
C6—C1—C2—C3	2.5 (6)	O3—S1—C8—C9	24.2 (4)
C1—C2—C3—C4	1.2 (7)	N1—S1—C8—C9	138.1 (3)
C2—C3—C4—C5	-3.6 (8)	C10—O2—C9—O1	5.4 (8)
C3—C4—C5—C6	2.4 (8)	C10—O2—C9—C8	-171.5 (5)
C4—C5—C6—C1	1.2 (7)	C7—C8—C9—O1	46.0 (7)
C4—C5—C6—C7	-172.6 (5)	S1—C8—C9—O1	-132.9 (5)
C2—C1—C6—C5	-3.6 (6)	C7—C8—C9—O2	-137.1 (5)
N1—C1—C6—C5	177.2 (4)	S1—C8—C9—O2	44.0 (5)

C2—C1—C6—C7	170.3 (4)	C1—N1—C12—C13	97.2 (5)
N1—C1—C6—C7	-8.9 (6)	S1—N1—C12—C13	-94.2 (4)
C5—C6—C7—C8	-164.7 (4)	N1—C12—C13—C14	3.5 (8)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg is the centroid of the C1—C6 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C12—H12A \cdots Cg ⁱ	0.97	2.95	3.576 (5)	123

Symmetry code: (i) $-x-1/2, y+3/2, z+1/2$.