



ORIGINAL RESEARCH

Open Access

Aetiology of auditory dysfunction in amusia: a systematic review

Daniel AJ Casey

Abstract

Background: Amusia, a music-specific agnosia, is a disorder of pitch interval analysis and pitch direction change recognition which results in a deficit in musical ability. The full range of aetiological factors which cause this condition is unknown, as is each cause's frequency. The objective of this study was to identify all causes of amusia, and to measure each of their frequencies.

Methods: Design: systematic review was conducted by search of multiple databases for articles related to the aetiology of amusic auditory dysfunction. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines for reporting of systematic reviews were followed, utilizing the PRISMA checklist and PRISMA flowchart methodology. Setting: Retrospective medical database review. Main outcome measures: evidence yielded from the systematic review process.

Results: The initial search protocol identified 5723 articles. Application of a classification review filter and exclusion of irrelevant or duplicates led to the initial identification of 56 relevant studies which detailed 301 patients. However, these studies were of poor quality. Because of this, synthesis and statistical analysis were not appropriate.

Conclusion: Although initially a large number of relevant studies were identified, and might point in future to potential diagnostic categories, it was not appropriate to synthesise and analyse them due to poor quality, considerable heterogeneity and small numbers. This suggests that large, high quality studies focussed directly on understanding the aetiology of amusia are required.

Keywords: Amusia, Aetiology, Auditory agnosia, Otology, Neurology

Background

Amusia is a music-specific auditory agnosia consisting of neurological deficit in musical ability [1]. Knoblauch's definition is 'the loss of a musical ability, such as the comprehension of music, the production of music, or the ability to read or write musical notation' [2].

However, a grey area lies between 'amusia' and auditory agnosias which impair auditory function related to pitch processing, interval analysis, timbre, rhythm or the emotional components of music but with sparing of gross musical ability [3]. This study will focus on studies in which there is a clear loss of musical ability and thus satisfy Knoblauch's definition.

Fry estimates that 4.2% of the UK adult population may be amusic [4]. As music and melody form a fundamental part of human experience, amusia is associated

with significant distress, especially for musical professionals. Removal of a stimulus which in some instances can activate neuroanatomical regions associated with intense pleasure can represent a severe handicap [5].

Thus, a systematic examination of the aetiology of amusia may be of considerable importance in forming a differential diagnosis in the clinic and in informing prevention strategies. Understanding the aetiological factors behind amusia can also shed light on musical auditory function and aid in deficit prediction. The neural basis of music is a matter of intense investigation.

However, no such aetiological categorization has yet been conducted. Here, a systematic review of the causes of amusia is performed.

Method

This systematic review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses

Correspondence: daniel.casey@merton.ox.ac.uk
Merton College, University of Oxford, Oxford OX1 4JD, UK

(PRISMA) guidelines for reporting of systematic reviews, utilizing the PRISMA checklist and PRISMA flowchart methodology [6].

Initial search protocol

Systematic review was conducted by search of multiple databases for articles related to the aetiology of amusic auditory dysfunction (Medline and The Cochrane Library). Key Terms used to identify the concept of amusic auditory dysfunction included amusia (sensory and motor), dysmusia, tone deaf(ness), tune deaf(ness), pitch processing deficit and note deaf(ness). Keywords used to identify aetiological frequency included aetiology, etiology, aetiologic(al) factor(s), aetiol*, etiol*, pathophysiology identification, risk factor(s) and cause(s). Aetiological keywords were combined with those specific to amusic auditory dysfunction. A secondary search was then conducted, and the grey literature was also searched.

Filter procedure

Initial search using the amusic key-terms, with and without aetiological key-term modifiers, yielded 5723 articles for assessment. Articles to form part of the systematic review were then filtered based on a classification review filter. Studies to be included had to be directly relevant to amusia, contain a definition of amusia consistent with the current literature (i.e. music specific auditory agnosia), and provide a full aetiological explanation for all subjects with accompanying diagnostic evidence.

After elimination of irrelevant articles and application of the above criteria, results were narrowed to 56 papers containing 301 patients ranging from 1878 to 2012, including articles in English, German, Spanish, French,

Italian, Russian and Japanese (non-English language articles were translated into English).

However, 40 of these 56 were single case reports; the remaining studies also contained small patient numbers (and/or had amusia as a secondary concern within the paper). This meant that the quality of evidence found was very poor. Due to this, no studies found were appropriate for synthesis and analysis.

Results

The PRISMA flow diagram (Figure 1) summarizes the article selection process. The initial search protocol identified 5723 articles. Application of the classification review filter and exclusion of irrelevant or duplicates led to the initial identification of 56 relevant studies [7-62]. These studies contained 301 patients. Articles had been published from 1878 to 2012 and included papers in English, German, Spanish, Italian, French, Russian and Japanese (non-English language articles were translated into English).

However, none were suitable for synthesis and further analysis because they were poor quality with regards to small numbers. In addition, considerable heterogeneity precluded synthesis.

Discussion

Although initially a large number of relevant studies were identified, and might point in future to potential diagnostic categories, it was not possible to synthesise them due to poor quality, considerable heterogeneity and small numbers.

40 of the 56 initially identified studies were single case reports: clearly, drawing conclusions from a synthesis of these would not be appropriate. The remaining studies




Figure 1 Article selection process following PRISMA guidelines.

Potential future diagnostic categories: classification of amusia aetiologies in 301 patients.




Figure 2 Potential future diagnostic categories: classification of amusia aetiologies in 301 patients. This figure is simply a record of the aetiologies identified in studies which were not suitable for analysis or synthesis but nonetheless contained amusic patients. Speculatively, some of these aetiological categories might be important when a full systematic review is possible in the future.

also contained relatively small patient numbers. In addition, these studies were extremely heterogeneous. Studies came from a wide range of time periods, with different reporting methods.

On this basis, to answer the question of amusia aetiology, it is necessary in the future to design and implement large, high quality studies with agreed diagnostic criteria.

Implications for research and future hypotheses

Understanding and quantifying amusia aetiology would be useful for assessment of this condition in a clinical setting. This paper suggests that this will not be possible until relevant high quality evidence is produced.

Before this evidence becomes available, it is interesting to think of the possible diagnostic categories which might emerge – classification of the 56 'low quality' (with respect to performing a systematic review) studies identified above suggests the following possibilities: vascular, congenital, iatrogenic, neoplastic, epilepsy-associated, degenerative, traumatic and idiopathic (Figure 2). Of course, which of these possibilities, if any, are relevant, awaits further investigation.

We might hypothesize that some cause of neural damage to key brain regions is required for the onset of amusia. Although it is inappropriate to draw any type of quantitative conclusion from the initial 56 studies identified above, categorization of the neuroanatomical areas involved suggests a host of possible key areas

Possible future diagnostic categories: neuroanatomical regions affected in 301 amusic patients.




Figure 3 Possible future diagnostic categories: neuroanatomical sites involved in 301 amusic cases.

(Figure 3). However, this is entirely speculative, and many of these possibilities may not be relevant. Hypothetically, the underlying commonality of any aetiological cause of amusia might be temporal lobe damage to a putative 'pitch centre' (or, at least, a key region involved in pitch processing) in lateral Heschl's gyrus (HG) [63].

Conclusion

In conclusion, due to the infrequent and irregular nature of the evidence related to amusia aetiology, all relevant studies suffer from small numbers and considerable heterogeneity. This, in addition to the other problems outlined above, suggests that large, high quality studies focussed directly on understanding the aetiology of amusia are required before a synthesis and analysis can take place.

Competing interests

The author declare that he has no competing interests.

Authors' contributions

DC was the sole author.

Received: 10 December 2012 Accepted: 19 April 2013

Published: 24 April 2013

References

- Pearce JM: Selected observations on amusia. *Eur Neurol* 2005, 54(3):145–148.
- Knoblauch A: Über Störungen der musikalischen Leistungsfähigkeit infolge von Gehirnläsionen. *Deutsch Arch Klin Med* 1888, 43:331–352.
- Stewart L, von Kriegstein K, Warren JD, Griffiths TD: Music and the brain: disorders of musical listening. *Brain* 2006, 129(Pt 10):2533–2553.
- Kalmus H, Fry DB: On tune deafness (Dysmelodia): frequency, development, genetics and musical background. *Ann Hum Genet* 1980, 43(4):369–382.
- Blood AJ, Zatorre RJ: Intensely pleasurable responses to music correlate with activity in brain regions implicated in reward and emotion. *Proc Natl Acad Sci USA* 2001, 98(20):11818–11823.
- Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group: Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009, 339:b2535.
- Allen G: Note-deafness. *Mind* 1878, 10:157–167.
- Anonymous: Amusia. *Br Med J* 1894, 2(1773):1441–1442.
- Smith PB: Auditory Aphasia with Amusia. *Br Med J* 1897, 1(1892):842–843.
- Kohl GF, Tschabitscher H: A case of amusia. *Wien Z Nervenheilkd Grenzgeb* 1953, 6(2–3):219–230.
- Scholer W: A case of pure word deafness with amusia. *Monatsschr Ohrenheilkd Laryngorhino* 1955, 8(2):106–109.
- Fukuda K, Hasegawa T: A case of visual agnosia with amusia (author's transl). *Seishin Shinkeigaku Zasshi* 1976, 78(5):417–430.
- Mazzucchi A, Marchini C, Budai R, Parma M: A case of receptive amusia with prominent timbre perception defect. *J Neurol Neurosurg Psychiatry* 1982, 45(7):644–647.
- Botez M, Wertheim N: Expressive aphasia and amusia following right frontal lesion in a right-handed man. *Brain* 1959, 82:186–202.
- Wertheim N, Botez M: Receptive amusia: a clinical analysis. *Brain* 1961, 84:19–30.
- Ulrich G, The syndrome of auditory agnosia: A case report and an attempt of neuropsychological qualification (author's transl). *Arch Psychiat Nervenkr* 1977, 224(3):221–233.
- Haguenauer JP, Schott B, Michel F, Dubreuil C, Romanet P: Three case histories of cortical and sub-cortical auditory lesions. Audiological and tomodensimetric confrontations (author's transl). *Ann Otolaryngol Chir Cervicofac* 1979, 96(3):185–196.
- Traugott NN, Beskadarov AV, Vasserman LI, Galunov VI, Dorofeeva SA: Clinico-experimental study of auditory-speech agnosia (case with anatomo-histologic verification. *Zh Nevropatol Psichiatr Im S S Korsakova* 1980, 80(12):1790–1798.
- Mavlov L: Amusia due to rhythm agnosia in a musician with left hemisphere damage: a non-auditory supramodal defect. *Cortex* 1980, 16(2):331–338.
- Buscaino GA, Grossi D, Orefice G: Pure verbal deafness with amusia. Apropos of a case. *Acta Neurol (Napoli)* 1981, 3(1):260–265. Italian.
- McFarland HR, Fortin D: Amusia due to right temporoparietal infarct. *Arch Neurol* 1982, 39(11):725–727.
- Takeda K, Bandou M, Nishimura Y: Motor amusia following a right temporal lobe hemorrhage—a case report. *Rinsho Shinkeigaku* 1990, 30(1):78–83. Review.
- Sidtis JJ, Feldmann E: Transient ischemic attacks presenting with a loss of pitch perception. *Cortex* 1990, 26(3):469–471.
- Griffiths TD, Rees A, Witton C, Cross PM, Shakir RA, Green GG: Spatial and temporal auditory processing deficits following right hemisphere infarction. A psychophysical study. *Brain* 1997, 120(Pt 5):785–794.
- Peretz I, Belleville S, Fontaine S: Dissociations between music and language functions after cerebral resection: A new case of amusia without aphasia. *Can J Exp Psychol* 1997, 51(4):354–368.
- Estañol B, Méndez A: A case of cortical amusia in a patient with musical ability. *Rev Neurol* 1998, 26(152):612–615.
- Schön D, Semenza C, Denes G: Naming of musical notes: a selective deficit in one musical clef. *Cortex* 2001, 37(3):407–421.
- Kohlmetz C, Altenmüller E, Schuppert M, Wieringa BM, Münte TF: Deficit in automatic sound-change detection may underlie some music perception deficits after acute hemispheric stroke. *Neuropsychologia* 2001, 39(11):1121–1124.
- McChesney-Atkins S, Davies KG, Montouris GD, Silver JT, Menkes DL: Amusia after right frontal resection for epilepsy with singing seizures: case report and review of the literature. *Epilepsy Behav* 2003, 4(3):343–347.
- Di Pietro M, Laganaro M, Leemann B, Schnider A: Receptive amusia: temporal auditory processing deficit in a professional musician following a left temporo-parietal lesion. *Neuropsychologia* 2004, 42(7):868–877.
- Satoh M, Takeda K, Murakami Y, Onouchi K, Inoue K, Kuzuhara S: A case of amusia caused by the infarction of anterior portion of bilateral temporal lobes. *Cortex* 2005, 41(1):77–83.
- Satoh M, Takeda K, Kuzuhara S: A case of auditory agnosia with impairment of perception and expression of music: cognitive processing of tonality. *Eur Neurol* 2007, 58(2):70–77.
- Lechevalier B, Rumbach L, Platel H, Lambert J: Pure amusia revealing an ischaemic lesion of right temporal planum. Participation of the right temporal lobe in perception of music. *Bull Acad Natl Med* 2006, 190(8):1697–1709. discussion.
- Peretz I, Cummings S, Dubé MP: The genetics of congenital amusia (tone deafness): a family-aggregation study. *Am J Hum Genet* 2007, 81(3):582–588. Epub 2007 Jul 20.
- Hyde KL, Lerch JP, Zatorre RJ, Griffiths TD, Evans AC, Peretz I: Cortical thickness in congenital amusia: when less is better than more. *J Neurosci* 2007, 27(47):13028–13032.
- Buklina SB, Skvortsova VB: Amusia and its topic specification. *Zh Nevropatol Psichiatr Im S S Korsakova* 2007, 107(9):4–10.
- Särkämö T, Tervaniemi M, Soinila S, Autti T, Silvennoinen HM, Laine M, Hietanen M: Cognitive deficits associated with acquired amusia after stroke: a neuropsychological follow-up study. *Neuropsychologia* 2009, 47(12):2642–2651. Epub 2009 Jun 3.
- Hyde KL, Zatorre RJ, Peretz I: Functional MRI evidence of an abnormal neural network for pitch processing in congenital amusia. *Cereb Cortex* 2011, 21(2):292–299. Epub 2010 May 21.
- Barquiero S, Gomez-Tortosa E, Baron M, Rabano A, Munoz DG, Jimenez-Escrig A: Amusia as an early manifestation of frontotemporal dementia caused by a novel progranulin mutation. *J Neurol* 2010, 257(3):475–477.
- Phillips-Silver J, Toivainen P, Gosselin N, Piché O, Nozaradan S, Palmer C, Peretz I: Born to dance but beat deaf: a new form of congenital amusia. *Neuropsychologia* 2011, 49(5):961–969. Epub 2011 Feb 21.
- Reed CL, Cahn SJ, Cory C, Szaflarski JP: Impaired perception of harmonic complexity in congenital amusia: a case study. *Cogn Neuropsychol* 2011, 28(5):305–321. Epub 2012 Jan 17.
- Confavreux C, Croisile B, Garassus P, Aimard G, Trillet M: Progressive amusia and aprosody. *Arch Neurol* 1992, 49(9):971–976.
- Piccirilli M, Sciarra T, Luzzi S: Modularity of music: evidence from a case of pure amusia. *J Neurol Neurosurg Psychiatry* 2000, 69(4):541–545.

44. Ayotte J, Peretz I, Hyde K: Congenital amusia: a group study of adults afflicted with a music-specific disorder. *Brain* 2002, 125(Pt 2):238–251.
45. Wilson SJ, Pressing JL, Wales RJ: Modelling rhythmic function in a musician post-stroke. *Neuropsychologia* 2002, 40(8):1494–1505.
46. Bautista RE, Ciampetti MZ: Expressive aprosody and amusia as a manifestation of right hemisphere seizures. *Epilepsia* 2003, 44(3):466–467.
47. Russell SM, Golfinos JG: Amusia following resection of a Heschl gyrus glioma. Case report. *J Neurosurg* 2003, 98(5):1109–1112.
48. Murayama J, Kashiwagi T, Kashiwagi A, Mimura M: Impaired pitch production and preserved rhythm production in a right brain-damaged patient with amusia. *Brain Cogn* 2004, 56(1):36–42.
49. Terao Y, Mizuno T, Shindoh M, Sakurai Y, Ugawa Y, Kobayashi S, Nagai C, Furubayashi T, Arai N, Okabe S, Mochizuki H, Hanajima R, Tsuji S: Vocal amusia in a professional tango singer due to a right superior temporal cortex infarction. *Neuropsychologia* 2006, 44(3):479–488.
50. Peretz I, Brattico E, Tervaniemi M: Abnormal electrical brain responses to pitch in congenital amusia. *Ann Neurol* 2005, 58(3):478–482.
51. Hyde KL, Zatorre RJ, Griffiths TD, Lerch JP, Peretz I: Morphometry of the amusic brain: a two-site study. *Brain* 2006, 129(Pt 10):2562–2570.
52. Hoffmann M: Isolated right temporal lobe stroke patients present with Geschwind Gastaut syndrome, frontal network syndrome and delusional misidentification syndromes. *Behav Neurol* 2008, 20(3):83–89.
53. Vanstone AD, Cuddy LL, Duffin JM, Alexander E: Exceptional preservation of memory for tunes and lyrics: case studies of amusia, profound deafness, and Alzheimer's disease. *Ann N Y Acad Sci* 2009, 1169:291–294.
54. Tillmann B, Burnham D, Nguyen S, Grimaud N, Gosselin N, Peretz I: Congenital Amusia (or Tone-Deafness) Interferes with Pitch Processing in Tone Languages. *Front Psychol* 2011, 2:120.
55. Hochman MS, Abrams KJ: Amusia for Pitch Caused by Right Middle Cerebral Artery Infarc. *J Stroke Cerebrovasc Dis* 2012. <http://dx.doi.org/10.1016/j.jstrokecerebrovasdis.2012.06.016>.
56. Geshwind N: The brain of a learning-disabled individual. *Ann Dyslexia* 1984, 34:319–327.
57. Lebrun MA, Moreau P, McNallt-Gagnon A, Mignault Goulet G, Peretz I: Congenital amusia in childhood: a case study. *Cortex* 2012, 48(6):683–688.
58. Eustache F, Lechevalier B, Viader F, Lambert J: Identification and discrimination disorders in auditory perception: a report on two cases. *Neuropsychologia* 1990, 28(3):257–270.
59. Hattiangadi N, Pillion JP, Slomine B, Christensen J, Trovato MK, Speedie LJ: Characteristics of auditory agnosia in a child with severe traumatic brain injury: a case report. *Brain Lang* 2005, 92(1):12–25.
60. Johannes S, Jöbges ME, Dengler R, Münte TF: Cortical auditory disorders: a case of non-verbal disturbances assessed with event-related brain potentials. *Behav Neurol* 1998, 11(1):55–73.
61. Mazzoni M, Moretti P, Pardossi L, Vista M, Muratorio A, Puglioli M: A case of music imperception. *J Neurol Neurosurg Psychiatry* 1993, 56(3):322.
62. Peretz I, Kolinsky R, Tramo M, Labrecque R, Hublet C, Demeurisse G, Belleville S: Functional dissociations following bilateral lesions of auditory cortex. *Brain* 1994, 117:1283–1301.
63. Penagos H, Melcher JR, Oxenham AJ: A neural representation of pitch salience in nonprimary human auditory cortex revealed with functional magnetic resonance imaging. *J Neurosci* 2004, 24:6810–6815.

doi:10.1186/1755-7682-6-16

Cite this article as: Casey: Aetiology of auditory dysfunction in amusia: a systematic review. *International Archives of Medicine* 2013 6:16.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

