

Draft Genome Sequences of *Clostridium tyrobutyricum* Strains FAM22552 and FAM22553, Isolated from Swiss Semihard Red-Smear Cheese

Michelangelo Storari,^a Daniel Wüthrich,^b Rémy Bruggmann,^b Hélène Berthoud,^a Emmanuelle Arias-Roth^a

Institute for Food Sciences, Agroscope, Bern, Switzerland^a; Interfaculty Bioinformatics Unit, University of Bern and Swiss Institute of Bioinformatics, Bern, Switzerland^b

***Clostridium tyrobutyricum* is the main microorganism responsible for late blowing defect in cheeses. Here, we present the draft genome sequences of two *C. tyrobutyricum* strains isolated from a Swiss semihard red-smear cheese. The two draft genomes comprise 3.05 and 3.08 Mbp and contain 3,030 and 3,089 putative coding sequences, respectively.**

Received 22 January 2015 Accepted 2 February 2015 Published 12 March 2015

Citation Storari M, Wüthrich D, Bruggmann R, Berthoud H, Arias-Roth E. 2015. Draft genome sequences of *Clostridium tyrobutyricum* strains FAM22552 and FAM22553, isolated from Swiss semihard red-smear cheese. *Genome Announc* 3(2):e00078-15. doi:10.1128/genomeA.00078-15.

Copyright © 2015 Storari et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Michelangelo Storari, michelangelo.storari@agroscope.admin.ch.

The spore-forming anaerobic bacterium *Clostridium tyrobutyricum* is the main cause of late blowing defect in hard and semihard cheeses (1). *Clostridium* spores contaminate milk during milking and survive the pasteurization process. The growth of *C. tyrobutyricum* during cheese maturation causes the formation of butyric acid, carbon dioxide, and molecular hydrogen, leading to the loss of the commercial value of the final product (2). The primary metabolism of *C. tyrobutyricum* as well as its nutritional requirements are still almost unknown, despite the huge negative economic impact that this microorganism has on the cheese manufacturing industry. To date, only the genomes of a strain used in industrial butyric acid production and a strain isolated from Grana Padano cheese have been sequenced (3, 4).

In the present study, we report the draft genome sequences of two *C. tyrobutyricum* strains (FAM22552 and FAM22553; deposited in the culture collection of Agroscope, Bern, Switzerland) isolated from a Swiss semihard red-smear cheese with typical symptoms of late blowing in 2012. Genomic DNA was extracted from liquid cultures of the two *C. tyrobutyricum* strains using the EZ1 DNA tissue kit (Qiagen) and sent for sequencing to Mi-crosynth AG (Switzerland). Libraries were prepared with the Nextera XT DNA sample preparation kit (Illumina) and sequenced using the Illumina MiSeq sequencing platform. Low-quality bases of raw reads (2 × 301 bp) were removed using Trimmomatic (version 0.30, SLIDINGWINDOW:4:15 MINLEN:127 [5]) before *de novo* assembly was performed. *De novo* assembly was carried out using SPAdes version 3.1.0 (*k*-mer sizes of 21, 33, 55, 77, 99, and 127 [6]). The obtained contigs were assembled to scaffolds using SSPACE version 3.0 (default parameters [7]). Trimmed reads (Trimmomatic, SLIDINGWINDOW:4:15 MINLEN:101 CROP:101) were mapped to the scaffolds with Bowtie2 version 2.1.0 (default parameters [8]) to determine the depth of coverage using SAMtools version 0.1.19 (idxstats [9]). Scaffolds with a mean depth of coverage lower than 20% of the mean depth of coverage of all scaffolds as well as contigs shorter than 200 bp were discarded. The *de novo* assembly consists of 58 (3.05 Mbp) and 62 (3.08 Mbp) scaffolds for FAM22552 and FAM22553, respectively.

Assembled genomes were annotated using the RAST server (10). In total, 3,030 putative coding sequences were annotated for FAM22552 and 3,089 for FAM22553.

C. tyrobutyricum grows in cheese using lactate and acetate as carbon and energy sources (11). A preliminary analysis of the two annotated genomes revealed the presence of putative genes coding for many enzymes needed to convert these two compounds into butyric acid, carbon dioxide, and molecular hydrogen. These include L- and D-lactate dehydrogenases, pyruvate-ferredoxin oxidoreductases, thiolases, hydroxybutyryl-CoA dehydrogenases, crotonases, hydroxybutyryl-CoA dehydrogenases, phosphotrans-acetylases, and acetate kinases (for a review, see reference 12). Studies are now under way to further clarify the metabolism of *C. tyrobutyricum* in cheese.

Nucleotide sequence accession numbers. The whole-genome shotgun projects have been deposited at DDBJ/EMBL/GenBank under the accession numbers JTER00000000 (for *C. tyrobutyricum* FAM22552) and JTES00000000 (for *C. tyrobutyricum* FAM22553). The versions described in this paper are the versions JTER01000000 and JTES01000000.

ACKNOWLEDGMENTS

This project was carried out with an industrial partner of the Swiss cheese industry and is supported by the Commission for Technology and Innovation (CTI), project number 13074-1.

REFERENCES

1. Klijn N, Nieuwenhof FF, Hoolwerf JD, van der Waals CB, Weerkamp AH. 1995. Identification of *Clostridium tyrobutyricum* as the causative agent of late blowing in cheese by species-specific PCR amplification. *Appl Environ Microbiol* 61:2919–2924.
2. Bachmann HP. 1999. Technologische Einflussfaktoren auf die Buttersäuregärung. *Agrarforschung* 6:137–140.
3. Jiang L, Zhu L, Xu X, Li Y, Li S, Huang H. 2013. Genome sequence of *Clostridium tyrobutyricum* ATCC 25755, a butyric acid-overproducing strain. *Genome Announc* 1(3):e00308-13. <http://dx.doi.org/10.1128/genomeA.00308-13>.
4. Bassi D, Fontana C, Gazzola S, Pietta E, Puglisi E, Cappa F, Coccinelli PS. 2013. Draft genome sequence of *Clostridium tyrobutyri-*

- cum* strain UC7086, isolated from Grana Padano cheese with late-blowing defect. *Genome Announc* 1(4):e00614-1. <http://dx.doi.org/10.1128/genomeA.00614-13>.
5. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <http://dx.doi.org/10.1093/bioinformatics/btu170>.
 6. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
 7. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27: 578–579. <http://dx.doi.org/10.1093/bioinformatics/btq683>.
 8. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <http://dx.doi.org/10.1038/nmeth.1923>.
 9. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* 25:2078–2079. <http://dx.doi.org/10.1093/bioinformatics/btp352>.
 10. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
 11. Ivy RA, Wiedman M. 2014. *Clostridium tyrobutyricum*, p 468–473. In Robinson R, Batt CA, Patel P (ed), *Encyclopedia of food microbiology*. Elsevier, Amsterdam, Netherlands.
 12. Liew FM, Köpke M, Simpson SD. 2013. Gas fermentation for commercial biofuels production, p 125–173. In Fang Z (ed), *Liquid, gaseous and solid biofuels—conversion techniques*. InTech, Rijeka, Croatia.