

REVIEW ARTICLE

Correspondence:

Jens Fedder, Centre of Andrology & Fertility Clinic, Odense University Hospital, Odense, Denmark.
E-mail: jens.fedder@rsyd.dk

Keywords:

artificial reproductive treatments, chromatin structure, DNA fragmentation, intra uterine insemination, male infertility, spermatozoa

Received: 15-Nov-2016

Revised: 13-Apr-2017

Accepted: 20-Apr-2017

doi: 10.1111/andr.12381

DNA fragmentation in spermatozoa: a historical review

^{1,2,3}A. S. Rex, ¹J. Aagaard and ^{2,3}J. Fedder 

¹Aagaard Gynaecological Clinic, Aarhus, Denmark, ²Centre of Andrology & Fertility Clinic, Odense University Hospital, Odense, Denmark, and ³Department of Clinical Research, University of Southern Denmark, Odense, Denmark

SUMMARY

Sperm DNA Fragmentation has been extensively studied for more than a decade. In the 1940s the uniqueness of the spermatozoa protein complex which stabilizes the DNA was discovered. In the fifties and sixties, the association between unstable chromatin structure and subfertility was investigated. In the seventies, the impact of induced DNA damage was investigated. In the 1980s the concept of sperm DNA fragmentation as related to infertility was introduced as well as the first DNA fragmentation test: the Sperm Chromatin Structure Assay (SCSA). The terminal deoxynucleotidyl transferase nick end labelling (TUNEL) test followed by others was introduced in the nineties. The association between DNA fragmentation in spermatozoa and pregnancy loss has been extensively investigated spurring the need for a therapeutic tool for these patients. This gave rise to an increased interest in the aetiology of DNA damage. The present decade continues within this research area. Some of the more novel methods recently submerging are sorting of cells with increased DNA fragmentation and hyaluronic acid (HA) binding techniques. The clinical value of these tests remains to be elucidated. In spite of half a century of research within the area, this analysis is not routinely implemented into the fertility clinics. The underlying causes are multiple. The abundance of methods has impeded the need for a clinical significant threshold. One of the most promising methods was commercialized in 2005 and has been reserved for larger licensed laboratories. Myriads of reviews and meta-analyses on studies using different assays for analysis of DNA fragmentation, different clinical Artificial Reproductive Treatments (ART), different definitions of successful ART outcome and small patient cohorts have been published. Although the area of DNA fragmentation in spermatozoa is highly relevant in the fertility clinics, the need for further studies focusing on standardization of the methods and clinical implementation persists.

INTRODUCTION

Paternal contribution to the fertilization and to the development of healthy offspring is of vital importance. There have been reports of an increased risk of schizophrenia or autism in offspring from fathers with increasing age (Sipos *et al.*, 2004; Reichenberg *et al.*, 2006) and an increased risk of cancer in offspring from fathers with increased level of sperm DNA fragmentation because of smoking (Ji *et al.*, 1997). Furthermore, some spontaneous dominant genetic diseases, epilepsy and some birth defects are linked to paternal contribution (reviewed in Aitken *et al.*, 2009). In a number of studies, an association between increased DNA fragmentation in the spermatozoa and subfertility has been reported. Comparing studies of fertile and infertile males have shown that the amount of DNA fragmentation is significantly higher in the infertile group (Evenson *et al.*, 1999;

Spano *et al.*, 2000; Saleh *et al.*, 2003; Alkhayal *et al.*, 2013; Oleszczuk *et al.*, 2013). An abnormal chromatin packing is more recurrent in men with normozoospermia undergoing ART treatment than in fertile men (Alkhayal *et al.*, 2013). If the man has increased DNA fragmentation in the spermatozoa, a prolonged Time To Pregnancy (TTP) (Evenson *et al.*, 1999), an increased risk of a missed abortion (Virro *et al.*, 2004; Lin *et al.*, 2008; Zini *et al.*, 2008; Kennedy *et al.*, 2011; Dar *et al.*, 2013) and a significantly reduced success rate in in vivo fertilization of the partner have been observed (Spano *et al.*, 2000; Bungum *et al.*, 2004, 2007; Giwercman *et al.*, 2010; Zini, 2011). When seeking fertility treatment, it seems that sperm DNA fragmentation is of vital importance when planning the course of treatment. A study included 131 couples seeking fertility treatment by intrauterine inseminations (IUI). Twenty-three of the male patients had an

increased amount of SCSA defined DNA fragmentation followed by a pregnancy rate of 4% in their partner (Bungum *et al.*, 2004). A later study including 387 cycles showed that the pregnancy rate dropped to 3% if the level of DNA fragmentation exceeded 30% (Bungum *et al.*, 2007). In a smaller Danish study including 48 couples, no pregnancies were observed in couples, where the male DNA fragmentation exceeded 27% (Boe-Hansen *et al.*, 2006). DNA fragmentation may affect the fertilization rate after in vitro fertilization (IVF). No clear association between increased amount of DNA fragmentation and intracytoplasmic sperm injection (ICSI) has been established. However, DNA fragmentation may affect the clinical pregnancy rate (Oleszczuk *et al.*, 2016) (Bungum *et al.*, 2007; Dar *et al.*, 2013).

It thus seems that an increase in DNA fragmentation primarily affects in vivo fertility, either by reducing natural conception or by a significant reduction in successful intrauterine inseminations. It is estimated that up to 20% of males with semen parameters otherwise suitable for IUI treatment present with a DFI >30%. On this basis, the authors behind this study recommend that IVF or ICSI being the first choice of treatment if the amount of DNA fragmentation exceeds 30% (Giwercman *et al.*, 2010; Bungum *et al.*, 2011). However, in the study from Oleszczuk *et al.* (2016) it was found that the fertilization rates might also be decreased after IVF when DFI by SCSA exceeds 30%. For high degree of DFI, it thus might be relevant to proceed the treatment using ICSI (Oleszczuk *et al.*, 2016).

Together, these studies provide important insight into the significance of sperm DNA fragmentation when treating couples for infertility.

With this in mind - why is sperm DNA fragmentation testing not a standard diagnostic tool in the treatment of the male fertility patient?

The journey regarding DNA fragmentation in spermatozoa has been long and began more than half a century ago.

HISTORICAL OVERVIEW

Forties, fifties and sixties

In 1946, Pollister and Mirsky discovered that a large part of the protein complexes surrounding the DNA in trout sperm was not composed of histones but of protamines (Pollister & Mirsky, 1946). Later Alfert found that the protamines replace the histones after meiosis in the maturation of the salmon spermatozoa (Alfert, 1956). Today it is estimated that only 5–15% of the chromatin in the human spermatozoa consist of histones and the major part consists of protamines (Castillo *et al.*, 2015). Alongside the discovery of the double helix in 1953, Leuchtenberger *et al.* (1953) discovered that the amount of DNA from infertile males had a significantly larger variation compared with fertile males. Already at this time, it was discovered that the quality of a sperm sample was more than a question of number and motility of the spermatozoa (Leuchtenberger *et al.*, 1953).

The seventies

During the seventies, an increasing interest in a possible association between exposure of DNA damaging agents and a possible reduction in fertility emerged. In 1970, Ringertz *et al.* used an assay where bull spermatozoa were heated and the subsequent denaturation of the DNA was detected with acridine orange followed by microfluorimetry. They realized that the

spermatozoa possessed an increased stability during the spermiogenesis (Ringertz *et al.*, 1970). A decrease in epididymal sperm count and weight of the testis was observed in mice after exposure to irradiation. Subsequently, an increased pre-implantation loss was observed in the female mice (Searle & Beechey, 1974).

The eighties

In the eighties, the technology for molecular biology advanced. Evenson *et al.* developed a flow cytometric assay for detection of DNA fragmentation in spermatozoa (Evenson *et al.*, 1980). They called the assay Sperm Chromatin Structure Assay (SCSA). The assay is based on the detection of DNA fragmentation by flow cytometry. The sperm DNA is denaturated by acid at sites of DNA strand breaks and subsequent stained with the fluorescent cationic dye Acridine Orange (AO). In this assay, AO attaches to the DNA in the ratio of approximately two AO molecules per phosphate group (Evenson & Jost, 1994). When the laser from the flow cytometer illuminates the cells, AO fluoresces with a green emission when bound to double stranded (db) DNA and a red emission when bound to single stranded (ss) DNA. Furthermore, the flow cytometer measures forward scatter and side scatter of the sample and this can help exclude debris from the sample. Usually a total of 5000–10,000 cells are analysed. DNA Fragmentation Index (DFI) is described as the percent wise ratio of red fluorescence to green + red fluorescence (Larson *et al.*, 2000; Evenson *et al.*, 2002; Larson-Cook *et al.*, 2003). SCSA also measures High DNA Stainability (HDS), which is believed to be an expression of immature spermatozoa containing excess histones or other abnormal proteins (Evenson *et al.*, 2002; Bungum *et al.*, 2004).

The nineties

The field of single cell electrophoresis was developed in the eighties and optimized during the nineties making it possible to detect DNA fragmentation by the comet assay. The comet assay was a novel diagnostic tool for DNA fragmentation and was used to emphasize that spermatozoa from infertile men were more susceptible to induced damage than spermatozoa from fertile men. In the comet assay, 200–300 cells are covered with agarose gel and subsequently lysed. If the DNA is embedded with breaks, the supercoiling of the DNA is released allowing the DNA to migrate towards the anode. This migration leaves a comet-like tail and the fluorescent intensity of the tail relates to the number of DNA breaks (Hughes *et al.*, 1996; Aravindan *et al.*, 1997).

In the nineties, TUNEL used to detect DNA fragmentation in human spermatozoa was developed. In this assay a terminal deoxynucleotidyl transferase labels the DNA strand breaks with fluorescent dUTP nucleotides. The assay can be performed using either flow cytometry or microscopy. Thus, both the neutral and alkaline comet assay and the TUNEL assay are considered 'direct' assays as they measure actual DNA strand breaks, whereas some of the other assays developed measure the DNA susceptibility to denature at sites of ss or ds DNA breaks or a differentiated binding of a dye to ds- or ssDNA (Gorczyca *et al.*, 1993; Zini & Sigman, 2009; Henkel *et al.*, 2010).

The zeroes

In the zeroes, other methods for determination of DNA fragmentation appeared. The DNA Breakage Detection-Fluorescence

in situ Hybridization (DBD-FISH) was developed for human spermatozoa. In this assay, the spermatozoa are fixed in an agarose matrix and the DNA is transformed into ssDNA by an alkaline unwinding solution. After the proteins are removed, the DNA is made accessible to hybridization with relevant probes that highlight the area for analysis. If the DNA strand contains increased amount of DNA breaks more probes will hybridize resulting in an increased fluorescence. This technique can be used for detecting DNA damage within specific sequence areas (Fernandez *et al.*, 2000). The Sperm Chromatin Dispersion (SCD) test was also developed in this decade and was used to detect spermatozoa with increased amount of DNA fragmentation. In this assay, the spermatozoa are embedded in an agarose matrix and exposed to a lysing solution. The relaxed DNA loops prevent dispersion into the surrounding area. With specific DNA fluorochromes and a fluorescence microscope, the dispersed DNA is seen as a halo surrounding the nuclei. If the DNA is fragmented, little dispersion is seen resulting in a small halo. This makes it possible to detect spermatozoa with increased amount of DNA fragmentation (Fernandez *et al.*, 2003). Two years later, an advanced SCD test was developed as a kit, Halosperm® (Fernandez *et al.*, 2005).

In 2006, Li *et al.* developed the γ H2AX assay to determine double strand breaks in human spermatozoa. The assay takes advantage of the fact that some protein-kinases induce phosphorylation of Ser139 on the histone H2AX. Phosphospecific antibodies are able to recognize the phosphorylated serine residue. These are subsequently quantified by a flow cytometer. Although most histones are replaced by protamines in human spermatozoa during the spermatogenesis, a small fraction remains in the nucleosome (around 15%). This fraction also contains the H2AX histone. In a recent study in 2015, Garolla *et al.* investigated the predictive value of the method. The pregnancy rate after ICSI was investigated and the method was compared with TUNEL. In this study, it was seen that the γ H2AX percentage was higher in the males from non-pregnant couples. This study also showed that the γ H2AX has a better predictive value than TUNEL. In the present form, the method seems to be rather time consuming, as samples need more than 3 h of preparation before the flow cytometric analysis can be performed. This compared to the strict protocol from SCSA where samples can be prepared within few minutes (Li *et al.*, 2006; Garolla *et al.*, 2015; Evenson, 2016).

The tenths

From the late zeroes and into the tenths the focus concerning DNA fragmentation shifted from development of methods to the aetiology of sperm DNA fragmentation. Furthermore, it became more evident that increased DNA fragmentation could be a valuable tool when deciding which type of fertility treatment the couple should be offered. Several studies added to the viewpoint that fertility treatment with IUI had very low chance of resulting in pregnancy if the SCSA-DFI in the spermatozoa was increased. However, the implantation rate after ICSI is not affected by increased amount of DNA fragmentation in the spermatozoa, it has been seen that the risk of early pregnancy loss is increased in these couples (Carrell *et al.*, 2003; Borini *et al.*, 2006; Gil-Villa *et al.*, 2009; Brahem *et al.*, 2011; Absalan *et al.*, 2012; Oleszczuk *et al.*, 2016).

In 2005, Greco *et al.* showed that ICSI with testicular sperm resulted in a significantly higher clinical pregnancy rate compared with ICSI where ejaculated sperm was used. This provided one of the first treatment options for male fertility patients with increased sperm DNA fragmentation. This study also gave insight to the aetiology of DNA damage as at least a part of the DNA damage seemed to appear after the spermatozoa have left the testis (Greco *et al.*, 2005). Recently, both Esteves *et al.* and Pabuccu *et al.* achieved similar results. Other methods such as intracytoplasmic morphologically selected sperm injection (IMSI) and motile sperm organelle morphology examination (MSOME) have been used in order to circumvent the negative effects of increased DNA fragmentation. These methods are based on a real time examination of the spermatozoa under an increased magnification (up to $\times 13,000$) which makes it possible to choose spermatozoa with better chromatin status and lower aneuploidy rate. If the spermatozoa showed a lack of vacuoles, the results improved further (Garolla *et al.*, 2008, 2014; Gosálvez *et al.*, 2013). In a recent study, Bradley *et al.* compared fertilization rates after physiological ICSI (PICS), IMSI and extraction of testicular sperm, respectively. They found that the use of testicular sperm significantly increased the fertilization rate, pregnancy rate and live birth rate after ICSI for patients with increased DNA fragmentation (Bradley *et al.*, 2016).

DNA fragmentation and infertility

A substantial amount of literature that strengthened the theory of a connection between increased DNA fragmentation and infertility had been published, and in 2015 Zini concluded that testing for DNA fragmentation should be a part of the routine male infertility diagnosing (Esteves *et al.*, 2015; Zini, 2015; Pabuccu *et al.*, 2016).

As the research in the area expanded, several studies show that the origin of DNA fragmentation can be very diverse. Link has been seen between increased DNA fragmentation and inadvertent effects during the spermiogenesis, increased amount of oxidative stress, sperm collection methods, storage temperature, varicocele, bacterial infections, age, temperature of the testes and reaction to medicine (reviewed in Gonzalez-Marin *et al.*, 2012). It is thus possible that the damage to the DNA happens in multiple steps. This has probably contributed to the blurred picture of DNA fragmentation. One theory is that the DNA is subjected to damaging events during the spermatogenesis. This could include nicks in the backbone of the DNA or poor packaging of the chromatin during the replacement of histones. Subsequently, the already weakened DNA is more susceptible to external stressors such as medication, temperature and Reactive Oxygen Species (ROS) (McPherson & Longo, 1993; Pradeepa & Rao, 2007).

Antioxidants

The role of antioxidants has been studied extensively in several areas in the last decade. Regarding spermatozoa, ROS are believed to play a part in the presence of DNA fragmentation. ROS play a positive role in several crucial functions such as proliferation and differentiation of cells. However, a pathogenic effect can occur when the balance between ROS and antioxidants are disturbed. This can result in an excess of ROS, for example in the reproductive tract or in the seminal plasma. It has been shown in several studies that antioxidants can have a

positive impact on some of the primary seminal parameters (Zini *et al.*, 2009; Abad *et al.*, 2013; Agarwal *et al.*, 2014). The distribution of dietary supplements to males with increased DFI have previously shown a significant reduction in DFI and an increase in the clinical pregnancy rate (Wright *et al.*, 2014). However, the overall effect of antioxidants remains controversial. This is mainly because of non-standardized assays for determination of ROS or antioxidant capacity, diversity in methods for determination of DNA fragmentation, lack of distinction between direct and indirect antioxidants and inadequate data on fertilization and pregnancy rates (Chen *et al.*, 2013).

Novel methods

In the present decade, the magnetic activated cell sorting (MACS) technique was enhanced. This method was used for the first time in connection to fertility treatment in the zeroes (Paasch *et al.*, 2007). In this assay, magnetic particles conjugated to proteins or antibodies target the cells of interest. This could be apoptotic surface markers like externalized phospholipid phosphatidylserine (PS). PS has a high affinity for annexin V, which cannot cross the membrane. Any conjugation between the two will therefore happen on spermatozoa with externalized PS, which is seen in apoptotic cells. The magnetic particles, and thereby the apoptotic cells, can subsequently be removed by a magnetic cell separation column (Said *et al.*, 2008).

A novel method where spermatozoa with increased amount of DNA fragmentation are separated by fluorescence-activated cell sorting has recently been presented. The spermatozoa are stained using a YO-PRO staining technique. The researchers behind the study showed that it is possible to separate the dead spermatozoa and the spermatozoa with increased amount of DNA fragmentation from the normal spermatozoa. This makes it possible to optimize the sperm sample before fertility treatment like ICSI is initiated (Ribeiro *et al.*, 2013). However, recent research has shown that methods where ICSI is optimized remains controversial (Tavalaee *et al.*, 2012; Troya & Zorrilla, 2015). Another novel method being investigated in the present decade is the possibility of detecting damage in spermatozoa by oligopeptides. A synthetic oligopeptide binds to the damaged DNA. The non-binding end of the oligopeptide consists of a rhodamine B dye that can be detected with fluorescence microscopy. There was seen a correlation of the amount of DNA damage detected with this method and the more classical methods such as SCD, comet and TUNEL (Enciso *et al.*, 2012). Another novel method that has been studied in the present decade is the HA binding technique. HA surrounds the oocyte only allowing spermatozoa with sufficient expression of specific receptors to fertilize it. It seems that there is an inverse association between the ability of spermatozoa to bind to HA and chromosomal abnormalities in the spermatozoa (Mokanszki *et al.*, 2012). In a study, it was found that HA binding test increased the chance of selecting a spermatozoon with a low amount of DNA fragmentation possibly optimizing the chance of pregnancy. A commercial kit has been developed thereby increasing the availability to the method (Parmegiani *et al.*, 2010) (Parmegiani *et al.*, 2012). However, in a recent meta-analysis it was not found that HA binding test increases fertilization rates after ICSI (Beck-Fruchter *et al.*, 2016) and further research in the area is thus needed for this test to have a relevance in the fertility clinics.

DNA fragmentation and pregnancy loss

Research continuously seems to focus on the possible association between sperm DNA fragmentation and recurrent pregnancy loss (Coughlan *et al.*, 2015; Leach *et al.*, 2015; Barih *et al.*, 2016; Zidi-Jrah *et al.*, 2016). Furthermore, an increasing interest has supervened regarding the types of fragmentations present in the DNA (Wei *et al.*, 2015) and how DNA fragmentation can be reduced in cryopreservation (Ghorbani *et al.*, 2016; Kably-Ambe *et al.*, 2016; Simonenko *et al.*, 2016). Recent papers have shown a possible correlation between an increased amount of DNA fragmentation and some of the natural antioxidants present in the seminal plasma like superoxide dismutase (Wdowiak *et al.*, 2015) and glutathione peroxidase (Dorostghoal *et al.*, 2017).

In spite of the effort in the last couples of decades, there is still a long way to go within the field of DNA fragmentation in spermatozoa. Substantial amounts of reviews and meta-analysis have been published, many of them imploring further studies with a controlled, randomized study population and more sensitive assays (Lewis *et al.*, 2013; Palermo *et al.*, 2014; Zhao *et al.*, 2014; Osman *et al.*, 2015).

In Fig. 1, an illustrative view of the historical development of DNA fragmentation in spermatozoa is presented.

DISCUSSION AND CONCLUSION

In spite of half a century's research and the widely accepted conviction that infertility and sperm DNA fragmentation are linked, this diagnostic tool is not yet a standard care in the fertility clinics.

Several issues contribute to this. The lack of uniformity in all assays, (except the SCSA) for analysis of DNA fragmentation and the absence of a clear clinical threshold, a myriad of studies using different assay, different clinical ART and diverse outcomes and small patient cohorts are all contributing factors. One of the obstacles is the difference in the methods used to assess the amount of DNA fragmentation present in the spermatozoa. Reviews and meta-analysis compare outcomes of fertility treatment across methods, which impede the progress of implementing the analysis in the fertility clinics. Another obstacle is the lack in knowledge concerning the aetiology of DNA fragmentation. There are several theories of the aetiology of DNA fragmentation. One is a two-step model for the development of DNA fragmentation in spermatozoa. In step one, an error in the spermatogenesis weakens the DNA and impairs the chromatin remodelling. This results in spermatozoa with low levels of nuclear protamine. In the second step, the vulnerable DNA is more susceptible to oxidative stress (Christensen & Birck, 2015). Another theory is that DNA fragmentation occurs after an interrupted apoptosis. An increased activity of apoptosis related proteins such as caspase 3 and 7, Fas and cPARP has been seen in samples with abnormal semen parameter or increased amount of DNA fragmentation. Apoptosis is believed to be activated by testicular conditions and oxidative stress. It is speculated that apoptosis could be the main pathway to DNA fragmentation in spermatozoa (Sakkas *et al.*, 2004; Manente *et al.*, 2015; Muratori *et al.*, 2015). An association between chromatin immaturity and DNA fragmentation has also been seen; which can be suggested as an accelerator for DNA fragmentation. Chromatin immaturity is believed to be caused by defects in the spermatogenesis (Sati *et al.*, 2008). The aetiology of DNA fragmentation is far from




Figure 1 Timeline. An illustrative view of the landmarks and development.

illuminated and the different theories are not mutually exclusive. It is possible that chromatin immaturity or oxidative stress triggers the activation of the apoptotic pathway. Furthermore, it is known that exogenous exposures, such as environment, lifestyle and health also contribute to increased amount of DNA fragmentation.

When relating to DNA fragmentation and infertility, the salient point is implementation in the fertility clinic. It is essential that this analysis is practicable in the daily work. Furthermore, uniformity and reproducibility across laboratories are of crucial importance. As the association between DNA fragmentation and pregnancy outcome is well established the analysis can be fully implemented as a diagnostic tool in the fertility clinics however, the aetiology of DNA fragmentation has not yet been fully elucidated.

When comparing outcomes of the different methods, there is a moderate correlation between SCSA, TUNEL and SCD with regard to levels of sperm DNA fragmentation. However, the Acridine Orange staining Technique (AOT) does not seem to have a clinical significance for fertility testing. In the AOT assay, the amount of DNA fragmentation is determined after a coloration with acridine orange and a microscopic evaluation. The method has recently been discredited by Evenson as a result of AO staining fluorescence fading and artefacts induced by glass/AO interactions (Evenson, 2016). Nonetheless SCSA also uses AO for the colouration of the DNA, the evaluation by microscopy vs. flow cytometer seems to be of crucial importance. Additionally, a study has shown that the neutral comet assay fails to distinguish between fertile donors and infertility patients. It does, however, relate to the risk of miscarriage. The alkaline comet assay seems to have a moderate correlation with SCSA, TUNEL and SCD. When the predictive values of the methods are assessed, there seems to be conflicting results regarding predictability for fertilization. It has been reported by Ribas-Maynou in 2013 that the alkaline comet assay has the highest sensitivity followed by the TUNEL, SCD and SCSA analysis and subsequently the neutral comet assay. Chohan *et al.* found a strong relationship between SCSA and TUNEL (Chohan *et al.*, 2006; Ribas-Maynou *et al.*,

2013). Furthermore, a systemic review and meta-analysis from 2016 claims that the comet assay and the TUNEL assay has the best predictability after IVF or ICSI of the methods assessed (Cissen *et al.*, 2016).

Other tests, like SCD are available as an easy to use kit but might be less accurate. However, the comet assay has shown the best predictability in some studies, it lacks a clear threshold and the methodology can change among laboratories. Furthermore, the comet assay as well as the SCD test, suffers by the fact that the evaluation of DNA fragmentation is estimated in only 200–300 cells. In addition, the comet assay is labour-intensive. The agreement between TUNEL and SCSA has been seen several times (Chohan *et al.*, 2006; Ribas-Maynou *et al.*, 2013; Evenson, 2016). When comparing the two flow cytometric assays, the TUNEL assay requires extensive preparation of the spermatozoa before analysis can be performed and there is currently a lack of a strict protocol. This inhibits the implementation of this method as a diagnostic tool in a clinical setting. SCSA has a strict protocol developed in 1980 and has been used for fertility assessment in both animal and human spermatozoa. The protocol is relatively easy and the analysis is not time consuming. It is possible to analyse up to 50 samples per day for an experienced lab technician. The analysis by flow cytometer allows evaluation of 10.000 spermatozoa within a minute or two resulting in a more robust analysis (Evenson *et al.*, 2002; Evenson & Wixon, 2006; Zini *et al.*, 2009; Evenson, 2013).

In 2005, the SCSA test was commercialized. Two European laboratories received license to perform the SCSA test. Fertility clinics were to ship sperm samples to these larger diagnostic centres in order to obtain a SCSA-DFI value (Evenson, 2011). As it is encouraged that DFI is determined using the commercial SCSAsoft[®] software the SCSA analysis have primarily been restricted to the larger licensed diagnostic laboratories (Evenson, 2011). In order for the analysis to be implemented into the individual fertility clinics, an investment in a flow cytometer is required. This has previously been an insurmountable cost. Recently, smaller bench-top flow cytometers having only a red and green fluorescent channel and a small air-cooled

blue laser has been developed. These are cheaper and require less space than the larger multichannel flow cytometers, making them more suitable for the smaller fertility clinics. More flexibility concerning software as well as a relocation from the larger commercial diagnostic labs to the individual fertility clinic will decrease the costs concerning the analysis thereby aiding the implementation of the analysis in the fertility clinics.

Bungum *et al.* (2011) estimates that 40% of all cases of unexplained infertility can be related to increased amount of DNA damage and suggests a treatment course where patients with DFI $\geq 30\%$ measured by SCSA should be referred directly to IVF/ICSI treatment. Furthermore, it is speculated that even a moderate increase in DFI (between 20–30% by SCSA) can give rise to a prolonged TTP – information that the treating physician can employ when counselling fertility patients and planning the course of treatment (Bungum *et al.*, 2011). As mentioned in the introduction, DNA fragmentation also seems to have implications for the offspring as it is linked to an increased risk of miscarriage or a number of pathogenic conditions in the offspring (reviewed in Aitken *et al.*, 2009; Sills & Christensen, 2015). In order for progress within this field a necessary next step is a combination of further clarification of the aetiology of DNA fragmentation and simplifying and standardizing the analysis. While it is very important to standardize all sperm fragmentation assays for utility in the ART clinic, a better understanding on the aetiology of sperm DNA fragmentation will be needed in order to develop effective therapeutic strategies for these patients. For couples where the male suffers from increased amount of DNA fragmentation, ICSI or TESA are currently the only methods that have shown any positive effect on the pregnancy rates. Both procedures are invasive and cannot cope with the increased amount of early pregnancy loss also seen in this group of patients. The goal must be to ease the diagnosis of these patients and to clarify the origin of the DNA fragmentation. This will make it possible to plan a treatment course with the aim of reducing the amount of DNA fragmentation and increase the rate of continuous pregnancy for these couples.

The predictive value of the analysis of DNA fragmentation in spermatozoa is often criticized. One point of criticism is that the method cannot predict all failure to conceive. As infertility is the couple's problem, one has to consider the fertility of the female as well. One single test of gamete dysfunction from just one partner of the couple cannot predict the outcome of the fertility treatment. Determination of DNA fragmentation is not a replacement of current diagnostic tools for infertility diagnosing. However, it is a valuable supplement adding independent information about the gamete status of the male partner and it is due time that this analysis becomes a standard tool in the fertility clinics.

ACKNOWLEDGEMENTS

This work is supported by grants from Innovation Fund Denmark and Odense University Hospital Ph.D. The work is also financially supported by Aagaard Gynaecological Clinic. Odense Patient data Explorative Network – OPEN has made valuable contribution to the project. Thanks to Jette Søndergaard from Aagaard Gynaecological Clinic for linguistic assistance of the manuscript.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHORS' CONTRIBUTIONS

JF and ASR were responsible for idea, design and outline of content of the paper. ASR wrote the manuscript draft. All authors revised the manuscript critically and approved the final version.

REFERENCES

- Abad C, Amengual MJ, Gosálvez J, Coward K, Hannaoui N, Benet J, García-Peiro A & Prats J. (2013) Effects of oral antioxidant treatment upon the dynamics of human sperm DNA fragmentation and subpopulations of sperm with highly degraded DNA. *Andrologia* 45, 211–216.
- Abalan F, Ghannadi A, Kazerooni M, Parifar R, Jamalzadeh F & Amiri S. (2012) Value of sperm chromatin dispersion test in couples with unexplained recurrent abortion. *J Assist Reprod Genet* 29, 11–14.
- Agarwal A, Durairajanayagam D & du Plessis SS. (2014) Utility of antioxidants during assisted reproductive techniques: an evidence based review. *Reprod Biol Endocrinol* 12, 112.
- Aitken RJ, De Iuliis GN & McLachlan RI. (2009) Biological and clinical significance of DNA damage in the male germ line. *Int J Androl* 32, 46–56.
- Alfert M. (1956) Chemical differentiation of nuclear proteins during spermatogenesis in the salmon. *J Biophys Biochem Cytol* 2, 109–114.
- Alkhayal A, San Gabriel M, Zeidan K, Alrabeeah K, Noel D, McGraw R, Bissonnette F, Kadoch IJ & Zini A. (2013) Sperm DNA and chromatin integrity in semen samples used for intrauterine insemination. *J Assist Reprod Genet* 30, 1519–1524.
- Aravindan GR, Bjordahl J, Jost LK & Evenson DP. (1997) Susceptibility of human sperm to in situ DNA denaturation is strongly correlated with DNA strand breaks identified by single-cell electrophoresis. *Exp Cell Res* 236, 231–237.
- Bareh GM, Jacoby E, Binkley P, Chang TC, Schenken RS & Robinson RD. (2016) Sperm deoxyribonucleic acid fragmentation assessment in normozoospermic male partners of couples with unexplained recurrent pregnancy loss: a prospective study. *Fertil Steril* 105, 329–36. e1.
- Beck-Fruchter R, Shalev E & Weiss A. (2016) Clinical benefit using sperm hyaluronic acid binding technique in ICSI cycles: a systematic review and meta-analysis. *Reprod Biomed Online* 32, 286–298.
- Boe-Hansen GB, Fedder J, Ersboll AK & Christensen P. (2006) The sperm chromatin structure assay as a diagnostic tool in the human fertility clinic. *Hum Reprod* 21, 1576–1582.
- Borini A, Tarozzi N, Bizzaro D, Bonu MA, Fava L, Flamigni C & Coticchio G. (2006) Sperm DNA fragmentation: paternal effect on early post-implantation embryo development in ART. *Hum Reprod* 21, 2876–2881.
- Bradley CK, McArthur SJ, Gee AJ, Weiss KA, Schmidt U & Toogood L. (2016) Intervention improves assisted conception intracytoplasmic sperm injection outcomes for patients with high levels of sperm DNA fragmentation: a retrospective analysis. *Andrology* 4, 903–910.
- Brahem S, Mehdi M, Landolsi H, Mougou S, Elghezal H & Saad A. (2011) Semen parameters and sperm DNA fragmentation as causes of recurrent pregnancy loss. *Urology* 78, 792–796.
- Bungum M, Humaidan P, Spano M, Jepson K, Bungum L & Giwercman A. (2004) The predictive value of sperm chromatin structure assay (SCSA) parameters for the outcome of intrauterine insemination, IVF and ICSI. *Hum Reprod* 19, 1401–1408.
- Bungum M, Humaidan P, Axmon A, Spano M, Bungum L, Erenpreiss J & Giwercman A. (2007) Sperm DNA integrity assessment in prediction of assisted reproduction technology outcome. *Hum Reprod* 22, 174–179.
- Bungum M, Bungum L & Giwercman A. (2011) Sperm chromatin structure assay (SCSA): a tool in diagnosis and treatment of infertility. *Asian J Androl* 13, 69–75.

- Carrell DT, Liu L, Peterson CM, Jones KP, Hatasaka HH, Erickson L & Campbell B. (2003) Sperm DNA fragmentation is increased in couples with unexplained recurrent pregnancy loss. *Arch Androl* 49, 49–55.
- Castillo J, Estanyol JM, Ballesca JL & Oliva R. (2015) Human sperm chromatin epigenetic potential: genomics, proteomics, and male infertility. *Asian J Androl* 17, 601–609.
- Chen SJ, Allam JP, Duan YG & Haidl G. (2013) Influence of reactive oxygen species on human sperm functions and fertilizing capacity including therapeutical approaches. *Arch Gynecol Obstet* 288, 191–199.
- Chohan KR, Griffin JT, Lafromboise M, De Jonge CJ & Carrell DT. (2006) Comparison of chromatin assays for DNA fragmentation evaluation in human sperm. *J Androl* 27, 53–59.
- Christensen P & Birck A. (2015) Comparison of methods for assesment of sperm DNA damage (fragmentation) and implications for the assisted reproductive technologies. In: *Screening the Single Euploid Embryo* (ed. E Scott Sills), pp. 53–71. Springer International Publishing, Switzerland.
- Cissen M, Wely MV, Scholten I, Mansell S, Bruin JP, Mol BW, Braat D, Repping S & Hamer G. (2016) Measuring sperm DNA fragmentation and clinical outcomes of medically assisted reproduction: a systematic review and meta-analysis. *PLoS ONE* 11, e0165125.
- Coughlan C, Clarke H, Cutting R, Saxton J, Waite S, Ledger W, Li T & Pacey AA. (2015) Sperm DNA fragmentation, recurrent implantation failure and recurrent miscarriage. *Asian J Androl* 17, 681–685.
- Dar S, Grover SA, Moskovtsev SI, Swanson S, Baratz A & Librach CL. (2013) In vitro fertilization-intracytoplasmic sperm injection outcome in patients with a markedly high DNA fragmentation index (>50%). *Fertil Steril* 100, 75–80.
- Dorostghoal M, Kazeminejad SR, Shahbazian N, Pourmehdi M & Jabbari A. (2017) Oxidative stress status and sperm DNA fragmentation in fertile and infertile men. *Andrologia* doi: 10.1111/and.12762.
- Enciso M, Pieczenik G, Cohen J & Wells D. (2012) Development of a novel synthetic oligopeptide for the detection of DNA damage in human spermatozoa. *Hum Reprod* 27, 2254–2266.
- Esteves SC, Sanchez-Martin F, Sanchez-Martin P, Schneider DT & Gosalvez J. (2015) Comparison of reproductive outcome in oligozoospermic men with high sperm DNA fragmentation undergoing intracytoplasmic sperm injection with ejaculated and testicular sperm. *Fertil Steril* 104, 1398–1405.
- Evenson D. (2011) Sperm Chromatin Structure Assay (SCSA[®]): 30 years of experience with the SCSA[®]. In: *Sperm Chromatin: Biological and Clinical Applications in Male Infertility and Assisted Reproduction* (eds A Agarwal & A Zini), pp. 125–149. New York: Springer Publishers.
- Evenson DP. (2013) Sperm chromatin structure assay (SCSA(R)). *Methods Mol Biol* 927, 147–164.
- Evenson DP. (2016) The Sperm Chromatin Structure Assay (SCSA(R)) and other sperm DNA fragmentation tests for evaluation of sperm nuclear DNA integrity as related to fertility. *Anim Reprod Sci* 169, 56–75.
- Evenson D & Jost L. (1994) Sperm chromatin structure assay: DNA denaturability. *Methods Cell Biol* 42, 159–176.
- Evenson DP & Wixon R. (2006) Clinical aspects of sperm DNA fragmentation detection and male infertility. *Theriogenology* 65, 979–991.
- Evenson DP, Darzynkiewicz Z & Melamed MR. (1980) Relation of mammalian sperm chromatin heterogeneity to fertility. *Science* 210, 1131–1133.
- Evenson DP, Jost LK, Marshall D, Zinaman MJ, Clegg E, Purvis K, de Angelis P & Claussen OP. (1999) Utility of the sperm chromatin structure assay as a diagnostic and prognostic tool in the human fertility clinic. *Hum Reprod* 14, 1039–1049.
- Evenson DP, Larson KL & Jost LK. (2002) Sperm chromatin structure assay: its clinical use for detecting sperm DNA fragmentation in male infertility and comparisons with other techniques. *J Androl* 23, 25–43.
- Fernandez JL, Vazquez-Gundin F, Delgado A, Goyanes VJ, Ramiro-Diaz J, de la Torre J & Gosalvez J. (2000) DNA breakage detection-FISH (DBD-FISH) in human spermatozoa: technical variants evidence different structural features. *Mutat Res* 453, 77–82.
- Fernandez JL, Muriel L, Rivero MT, Goyanes V, Vazquez R & Alvarez JG. (2003) The sperm chromatin dispersion test: a simple method for the determination of sperm DNA fragmentation. *J Androl* 24, 59–66.
- Fernandez JL, Muriel L, Goyanes V, Segrelles E, Gosalvez J, Enciso M, LaFromboise M & De Jonge C. (2005) Simple determination of human sperm DNA fragmentation with an improved sperm chromatin dispersion test. *Fertil Steril* 84, 833–842.
- Garolla A, Fortini D, Menegazzo M, De Toni L, Nicoletti V, Moretti A, Selice R, Engl B & Foresta C. (2008) High-power microscopy for selecting spermatozoa for ICSI by physiological status. *Reprod Biomed Online* 17, 610–616.
- Garolla A, Cosci I, Menegazzo M, De Palo R, Ambrosini G, Sartini B, Pizzol D & Foresta C. (2014) Sperm selected by both birefringence and motile sperm organelle morphology examination have reduced deoxyribonucleic acid fragmentation. *Fertil Steril* 101, 647–652.
- Garolla A, Cosci I, Bertoldo A, Sartini B, Boudjema E & Foresta C. (2015) DNA double strand breaks in human spermatozoa can be predictive for assisted reproductive outcome. *Reprod Biomed Online* 31, 100–107.
- Ghorbani M, Vatannejad A, Khodadadi I, Amiri I & Tavilani H. (2016) Protective effects of glutathione supplementation against oxidative stress during cryopreservation of human spermatozoa. *Cryo Lett* 37, 34–40.
- Gil-Villa AM, Cardona-Maya W, Agarwal A, Sharma R & Cadavid A. (2009) Role of male factor in early recurrent embryo loss: do antioxidants have any effect? *Fertil Steril* 92, 565–571.
- Giwercman A, Lindstedt L, Larsson M, Bungum M, Spano M, Levine RJ & Rylander L. (2010) Sperm chromatin structure assay as an independent predictor of fertility in vivo: a case-control study. *Int J Androl* 33, e221–e227.
- Gonzalez-Marin C, Gosalvez J & Roy R. (2012) Types, causes, detection and repair of DNA fragmentation in animal and human sperm cells. *Int J Mol Sci* 13, 14026–14052.
- Gorczyca W, Traganos F, Jesionowska H & Darzynkiewicz Z. (1993) Presence of DNA strand breaks and increased sensitivity of DNA in situ to denaturation in abnormal human sperm cells: analogy to apoptosis of somatic cells. *Exp Cell Res* 207, 202–205.
- Gosalvez A, Migueles B, López-Fernández C, Sánchez-Martín F & Sánchez-Martín P. (2013) Single sperm selection and DNA fragmentation analysis: the case of MSOME/IMSI. *Nat Sci* 5, 7–14.
- Greco E, Scarselli F, Iacobelli M, Rienzi L, Ubaldi F, Ferrero S, Franco G, Anniballo N, Mendoza C & Tesarik J. (2005) Efficient treatment of infertility due to sperm DNA damage by ICSI with testicular spermatozoa. *Hum Reprod* 20, 226–230.
- Henkel R, Hoogendijk CF, Bouic PJ & Kruger TF. (2010) TUNEL assay and SCSA determine different aspects of sperm DNA damage. *Andrologia* 42, 305–313.
- Hughes CM, Lewis SE, McKelvey-Martin VJ & Thompson W. (1996) A comparison of baseline and induced DNA damage in human spermatozoa from fertile and infertile men, using a modified comet assay. *Mol Hum Reprod* 2, 613–619.
- Ji BT, Shu XO, Linet MS, Zheng W, Wacholder S, Gao YT, Ying DM & Jin F. (1997) Paternal cigarette smoking and the risk of childhood cancer among offspring of nonsmoking mothers. *J Natl Cancer Inst* 89, 238–244.
- Kably-Ambe A, Carballo-Mondragon E, Roque-Sanchez AM, Duran-Monteros L & Amaro-Hernandez EA. (2016) Evaluation of semen parameters in long term cryopreserved samples for over 10 years. *Ginecol Obstet Mex* 84, 1–6.
- Kennedy C, Ahlering P, Rodriguez H, Levy S & Sutovsky P. (2011) Sperm chromatin structure correlates with spontaneous abortion and

- multiple pregnancy rates in assisted reproduction. *Reprod Biomed Online* 22, 272–276.
- Larson KL, DeJonge CJ, Barnes AM, Jost LK & Evenson DP. (2000) Sperm chromatin structure assay parameters as predictors of failed pregnancy following assisted reproductive techniques. *Hum Reprod* 15, 1717–1722.
- Larson-Cook KL, Brannian JD, Hansen KA, Kasperon KM, Aamold ET & Evenson DP. (2003) Relationship between the outcomes of assisted reproductive techniques and sperm DNA fragmentation as measured by the sperm chromatin structure assay. *Fertil Steril* 80, 895–902.
- Leach M, Aitken RJ & Sacks G. (2015) Sperm DNA fragmentation abnormalities in men from couples with a history of recurrent miscarriage. *Aust N Z J Obstet Gynaecol* 55, 379–383.
- Leuchtenberger C, Schrader F, Weir DR & Gentile DP. (1953) The desoxyribosenucleic acid (DNA) content in spermatozoa of fertile and infertile human males. *Chromosoma* 6, 61–78.
- Lewis SE, John Aitken R, Conner SJ, Iulius GD, Evenson DP, Henkel R, Giwercman A & Gharagozloo P. (2013) The impact of sperm DNA damage in assisted conception and beyond: recent advances in diagnosis and treatment. *Reprod Biomed Online* 27, 325–337.
- Li Z, Yang J & Huang H. (2006) Oxidative stress induces H2AX phosphorylation in human spermatozoa. *FEBS Lett* 580, 6161–6168.
- Lin MH, Kuo-Kuang Lee R, Li SH, Lu CH, Sun FJ & Hwu YM. (2008) Sperm chromatin structure assay parameters are not related to fertilization rates, embryo quality, and pregnancy rates in vitro fertilization and intracytoplasmic sperm injection, but might be related to spontaneous abortion rates. *Fertil Steril* 90, 352–359.
- Manente L, Pecoraro S, Picillo E, Gargiulo U, Gargiulo P, De Luca A & Politano L. (2015) Molecular evidence of apoptotic pathway activation in semen samples with high DNA fragmentation. *In Vivo* 29, 289–294.
- McPherson S & Longo FJ. (1993) Chromatin structure-function alterations during mammalian spermatogenesis: DNA nicking and repair in elongating spermatids. *Eur J Histochem* 37, 109–128.
- Mokanszki A, Molnar Z, Ujfalusi A, Balogh E, Bazsane ZK, Varga A, Jakab A & Olah E. (2012) Correlation study between sperm concentration, hyaluronic acid-binding capacity and sperm aneuploidy in Hungarian patients. *Reprod Biomed Online* 25, 620–626.
- Muratori M, Tamburrino L, Marchiani S, Cambi M, Olivito B, Azzari C, Forti G & Baldi E. (2015) Investigation on the origin of sperm DNA fragmentation: role of apoptosis, immaturity and oxidative stress. *Mol Med* 21, 109–122.
- Oleszczuk K, Augustinsson L, Bayat N, Giwercman A & Bungum M. (2013) Prevalence of high DNA fragmentation index in male partners of unexplained infertile couples. *Andrology* 1, 357–360.
- Oleszczuk K, Giwercman A & Bungum M. (2016) Sperm chromatin structure assay in prediction of in vitro fertilization outcome. *Andrology* 4, 290–296.
- Osman A, Alsomait H, Seshadri S, El-Toukhy T & Khalaf Y. (2015) The effect of sperm DNA fragmentation on live birth rate after IVF or ICSI: a systematic review and meta-analysis. *Reprod Biomed Online* 30, 120–127.
- Paasch U, Grunewald S & Glander HJ. (2007) Sperm selection in assisted reproductive techniques. *Soc Reprod Fertil Suppl* 65, 515–525.
- Pabuccu EG, Caglar GS, Tangal S, Haliloglu AH & Pabuccu R. (2016) Testicular versus ejaculated spermatozoa in ICSI cycles of normozoospermic men with high sperm DNA fragmentation and previous ART failures. *Andrologia* 49, e12609.
- Palermo GD, Neri QV, Cozzubbo T & Rosenwaks Z. (2014) Perspectives on the assessment of human sperm chromatin integrity. *Fertil Steril* 102, 1508–1517.
- Parmegiani L, Cognigni GE, Bernardi S, Troilo E, Ciampaglia W & Filicori M. (2010) “Physiologic ICSI”: hyaluronic acid (HA) favors selection of spermatozoa without DNA fragmentation and with normal nucleus, resulting in improvement of embryo quality. *Fertil Steril* 93, 598–604.
- Parmegiani L, Cognigni GE, Bernardi S, Troilo E, Taraborrelli S, Arnone A, Maccarini AM & Filicori M. (2012) Comparison of two ready-to-use systems designed for sperm-hyaluronic acid binding selection before intracytoplasmic sperm injection: PICSI vs. Sperm Slow: a prospective, randomized trial. *Fertil Steril* 98, 632–637.
- Pollister AW & Mirsky AE. (1946) The nucleoprotamine of trout sperm. *J Gen Physiol* 30, 101–116.
- Pradeepa MM & Rao MR. (2007) Chromatin remodeling during mammalian spermatogenesis: role of testis specific histone variants and transition proteins. *Soc Reprod Fertil Suppl* 63, 1–10.
- Reichenberg A, Gross R, Weiser M, Bresnahan M, Silverman J, Harlap S, Rabinowitz J, Shulman C, Malaspina D, Lubin G, Knobler HY, Davidson M & Susser E. (2006) Advancing paternal age and autism. *Arch Gen Psychiatry* 63, 1026–1032.
- Ribas-Maynou J, Garcia-Peiro A, Fernandez-Encinas A, Abad C, Amengual MJ, Prada E, Navarro J & Benet J. (2013) Comprehensive analysis of sperm DNA fragmentation by five different assays: TUNEL assay, SCSA, SCD test and alkaline and neutral Comet assay. *Andrology* 1, 715–722.
- Ribeiro SC, Sartorius G, Pletscher F, de Geyter M, Zhang H & de Geyter C. (2013) Isolation of spermatozoa with low levels of fragmented DNA with the use of flow cytometry and sorting. *Fertil Steril* 100, 686–694.
- Ringertz NR, Gledhill BL & Darzynkiewicz Z. (1970) Changes in deoxyribonucleoprotein during spermiogenesis in the bull. Sensitivity of DNA to heat denaturation. *Exp Cell Res* 62, 204–218.
- Said TM, Agarwal A, Zborowski M, Grunewald S, Glander HJ & Paasch U. (2008) Utility of magnetic cell separation as a molecular sperm preparation technique. *J Androl* 29, 134–142.
- Sakkas D, Seli E, Manicardi GC, Nijs M, Ombelet W & Bizzaro D. (2004) The presence of abnormal spermatozoa in the ejaculate: did apoptosis fail? *Hum Fertil (Camb)* 7, 99–103.
- Saleh RA, Agarwal A, Nada EA, El-Tonsy MH, Sharma RK, Meyer A, Nelson DR & Thomas AJ. (2003) Negative effects of increased sperm DNA damage in relation to seminal oxidative stress in men with idiopathic and male factor infertility. *Fertil Steril* 79(Suppl 3), 1597–1605.
- Sati L, Ovari L, Bennett D, Simon SD, Demir R & Huszar G. (2008) Double probing of human spermatozoa for persistent histones, surplus cytoplasm, apoptosis and DNA fragmentation. *Reprod Biomed Online* 16, 570–579.
- Searle AG & Beechey CV. (1974) Sperm-count, egg-fertilization and dominant lethality after X-irradiation of mice. *Mutat Res* 22, 63–72.
- Simonenko EY, Garmaeva SB, Yakovenko SA, Grigorieva AA, Tverdislov VA, Mironova AG & Aprishko VP. (2016) Influence of storage temperature and cryopreservation conditions on the extent of human sperm DNA fragmentation. *Biofizika* 61, 316–320.
- Sipos A, Rasmussen F, Harrison G, Tynelius P, Lewis G, Leon DA & Gunnell D. (2004) Paternal age and schizophrenia: a population based cohort study. *BMJ* 329, 1070.
- Spano M, Bonde JP, Hjollund HI, Kolstad HA, Cordelli E & Leter G. (2000) Sperm chromatin damage impairs human fertility. The Danish First Pregnancy Planner Study Team. *Fertil Steril* 73, 43–50.
- Tavalaee M, Deemeh MR, Arbabian M & Nasr-Esfahani MH. (2012) Density gradient centrifugation before or after magnetic-activated cell sorting: which technique is more useful for clinical sperm selection? *J Assist Reprod Genet* 29, 31–38.
- Troya J & Zorrilla I. (2015) Annexin V-MACS in infertile couples as method for separation of sperm without DNA fragmentation. *JBRA Assist Reprod* 19, 66–69.
- Virro MR, Larson-Cook KL & Evenson DP. (2004) Sperm chromatin structure assay (SCSA) parameters are related to fertilization, blastocyst development, and ongoing pregnancy in vitro fertilization and intracytoplasmic sperm injection cycles. *Fertil Steril* 81, 1289–1295.

- Wdowiak A, Bakalczuk S & Bakalczuk G. (2015) Decreased activity of superoxide dismutase in the seminal plasma of infertile men correlates with increased sperm deoxyribonucleic acid fragmentation during the first hours after sperm donation. *Andrology* 3, 748–755.
- Wei RX, Chen JW, Huang JH, Zhang XX & Cui Y. (2015) Type of sperm DNA strand breaks in infertile men and its clinical implication. *Zhonghua Nan Ke Xue* 21, 604–609.
- Wright C, Milne S & Leeson H. (2014) Sperm DNA damage caused by oxidative stress: modifiable clinical, lifestyle and nutritional factors in male infertility. *Reprod Biomed Online* 28, 684–703.
- Zhao J, Zhang Q, Wang Y & Li Y. (2014) Whether sperm deoxyribonucleic acid fragmentation has an effect on pregnancy and miscarriage after in vitro fertilization/intracytoplasmic sperm injection: a systematic review and meta-analysis. *Fertil Steril* 102, 998–1005. e8
- Zidi-Jrah I, Hajlaoui A, Mougou-Zerelli S, Kammoun M, Meniaoui I, Sallem A, Brahem S, Fekih M, Bibi M, Saad A & Ibala-Romdhane S. (2016) Relationship between sperm aneuploidy, sperm DNA integrity, chromatin packaging, traditional semen parameters, and recurrent pregnancy loss. *Fertil Steril* 105, 58–64.
- Zini A. (2011) Are sperm chromatin and DNA defects relevant in the clinic? *Syst Biol Reprod Med* 57, 78–85.
- Zini A. (2015) Use of testicular sperm for intracytoplasmic sperm injection in oligozoospermic couples with high levels of sperm deoxyribonucleic acid fragmentation. *Fertil Steril* 104, 1376–1377.
- Zini A & Sigman M. (2009) Are tests of sperm DNA damage clinically useful? Pros and cons *J Androl* 30, 219–229.
- Zini A, Boman JM, Belzile E & Ciampi A. (2008) Sperm DNA damage is associated with an increased risk of pregnancy loss after IVF and ICSI: systematic review and meta-analysis. *Hum Reprod* 23, 2663–2668.
- Zini A, San GM & Baazeem A. (2009) Antioxidants and sperm DNA damage: a clinical perspective. *J Assist Reprod Genet* 26, 427–432.