



# Redox-Sensitive Induction of Src/PI3-kinase/Akt and MAPKs Pathways Activate eNOS in Response to EPA:DHA 6:1

Faraj Zgheel, Mahmoud Alhosin, Sherzad Rashid, Mélanie Burban, Cyril Auger, Valérie B. Schini-Kerth\*

CNRS UMR 7213 Laboratoire de Biophotonique et Pharmacologie, Université de Strasbourg, Faculté de Pharmacie, Illkirch, France

## Abstract

**Aims:** Omega-3 fatty acid products containing eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) have vasoprotective effects, in part, by stimulating the endothelial formation of nitric oxide (NO). This study determined the role of the EPA:DHA ratio and amount, and characterized the mechanism leading to endothelial NO synthase (eNOS) activation.

**Methods and Results:** EPA:DHA 6:1 and 9:1 caused significantly greater endothelium-dependent relaxations in porcine coronary artery rings than EPA:DHA 3:1, 1:1, 1:3, 1:6, 1:9, EPA and DHA alone, and EPA:DHA 6:1 with a reduced EPA + DHA amount, which were inhibited by an eNOS inhibitor. Relaxations to EPA:DHA 6:1 were insensitive to cyclooxygenase inhibition, and reduced by inhibitors of either oxidative stress, Src kinase, PI3-kinase, p38 MAPK, MEK, or JNK. EPA:DHA 6:1 induced phosphorylation of Src, Akt, p38 MAPK, ERK, JNK and eNOS; these effects were inhibited by MnTMPyP. EPA:DHA 6:1 induced the endothelial formation of ROS in coronary artery sections as assessed by dihydroethidium, and of superoxide anions and hydrogen peroxide in cultured endothelial cells as assessed by electron spin resonance with the spin probe CMH, and the Amplex Red based assay, respectively.

**Conclusion:** Omega-3 fatty acids cause endothelium-dependent NO-mediated relaxations in coronary artery rings, which are dependent on the EPA:DHA ratio and amount, and involve an intracellular activation of the redox-sensitive PI3-kinase/Akt and MAPKs pathways to activate eNOS.

**Citation:** Zgheel F, Alhosin M, Rashid S, Burban M, Auger C, et al. (2014) Redox-Sensitive Induction of Src/PI3-kinase/Akt and MAPKs Pathways Activate eNOS in Response to EPA:DHA 6:1. PLoS ONE 9(8): e105102. doi:10.1371/journal.pone.0105102

**Editor:** Yulia Komarova, University of Illinois at Chicago, United States of America

**Received:** October 21, 2013; **Accepted:** July 21, 2014; **Published:** August 18, 2014

**Copyright:** © 2014 Zgheel et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported, in part, by an unrestricted research grant from Pivotal Therapeutics Inc., Toronto, Canada. Faraj Zgheel is supported by a fellowship from the Higher Education Ministry, Libya, and Sherzad Rashid by a fellowship from the Higher Education Ministry, KRG, Iraq. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** Regarding the Pivotal Therapeutics Inc., the authors confirm that they received an unrestricted research grant from this company. However, this does not alter their adherence to all the PLOS ONE policies on sharing data and materials.

\* Email: valerie.schini-kerth@unistra.fr

## Introduction

Many prospective cohort studies and meta-analyses have provided evidence that fish consumption, especially oily fish (e.g., salmon, trout, herring, sardines, and mackerel), with high amounts of omega-3 fatty acids, including eicosapentaenoic acid (EPA, C20:5  $\omega$ -3) and docosahexaenoic acid (DHA, C22:6  $\omega$ -3), or supplementation with omega-3 fatty acids reduce cardiovascular mortality in patients with heart disease [1,2]. In the GISSI trial published in 1999 for patients surviving a recent myocardial infarction, intake of 289 mg EPA plus 577 mg DHA per day reduced significantly fatal cardiovascular diseases by 30%, fatal coronary artery diseases by 35%, and sudden death by 45% [3]. However, more recent double-blind trials did not show an effect of an additional amount of EPA plus DHA on major cardiovascular endpoints in patients with coronary artery disease or after myocardial infarction [4,5]. Such differences have been explained by differences in study design, the fact that patients in the more recent trials were optimally treated not only by antithrombotics but also by antihypertensives and statins, and possibly also due to the use of different doses, sources (oily fish or fish-oil supplements),

and formulation of EPA and/or DHA [6]. The beneficial effect of omega-3 fatty acids on the cardiovascular system has been suggested to involve several mechanisms and in particular their ability to induce anti-inflammatory and anti-arrhythmic effects, to improve the lipid profile by lowering triglyceride levels, to prevent plaque development and possibly also to promote plaque stabilization, and to inhibit platelet aggregation [6]. In addition, the beneficial effect might also be due to their ability to improve endothelial dysfunction, which may represent a very early step in atherogenesis and is characterized by blunted endothelium-dependent vasodilatation mostly due to a reduced bioavailability of endothelium-derived nitric oxide (NO), and, often, endothelium-derived hyperpolarization (EDH). Indeed, chronic intake of fish oils increased endothelium-dependent relaxations in normal and in hypercholesterolemic and atherosclerotic porcine coronary arteries [7,8]. An improved endothelial function as assessed by flow-mediated vasodilatation was observed in humans after intake of fish-derived products or omega-3 fatty acids [9,10]. Moreover, intake of EPA and DHA improved the postprandial macro- and microvascular function in patients with type 2 diabetes mellitus [11]. EPA has been shown to cause endothelium-dependent

relaxations of isolated sheep pulmonary arteries, and to increase endothelial NO synthase (eNOS) activity and NO formation in cultured endothelial cells [12–14]. In addition, EPA but not DHA has also been shown to cause cerebral microvascular vasodilatation involving arachidonic acid metabolites [15]. On the other hand, DHA but not EPA supplementation improved forearm microvascular reactivity in overweight hyperlipidemic men [16]. In order to better understand the omega-3 fatty acid-induced endothelial protection, the ability of different omega-3 fatty acid products (EPA, DHA, different ratios of EPA:DHA, different content of EPA and DHA) to cause endothelium-dependent relaxations in porcine coronary arteries was evaluated. Thereafter, the endothelium-dependent relaxation to one of the most active omega-3 products was further characterized and in particular the role of endothelium-derived NO, EDH and vasoactive prostanoids, and the signal transduction pathway leading to eNOS activation were determined.

## Materials and Methods

### Vascular reactivity studies

Pig hearts were collected from the local slaughterhouse (Covpial, Holtzheim) and vascular reactivity was assessed as indicated previously [17,18]. Briefly, left circumflex coronary arteries were excised, cleaned of loose connective tissue and flushed with PBS without calcium to remove remaining blood. Rings of porcine coronary arteries (4–5 mm in length) were then suspended in organ baths containing oxygenated (95% O<sub>2</sub>; 5% CO<sub>2</sub>) Krebs bicarbonate solution (composition in mM: NaCl 119, KCl 4.7, KH<sub>2</sub>PO<sub>4</sub> 1.18, MgSO<sub>4</sub> 1.18, CaCl<sub>2</sub> 1.25, NaHCO<sub>3</sub> 25 and D-glucose 11, pH 7.4, 37°C) for the determination of changes in isometric tension (basal tension 5 g). The integrity of the endothelium was checked with bradykinin (0.3 μM). Rings were contracted (about 80% maximal contraction) with U46619, a thromboxane A<sub>2</sub> receptor agonist, before construction of a concentration-relaxation curve to an omega-3 fatty acid product. In some experiments, rings were exposed to an inhibitor for 30 min before the addition of U46619.

### Culture of coronary artery endothelial cells

Endothelial cells from coronary arteries were isolated from porcine left circumflex coronary arteries by collagenase treatment (type I, Worthington, 1 mg/ml for 12 min at 37°C), and cultured in culture dishes containing medium MCDB 131 (Invitrogen, Saint Aubin, France) supplemented with 15% fetal calf serum, penicillin (100 U/ml), streptomycin (100 U/ml), fungizone (250 μg/ml), and L-glutamine (2 mM) (all from Cambrex, Wiesbaden, Germany), and grown for 48–72 h. Cells (first passage) were exposed to serum-free culture medium in the presence of 0.1% bovine serum albumin (Euromedex, Souffelweyersheim, France) for 6 h prior to treatment.

### Western blot analysis

Endothelial cells were lysed in extraction buffer (composition in mM: Tris/HCl 20 (Euromedex), NaCl 150, Na<sub>3</sub>VO<sub>4</sub> 1, sodium pyrophosphate 10, NaF 20, okadaic acid 0.01, a tablet of protease inhibitor (Complete, Roche, Meylan, France) and 1% Triton X-100 (Euromedex); pH 7.5). Total proteins (10 μg) were separated on 10–12% SDS-polyacrylamide gels at 100 V for 2 h. Separated proteins were transferred electrophoretically onto polyvinylidene difluoride membranes (GE Healthcare, Velizy-Villacoublay, France) at 100 V for 120 min. Membranes were blocked with buffer containing 5% bovine serum albumin, Tris-buffered saline solution (Euromedex) and 0.1% Tween 20 (Sigma-Aldrich, Saint-

Quentin Fallavier, France) for 1 h. Membranes were incubated with a primary antibody (p-Src Ser17, p-Akt Ser473, p-p38 MAPK Thr180/Tyr182, p-JNK Thr183/Tyr185, p-ERK Thr202/Tyr204, p-eNOS Ser1177 from rabbit; eNOS and β-tubulin from mouse, Cell Signaling Technology; dilution of 1:1,000) overnight at 4°C. After washing, membranes were incubated with the appropriate horseradish peroxidase-conjugated secondary antibody (diluted to 1:10,000 for anti-mouse antibody and 1:5,000 for anti-rabbit antibody, Cell Signaling Technology, Danvers, MA, USA) at room temperature for 60 min. Immunoreactive bands were detected by enhanced chemiluminescence (GE Healthcare).

### Determination of the vascular formation of ROS

**Dihydroethidium staining.** Porcine coronary arteries (5–7 mm length) were embedded in OCT compound (Sakura Finetek, Villeneuve d'Ascq, France) and frozen in a nitrogen bath for cryostat sections. The redox-sensitive fluorescent dye dihydroethidium (DHE, 2.5 μM) was applied onto 25 μm unfixed cryosections of coronary arteries for 30 min at 37°C in a light-protected humidified chamber before the addition of EPA:DHA 6:1 (0.4% v/v) for 30 min. In some experiments, sections were incubated with a pharmacological tool for 30 min at 37°C before the addition of DHE. Sections were then washed, mounted in fluorescence mounting medium (Dako, Les Ulis, France) and cover-slipped. Images were obtained using a Leica SP2 UV DM IRBE laser scanning confocal microscope (Leica Microsystem, Nanterre, France). Quantification of staining levels was performed using FIJI GPL v2 software.

**Electron spin resonance (ESR).** ESR was used to determine whether EPA:DHA 6:1 stimulates the endothelial formation of superoxide anions in cultured endothelial cells using the superoxide anion cell-permeable spin probe CMH following the method described [19].

Endothelial cells were incubated with either solvent, SOD or PEG-SOD followed by treatment with EPA:DHA 6:1 (0.4% v/v) in Krebs-HEPES solution containing 500 μM of 1-hydroxy-3-methoxycarbonyl-2,2,5,5-tetramethylpyrrolidin (CMH, Noxygen, Germany), deferoxamine (25 μM) and DETC (5 μM) under constant temperature (37°C) for 30 min. The reaction was stopped by putting the samples on ice, subsequently they were frozen in liquid N<sub>2</sub> and analyzed by ESR at 77 K in a Dewar flask using a MS100 spectrometer (Magnetech Ltd., Berlin, Germany). Instrument settings were: temperature 77 K, microwave frequency 9.34 GHz, microwave power 10 mW, amplitude of modulation 100 mT, modulation of frequency 100 kHz, and 60 s of sweep time. Superoxide anions cannot be directly measured by ESR due to their short half-life. Therefore, the CMH spin-trapping reagent has been used. It reacts with superoxide anions to generate the stable CM<sup>•</sup> radical, which gives a characteristic triplet signal ESR spectra, with an amplitude directly proportional to the quantity of CM<sup>•</sup> present in the sample. The quantification of the signal is based on the mean of the height (amplitude) of the three signals. Values are expressed in signal amplitude (Amplitude, arbitrary units).

**Amplex Red assay.** Hydrogen peroxide measurements in cultured endothelial cells were made using the horseradish peroxidase-linked Amplex Red fluorescence assay kit as recommended (Invitrogen, Paisley, UK). Endothelial cells were incubated with either solvent, catalase or PEG-catalase for 30 min and then they were treated with EPA:DHA 6:1 (0.4% v/v) for 45 min. Amplex Red (50 μM) and horseradish peroxidase type II (0.1 U/ml) were added to the supernatant samples (100 μl). Fluorescence

readings were made in triplicate in a 96-well plate, and Ex/Em = 530/580 nm.

## Chemicals

All products were from Sigma-Aldrich (Saint-Quentin Fallavier, France), except wortmannin, PP2 and MnTMPyP from Enzo Life Sciences (Lausen, Switzerland). Omega-3 fatty acid products (Table 1) were provided by Pivotal Therapeutics, Inc (Woodbridge, ON, Canada).

## Statistical analysis

All values are expressed as means  $\pm$  SEM of *n* different experiments. Statistical analysis was performed using Student's paired *t* test or a two-way analysis of variance test followed by Bonferroni's post-hoc test as appropriate using Graphpad Prism software (version 5.04 for Windows, GraphPad Software, Inc., CA, USA). *P* < 0.05 was considered to be statistically significant.

## Results

### The omega-3 fatty acid-induced endothelium-dependent relaxations are dependent on the EPA:DHA ratio and content

To study the vasoactive effect of omega-3 fatty acid products, coronary artery rings with endothelium were contracted with U46619 before the addition of increasing concentrations of an omega-3 fatty acid product. Both EPA and DHA, the two major omega-3 fatty acid products, induced concentration-dependent relaxations in coronary artery rings with endothelium whereas no such effect was observed in those without endothelium (Figures 1A and B). EPA and DHA induced similar relaxations except at 0.4% (v/v) where relaxations to EPA were slightly but significantly greater than those to DHA ( $77.7 \pm 10.3$  and  $64.6 \pm 12.8\%$ , respectively, Figures 1A and B). Next, the ability of different EPA:DHA ratios to induce endothelium-dependent relaxations was determined. EPA:DHA with different ratios induced concentration-dependent relaxations in intact coronary artery rings (Figures 1A and B). However, relaxations to EPA:DHA 6:1 and 9:1 were significantly greater than those to EPA:DHA 3:1, 1:1, 1:3, 1:6, and 1:9 (Figures 1A and B). An increased relaxation was also observed in response to an EPA:DHA 6:1 product having a high EPA plus DHA content (694:121 mg/g) compared to one having a low content (352:65 mg/g, Figure 1C). Altogether, these findings indicate that omega-3 fatty acids are inducers of endothelium-dependent relaxations in coronary artery rings, and that this effect is dependent on the EPA:DHA ratio and on the EPA plus DHA amount.

### EPA:DHA 6:1 induces endothelium-dependent relaxations involving predominantly NO and also, to some extent, endothelium-dependent hyperpolarization

Next, experiments were performed to determine the role of the different endothelium-derived relaxing factors in the EPA:DHA 6:1-induced relaxation of the coronary artery. Relaxations to EPA:DHA 6:1 were slightly but significantly inhibited by L-NA (a competitive inhibitor of eNOS), not affected by inhibitors of endothelium-dependent hyperpolarization-mediated responses, TRAM 34 plus apamin (inhibitors of  $Ca^{2+}$ -dependent potassium channels of intermediate and low conductance  $IK_{Ca}$  and  $SK_{Ca}$ , respectively), and abolished by the combination of L-NA plus TRAM 34 and apamin (Figure 1D). In addition, they were not affected by indomethacin (cyclooxygenase inhibitor; relaxations at 0.4% v/v were  $94.8 \pm 3.2\%$  and  $95.8 \pm 3.2\%$  in the absence and presence of indomethacin, *n* = 5). These findings indicate that EPA:DHA 6:1 induces endothelium-dependent relaxations in coronary artery rings involving predominantly NO, and also, to some extent, EDH but not vasoactive prostanoids.

### EPA:DHA 6:1 induces redox-sensitive NO-mediated relaxations involving Src kinase, PI3-kinase, and MAPKs


Besides the calcium-calmodulin complex, the redox-sensitive Src/PI3-kinase/Akt and MAPKs pathways have also been shown to activate eNOS in endothelial cells in response to both physiological activators such as shear stress and estrogens, and natural products such as grape-derived polyphenols and tea catechins [18,20-23]. Since the EPA-induced activation of eNOS is a calcium-independent event [12], experiments were performed to determine the role of redox-sensitive Src/PI3-kinase/Akt and MAPKs pathways in the NO-mediated relaxations to EPA:DHA 6:1. For this purpose, all subsequent experiments were performed in the presence of indomethacin and TRAM34 plus apamin to rule out the formation of vasoactive prostanoids and EDH-mediated responses, respectively. Relaxations to EPA:DHA 6:1 were significantly reduced by the antioxidant N-acetylcysteine, the membrane permeant mimetic of SOD and the membrane permeant catalase (MnTMPyP and PEG-catalase, respectively) and native SOD and catalase (Figures 2 A and B). In addition, they were also reduced by PP2 (a Src kinase inhibitor), wortmannin (a PI3-kinase inhibitor), SB203580 (a p38 MAPK inhibitor), SP600125 (a JNK inhibitor), PD98059 (a MEK inhibitor, Figures 2C and D). In contrast to EPA:DHA 6:1, relaxations to bradykinin were not affected by either PD98059, SP600125, MnTMPyP, PP2 or wortmannin (data not shown) [24-26]. Thus, these findings indicate that the EPA:DHA 6:1-induced NO-mediated relaxation is critically dependent on a redox-

**Table 1.** Characteristics of the different omega-3 fatty acid products.

Fatty acids	Purity (in %)	Ratio EPA:DHA	Content of Omega-3 EPA-DHA as EE		Sum Omega-3 as EE (mg/g)
			EPA (mg/g)	DHA (mg/g)	
EPA:DHA	93.4	1:1	476	386	934
EPA:DHA	91.3	6:1	694	121	913
EPA:DHA	46.0	6:1	352	65	460
EPA	99.5		991		995
DHA	98.6			945	986

Fatty acids contents are given in mg/g expressed as ethyl ester (EE).

doi:10.1371/journal.pone.0105102.t001




**Figure 1. Omega-3 fatty acids cause endothelium-dependent relaxations, which are dependent on the EPA:DHA ratio and the EPA plus DHA content.** (A, B, C) Coronary artery rings with endothelium were contracted with U46619 before the addition of increasing concentrations of an omega-3 fatty acid product. (D) Rings were incubated with either L-NA (NO synthase inhibitor), TRAM34 plus apamin (inhibitors of endothelium-dependent hyperpolarization) or the three inhibitors together for 30 min before the contraction to U46619 and the subsequent relaxation to EPA:DHA 6:1. All experiments were performed in the presence of indomethacin (10  $\mu$ M) to prevent the formation of vasoactive prostanoids. Results are expressed as means  $\pm$  SEM of 5 different experiments. \* $P$ <0.05 versus (A, B) EPA:DHA 6:1 with endothelium, (C) EPA:DHA 6:1 (694:121 mg/g) with endothelium, and (D) control. doi:10.1371/journal.pone.0105102.g001

sensitive event involving Src kinase/PI3-kinase and MAPKs pathways.

**EPA:DHA 6:1 induces the redox-sensitive Src/PI3-kinase/Akt- and MAPKs-dependent phosphorylation of eNOS at Ser 1177**

To obtain further evidence that EPA:DHA 6:1 activates Src, Akt, MAPKs and eNOS, endothelial cells were cultured from coronary arteries and the phosphorylation level of these proteins at activator sites (Ser17 for p-Src, Ser473 for p-Akt, Thr183/Tyr185 for p-JNK, Thr180/Tyr182 for p-p38 MAPK, Thr202/Tyr204 for p-ERK, and Ser1177 for p-eNOS) was examined by Western blot analysis. EPA:DHA 6:1 at 0.4% v/v caused a time-dependent phosphorylation of Src, Akt and eNOS with significantly increased

levels observed within 5 min and, thereafter, these signals remained elevated at least up to 2 h (Figure 3A). In addition, EPA:DHA 6:1 increased to a greater extent the phosphorylation of Src, Akt and eNOS than EPA:DHA 1:1 (Figure 3B). EPA:DHA 6:1 (0.4% v/v) also significantly increased the phosphorylation level of p38 MAPK, JNK and ERK (Figure 4B). The possibility that the redox-sensitive event triggers the Src/PI3-kinase/Akt and MAPKs pathways leading to eNOS phosphorylation was examined. The EPA:DHA 6:1-induced phosphorylation of Src, Akt and eNOS was markedly reduced by MnTMPyP and PEG-catalase and not significantly affected by native SOD and catalase (Figure 4A). The EPA:DHA 6:1-induced phosphorylation of JNK, p38 MAPK and ERK was also markedly reduced by MnTMPyP whereas SOD, catalase had only minor effects



**Figure 2. The EPA:DHA 6:1-induced endothelium-dependent NO-mediated relaxation involves a redox-sensitive event.** Coronary artery rings with endothelium were incubated either with (A) an antioxidant (N-acetylcysteine), an extracellular scavenger of superoxide anions (SOD), a membrane permeant mimetic of superoxide dismutase (MnTMPyP), (B) extracellular and membrane permeant analogue of catalase (catalase and PEG-catalase, respectively), (C) an inhibitor of either Src kinase (PP2) or PI3-kinase (wortmannin), (D) an inhibitor of either p38 MAPK (SB203580), MEK (PD98059) or JNK (SP600125) for 30 min before the addition of U46619 and the subsequent relaxation to EPA:DHA 6:1. All experiments were performed in the presence of indomethacin (10  $\mu$ M) and TRAM34 plus apamin (both at 100 nM) to rule out the formation of vasoactive prostanoids and endothelium-dependent hyperpolarization-mediated responses, respectively. Results are presented as means  $\pm$  SEM of 5 different experiments. \* $P$ <0.05 versus control.


doi:10.1371/journal.pone.0105102.g002

(Figure 4B). PEG-catalase did not significantly affect the phosphorylation of JNK and p38 MAPK but reduced that of ERK (Figure 4B). Thus, these findings suggest that intracellular ROS, in particular superoxide anions, are upstream activators of the Src/PI3-kinase/Akt and MAPKs pathways leading to eNOS activation in response to EPA:DHA 6:1.

#### EPA:DHA 6:1 induces a pro-oxidant response in coronary artery endothelial cells

Experiments were performed to determine whether EPA:DHA 6:1 stimulates the endothelial formation of ROS in coronary artery sections using the redox-sensitive fluorescent probe dihydroethidium (DHE). EPA:DHA 6:1 markedly increased the fluorescence signal in the endothelium whereas the signal in the smooth muscle

was minimally affected (Figure 5A). The stimulatory effect of EPA:DHA 6:1 was markedly reduced by MnTMPyP and PEG-SOD, and, to a lesser extent, by PEG-catalase, and not affected by SOD and catalase (Figure 5A). In addition, EPA:DHA 6:1 significantly increased the formation of superoxide anions in cultured endothelial cells as assessed using ESR and the cell permeable spin probe CMH, and this effect was inhibited by SOD and PEG-SOD (Figure 5B). Moreover, EPA:DHA 6:1 significantly increased the formation of hydrogen peroxide in cultured endothelial cells as assessed using the Amplex Red based assay, and this effect was inhibited by PEG-catalase but not by catalase (Figure 5C). Altogether, these findings indicate that EPA:DHA 6:1 induces a pro-oxidant response in endothelial cells involving both superoxide anions and hydrogen peroxide.




**Figure 3. EPA:DHA 6:1 induces the phosphorylation of Src, Akt and eNOS in porcine coronary artery endothelial cells in a time- and EPA:DHA ratio-dependent manner.** (A) Cultured endothelial cells were exposed to EPA:DHA 6:1 (0.4% v/v) for the indicated times at 37°C. (B) Cultured endothelial cells were exposed to products with either an EPA:DHA ratio of 6:1 or 1:1 at the concentration of 0.4% (v/v) for 30 min at 37°C. Representative immunoblots of p-Src, p-Akt and p-eNOS proteins as assessed by Western blot analysis (top), and corresponding cumulative data (bottom). Results are shown as means  $\pm$  SEM of 3 different experiments. \* $P$ <0.05 versus control, \*\* $P$ <0.05 versus EPA:DHA 6:1 treatment. doi:10.1371/journal.pone.0105102.g003

## Discussion

The major findings of the present study indicate that the ability of omega-3 fatty acid products to cause endothelium-dependent relaxations involving predominantly NO and also, to some extent endothelium-dependent hyperpolarization in coronary arteries, depends on the formulation and the content of EPA and DHA, and that the EPA:DHA 6:1 is a superior formulation than EPA:DHA 1:1. They further indicate that the stimulatory effect of EPA:DHA 6:1 leading to eNOS activation involves the Src/PI3-kinase/Akt and MAPKs pathways with the subsequent phosphorylation of eNOS at the activator site Ser1177. Moreover, a redox-sensitive intracellular event involving superoxide anions and hydrogen peroxide has been identified as an early event triggering the activation of the Src/PI3-kinase/Akt and MAPKs pathways in endothelial cells. Thus, endothelial cells appear to act as special

sensors of omega-3 fatty acids in the arterial wall to increase in a sustained manner the eNOS-derived formation of NO, and, hence, vascular protection.

The evaluation of the ability of different omega-3 fatty acid products to relax porcine coronary arteries has indicated that EPA, DHA and formulations with different ratios of EPA:DHA induced pronounced relaxations in intact arterial preparations whereas they had little effects in rings without endothelium. Moreover, although EPA and DHA alone evoked similar concentration-relaxation curves up to 0.2% (v/v), their combination in the formulation EPA:DHA 6:1 and 9:1 demonstrated superiority to the formulation EPA:DHA 3:1, 1:1, 1:3, 1:6 and 1:9. The mechanism accounting for such a synergistic effect remains unclear. Potential explanations could be that the integration of omega-3 fatty acids into membrane lipids and/or that the stability




**Figure 4. The EPA:DHA 6:1-induced phosphorylation of Src, Akt, eNOS, JNK, p38 MAPK and ERK is a redox-sensitive event involving predominantly superoxide anions in endothelial cells.** Endothelial cells were incubated either with solvent, MnTMPyP (100  $\mu$ M), SOD (500 U/ml) or PEG-catalase (500 U/ml) for 30 min before the addition of EPA:DHA 6:1 (0.4% v/v) for 30 min. Representative immunoblots for detection of (A) p-Src, p-Akt and p-eNOS and (B) p-JNK, p-p38 MAPK and p-ERK (top), and corresponding cumulative data (bottom). Results are shown as means  $\pm$  SEM of 3-4 different experiments except for p-ERK (n=2). \* $P$ <0.05 versus control, # $P$ <0.05 versus EPA:DHA 6:1 treatment. doi:10.1371/journal.pone.0105102.g004

of the omega-3 fatty acids are dependent on an optimal EPA:DHA ratio. Indeed, EPA but not DHA has also been shown to cause cerebral microvascular vasodilatation involving arachidonic acid metabolites [15], whereas DHA but not EPA supplementation improved forearm microvascular reactivity in overweight hyperlipidemic men [16]. The characterization of the omega-3 fatty acid-induced endothelium-dependent relaxation has indicated a predominant role of NO, and also, to some extent, endothelium-dependent hyperpolarization whereas vasoactive prostanoids are not involved. Previous studies have also shown an increased endothelium-dependent relaxation or vasodilatation in response to the acute addition of EPA [12], and the chronic administration of EPA (3.5 g daily) and DHA (1.5 g daily) in pigs [7,27], and an increased vasodilatation to EPA (1.8 g daily) in humans [28].

Although omega-3 fatty acids are potent inducers of the endothelial formation of NO, the underlying mechanism remains poorly characterized. Activation of eNOS can be triggered by elevations in intracellular calcium by calcium elevating agonists such as bradykinin and thrombin, subsequently to the binding of the calcium-calmodulin complex to eNOS and the dissociation of the enzyme from its inhibitor caveolin [29]. Although EPA

induced elevations in intracellular calcium in endothelial cells, these responses were very small and did not contribute to eNOS activation [12]. EPA and DHA have also been shown to alter caveolae microenvironment not only by modifying membrane lipid composition but also by changing distribution of major structural proteins, which may eventually lead to eNOS activation [30,31]. The present findings provide evidence for an additional mechanism of eNOS activation by omega-3 fatty acids via the Src/PI3-kinase/Akt and MAPKs pathways leading to the phosphorylation of eNOS at the activator site Ser1177. Indeed, inhibitors of either Src kinase, PI3-kinase, JNK, p38 MAPK or MEK reduced NO-mediated relaxations to EPA:DHA 6:1, and EPA:DHA 6:1 increased the phosphorylation level of activator sites of Src, Akt, JNK, p38 MAPK, ERK and eNOS in endothelial cells. Of interest, the Src/PI3-kinase/Akt pathway has also been shown to mediate the stimulatory effect of other natural products containing high levels of polyphenols such as grape-derived products and tea catechins on eNOS [17,18,22,23]. Similarly to polyphenols [17,18,22,23], the intracellular formation of ROS, in particular superoxide anions and hydrogen peroxide, mediates the stimulatory effect on the Src/PI3-kinase/Akt and MAPKs



**Figure 5. EPA:DHA 6:1 induces the formation of ROS involving superoxide anions and hydrogen peroxide in endothelial cells.** A) The EPA:DHA 6:1-induced oxidative stress was assessed in coronary artery sections with endothelium using the redox-sensitive probe dihydroethidium (2.5 μM for 30 min). The EPA:DHA 6:1-induced formation of B) superoxide anions in cultured endothelial cells was assessed using electron spin resonance and the cell permeable spin probe CMH, and C) of hydrogen peroxide using the Amplex Red assay. B) Top: Representative original traces showing the three signals used for quantification (vertical bars). Bottom: Corresponding cumulative data. Means of the amplitude (height) of the three signals was measured for each spectrum. Results are shown as means ± SEM of 3-5 different experiments. \* $P < 0.05$  versus control, # $P < 0.05$  versus EPA: DHA 6:1 treatment. doi:10.1371/journal.pone.0105102.g005

pathways leading to eNOS activation in response to EPA:DHA 6:1. Indeed, inhibitors of intracellular oxidative stress and in particular the membrane permeant mimetic of SOD, MnTMPyP, markedly reduced the relaxation and the phosphorylation of Src, Akt, JNK, p38 MAPK, ERK and eNOS in response to EPA:DHA 6:1. Moreover, the redox-sensitive probe dihydroethidium provided direct evidence that EPA:DHA 6:1 induced the formation of ROS in the native endothelium whereas the signal in the smooth muscle was minimally affected. Furthermore, EPA:DHA 6:1-induced the generation of superoxide anions and hydrogen peroxide in cultured endothelial cells as assessed using the ESR technique with the cell permeable spin probe CMH and the Amplex Red assay, respectively. Native SOD and catalase did not affect the EPA:DHA 6:1-induced phosphorylation of target proteins and the increased DHE fluorescence signal in the native endothelium of coronary artery sections. However, both compounds reduced NO-mediated relaxations, such a difference is most likely due to different experimental conditions and in particular to the high partial oxygen pressure in the vascular reactivity studies (aerated with 95% O<sub>2</sub>) compared to that in

experiments using cultured endothelial cells (aerated with 18-20% O<sub>2</sub>) and coronary artery sections (ambient air, 21% O<sub>2</sub>). Besides endothelial cells, the induction of oxidative stress in response to omega-3 fatty acids has also been observed in several types of cancer cells including MCF-7 [32], and vascular smooth muscle cells to induce apoptosis [33]. The oxidative stress-induced apoptosis has been suggested to involve lipid peroxidation [32], and a calcium-regulated event [33].

In conclusion, the ability of omega-3 fatty acids to stimulate the endothelial formation of NO is dependent on the ratio and amount of the EPA:DHA formulation, and it involves an original pathway triggered by the intracellular oxidative stress leading to the activation of the Src/PI3-kinase/Akt and MAPKs pathways to increase ultimately eNOS activity.

### Author Contributions

Conceived and designed the experiments: CA VSK. Performed the experiments: FZ MA SR MB. Analyzed the data: FZ MA SR MB CA VSK. Wrote the paper: FZ MA CA VSK.

### References

- He K, Song Y, Daviglius ML, Liu K, Van Horn L, et al. (2004) Accumulated evidence on fish consumption and coronary heart disease mortality: a meta-analysis of cohort studies. *Circulation* 109: 2705–2711.
- Hu FB, Bronner L, Willett WC, Stampfer MJ, Rexrode KM, et al. (2002) Fish and omega-3 fatty acid intake and risk of coronary heart disease in women. *JAMA* 287: 1815–1821.
- GISSI (1999) Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial.



- Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto miocardico. *Lancet* 354: 447–455.
4. Galan P, Kesse-Guyot E, Czernichow S, Briancon S, Blacher J, et al. (2010) Effects of B vitamins and omega 3 fatty acids on cardiovascular diseases: a randomised placebo controlled trial. *BMJ* 341: c6273.
  5. Kromhout D, Giltay EJ, Geleijnse JM, Alpha Omega Trial G (2010) n-3 fatty acids and cardiovascular events after myocardial infarction. *N Engl J Med* 363: 2015–2026.
  6. Kromhout D, Yasuda S, Geleijnse JM, Shimokawa H (2012) Fish oil and omega-3 fatty acids in cardiovascular disease: do they really work? *Eur Heart J* 33: 436–443.
  7. Shimokawa H, Aarhus LL, Vanhoutte PM (1988) Dietary omega 3 polyunsaturated fatty acids augment endothelium-dependent relaxation to bradykinin in coronary microvessels of the pig. *Br J Pharmacol* 95: 1191–1196.
  8. Shimokawa H, Lam JY, Chesebro JH, Bowie EJ, Vanhoutte PM (1987) Effects of dietary supplementation with cod-liver oil on endothelium-dependent responses in porcine coronary arteries. *Circulation* 76: 898–905.
  9. de Berrazueta JR, Gomez de Berrazueta JM, Amado Senaris JA, Pena Sarabia N, Fernandez Viadero C, et al. (2009) A diet enriched with mackerel (*Scomber scombrus*)-derived products improves the endothelial function in a senior population (Prevencion de las Enfermedades Cardiovasculares: Estudio Santana—PECES project). *Eur J Clin Invest* 39: 165–173.
  10. Wang Q, Liang X, Wang L, Lu X, Huang J, et al. (2012) Effect of omega-3 fatty acids supplementation on endothelial function: a meta-analysis of randomized controlled trials. *Atherosclerosis* 221: 536–543.
  11. Stirban A, Nandreaan S, Gotting C, Tamler R, Pop A, et al. (2010) Effects of n-3 fatty acids on macro- and microvascular function in subjects with type 2 diabetes mellitus. *Am J Clin Nutr* 91: 808–813.
  12. Omura M, Kobayashi S, Mizukami Y, Mogami K, Todoroki-Ikeda N, et al. (2001) Eicosapentaenoic acid (EPA) induces Ca(2+)-independent activation and translocation of endothelial nitric oxide synthase and endothelium-dependent vasorelaxation. *FEBS Lett* 487: 361–366.
  13. Singh TU, Kathirvel K, Choudhury S, Garg SK, Mishra SK (2010) Eicosapentaenoic acid-induced endothelium-dependent and -independent relaxation of sheep pulmonary artery. *Eur J Pharmacol* 636: 108–113.
  14. Boulanger C, Schini VB, Hendrickson H, Vanhoutte PM (1990) Chronic exposure of cultured endothelial cells to eicosapentaenoic acid potentiates the release of endothelium-derived relaxing factor(s). *Br J Pharmacol* 99: 176–180.
  15. Ellis EF, Police RJ, Yancey LM, Grabeel MN, Heizer ML, et al. (1990) Effect of fish oil n-3 fatty acids on cerebral microcirculation. *Am J Physiol* 258: H1780–1785.
  16. Mori TA, Watts GF, Burke V, Hilde E, Puddey IB, et al. (2000) Differential effects of eicosapentaenoic acid and docosahexaenoic acid on vascular reactivity of the forearm microcirculation in hyperlipidemic, overweight men. *Circulation* 102: 1264–1269.
  17. Auger C, Kim JH, Chabert P, Chaabi M, Anselm E, et al. (2010) The EGCg-induced redox-sensitive activation of endothelial nitric oxide synthase and relaxation are critically dependent on hydroxyl moieties. *Biochem Biophys Res Commun* 393: 162–167.
  18. Anselm E, Chataigneau M, Ndiaye M, Chataigneau T, Schini-Kerth VB (2007) Grape juice causes endothelium-dependent relaxation via a redox-sensitive Src- and Akt-dependent activation of eNOS. *Cardiovasc Res* 73: 404–413.
  19. Ganster F, Burbhan M, de la Bourdonnaye M, Fizanne L, Douay O, et al. (2010) Effects of hydrogen sulfide on hemodynamics, inflammatory response and oxidative stress during resuscitated hemorrhagic shock in rats. *Crit Care* 14: R165.
  20. Fisslthaler B, Dimmeler S, Hermann C, Busse R, Fleming I (2000) Phosphorylation and activation of the endothelial nitric oxide synthase by fluid shear stress. *Acta Physiol Scand* 168: 81–88.
  21. Haynes MP, Sinha D, Russell KS, Collinge M, Fulton D, et al. (2000) Membrane estrogen receptor engagement activates endothelial nitric oxide synthase via the PI3-kinase-Akt pathway in human endothelial cells. *Circ Res* 87: 677–682.
  22. Kim JA, Formoso G, Li Y, Potenza MA, Marasciulo FL, et al. (2007) Epigallocatechin gallate, a green tea polyphenol, mediates NO-dependent vasodilation using signaling pathways in vascular endothelium requiring reactive oxygen species and Fyn. *J Biol Chem* 282: 13736–13745.
  23. Ndiaye M, Chataigneau M, Lobysheva I, Chataigneau T, Schini-Kerth VB (2005) Red wine polyphenol-induced, endothelium-dependent NO-mediated relaxation is due to the redox-sensitive PI3-kinase/Akt-dependent phosphorylation of endothelial NO-synthase in the isolated porcine coronary artery. *FASEB J* 19: 455–457.
  24. Ndiaye M, Chataigneau T, Andriantsitohaina R, Stoclet JC, Schini-Kerth VB (2003) Red wine polyphenols cause endothelium-dependent EDHF-mediated relaxations in porcine coronary arteries via a redox-sensitive mechanism. *Biochem Biophys Res Commun* 310: 371–377.
  25. Ndiaye M, Chataigneau T, Chataigneau M, Schini-Kerth VB (2004) Red wine polyphenols induce EDHF-mediated relaxations in porcine coronary arteries through the redox-sensitive activation of the PI3-kinase/Akt pathway. *Br J Pharmacol* 142: 1131–1136.
  26. Madeira SV, Auger C, Anselm E, Chataigneau M, Chataigneau T, et al. (2009) eNOS activation induced by a polyphenol-rich grape skin extract in porcine coronary arteries. *J Vasc Res* 46: 406–416.
  27. Shimokawa H, Vanhoutte PM (1989) Dietary omega 3 fatty acids and endothelium-dependent relaxations in porcine coronary arteries. *Am J Physiol* 256: H968–973.
  28. Tagawa H, Shimokawa H, Tagawa T, Kuroiwa-Matsumoto M, Hirooka Y, et al. (1999) Long-term treatment with eicosapentaenoic acid augments both nitric oxide-mediated and non-nitric oxide-mediated endothelium-dependent forearm vasodilatation in patients with coronary artery disease. *J Cardiovasc Pharmacol* 33: 633–640.
  29. Michel JB, Feron O, Sase K, Prabhakar P, Michel T (1997) Caveolin versus calmodulin. Counterbalancing allosteric modulators of endothelial nitric oxide synthase. *J Biol Chem* 272: 25907–25912.
  30. Li Q, Zhang Q, Wang M, Liu F, Zhao S, et al. (2007) Docosahexaenoic acid affects endothelial nitric oxide synthase in caveolae. *Arch Biochem Biophys* 466: 250–259.
  31. Li Q, Zhang Q, Wang M, Zhao S, Ma J, et al. (2007) Eicosapentaenoic acid modifies lipid composition in caveolae and induces translocation of endothelial nitric oxide synthase. *Biochimie* 89: 169–177.
  32. Kang KS, Wang P, Yamabe N, Fukui M, Jay T, et al. (2010) Docosahexaenoic acid induces apoptosis in MCF-7 cells in vitro and in vivo via reactive oxygen species formation and caspase 8 activation. *PLoS One* 5: e10296.
  33. Crnkovic S, Riederer M, Lechleitner M, Hallstrom S, Malli R, et al. (2012) Docosahexaenoic acid-induced unfolded protein response, cell cycle arrest, and apoptosis in vascular smooth muscle cells are triggered by Ca(2+)-dependent induction of oxidative stress. *Free Radic Biol Med* 52: 1786–1795.