

Complete Genome Sequences of One Human Respiratory Syncytial Antigenic Group A Virus from China and Its Four Mouse-Adapted Isolates

Ke Zhang,^{a,b} Jie He,^c Cun Li,^a Michael E. Bose,^c Kelly J. Henrickson,^c Jie Zhou,^a Bo-Jian Zheng^a

Department of Microbiology, University of Hong Kong, Hong Kong, China^a; Department of Parasitology, Guiyang Medical University, Guizhou, China^b; Department of Pediatrics, Medical College of Wisconsin, Milwaukee, Wisconsin, USA^c

In this study, one human respiratory syncytial antigenic group A virus (HRSV-A-GZ08-0) and its four BALB/c mouse-adapted isolates were sequenced and elucidated. Nineteen nucleotides were mutated between HRSV-A-GZ08-0 and the four mouse-adapted isolates.

Received 18 January 2015 Accepted 28 January 2015 Published 5 March 2015

Citation Zhang K, He J, Li C, Bose ME, Henrickson KJ, Zhou J, Zheng B-J. 2015. Complete genome sequences of one human respiratory syncytial antigenic group A virus from China and its four mouse-adapted isolates. *Genome Announc* 3(2):e00062-15. doi:10.1128/genomeA.00062-15.

Copyright © 2015 Zhang et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Bo-Jian Zheng, bzheng@hkucc.hku.hk.

Worldwide, human respiratory syncytial virus (HRSV) is a main cause of acute lower respiratory tract infection (ALRTI), which can be severe and lead to hospitalization for children under 5 years of age and immunocompromised adults (1). HRSV, known as a nonsegmented enveloped negative-strand RNA virus, belongs to the subfamily *Pneumovirinae* of the family *Paramyxoviridae* with antigenic groups A and B. Currently, antigenic group A is further divided into GA1 to GA7, NA1 to NA2, ON1, and SAA1 genotypes, while group B consists of BA1 to BA10, GB1 to GB4, and SAB1 to SAB3 (2). The initiate sites of 3' leader (Le) and 5' trailer (Tr) sequences in the HRSV genome are fully conserved (3). With strong promoters enclosed, Le plays a key role in replication, transcription, and encapsidation, whereas Tr focuses more on genome RNA generation via the mediation from viral RNA-dependent RNA polymerase (4, 5). The mutation in Le or Tr may cause significant modification in HRSV biology.

The HRSV strain HRSV-A-GZ08-0 was clinically isolated in Guangzhou Children's Hospital, Guangzhou, China, in 2008. Due to severe cytopathogenic effects observed *in vitro*, HRSV-A-GZ08-0 was considered as a candidate strain for establishing an HRSV infection mouse model, so this strain was serially passed in BALB/c mice for over 50 passages. Four mouse-adapted clones (HRSV-A-GZ08-10, HRSV-A-GZ08-12, HRSV-A-GZ08-18, and HRSV-A-GZ08-19) were isolated from one mouse because of high titer generation and purified by plaque assay. The full-length sequencing data of these five HRSV-A isolates were acquired by custom-designed primers, RNA extraction, reverse transcription PCR, and reading by 3730 DNA Analyzer (Thermo, Fisher Scientific, Waltham, MA, USA). The DNA 7500 kit and 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA) were used as quality controls for concentration of amplicon. Sequence assembly and consensus calling were performed using SeqMan Pro of the Lasergene 11 program suite (DNASTAR, Madison, WI, USA) (6). The nucleotide and protein annotations were presented with GenBank reference KC731483 and CLC Genomic Workbench version 7.0 (Qiagen, Hilden, Germany).

The five newly sequenced HRSVs possessed genomes from 15,193 bp to 15,201 bp, including Le and partial Tr regions. Eleven open reading frames were deduced from genomes for expression of NS1, NS2, N, P, M, SH, G, F, M2-1, M2-2, and L proteins. The four mouse-adapted isolates had nearly identical nucleotide information. Compared with HRSV-A-GZ08-0, they all possessed 5, 4, 3, and 2 nucleotide mutations in G, F, L, and NS1 genes, while N, P, and M genes exhibited 1 mutation, respectively. Notably, 1 mutation (A42G) was found in the Le region of all four mouse isolates and another mutation (A37G) occurred only in HRSV-A-GZ08-19. The nucleotide sequence identities of HRSV-A-GZ08-0 to GenBank whole-genome information of HRSV-A were from 96.97% to 97.30% (7, 8). The phylogenetic analysis showed that HRSV-A-GZ08-0 was assigned as GA2 genotype.

The genome information of HRSV-A-GZ08-0 and its four mouse-adapted isolates would provide valuable information for HRSV mice model research.

Nucleotide sequence accession numbers. The complete sequences of HRSV-A-GZ08-0, HRSV-A-GZ08-10, HRSV-A-GZ08-12, HRSV-A-GZ08-18, and HRSV-A-GZ08-19 were deposited in GenBank under the accession numbers [KP218910](https://pubmed.ncbi.nlm.nih.gov/26119745/), [KP119745](https://pubmed.ncbi.nlm.nih.gov/26119745/), [KP119746](https://pubmed.ncbi.nlm.nih.gov/26119746/), [KP119747](https://pubmed.ncbi.nlm.nih.gov/26119747/), and [KP119748](https://pubmed.ncbi.nlm.nih.gov/26119748/), respectively.

ACKNOWLEDGMENTS

This study was partially supported by the Research Fund for the Control of Infectious Diseases (grant 10091222) (now named the Health and Medical Research Fund), Hong Kong SAR.

REFERENCES

- Nair H, Nokes DJ, Gessner BD, Dherani M, Madhi SA, Singleton RJ, O'Brien KL, Roca A, Wright PF, Bruce N, Chandran A, Theodoratou E, Sutanto A, Sedyaningsih ER, Ngama M, Muniyoki PK, Kartasmita C, Simões EA, Rudan I, Weber MW, Campbell H. 2010. Global burden of acute lower respiratory infections due to respiratory syncytial virus in young children: a systematic review and meta-analysis. *Lancet* 375: 1545–1555. [http://dx.doi.org/10.1016/S0140-6736\(10\)60206-1](http://dx.doi.org/10.1016/S0140-6736(10)60206-1).

2. Valley-Omar Z, Muloiwa R, Hu N-C, Eley B, Hsiao N-Y. 2013. Novel respiratory syncytial virus subtype ON1 among children, Cape Town, South Africa, 2012. *Emerg Infect Dis* 19:668–670. <http://dx.doi.org/10.3201/eid1904.121465>.
3. Noton SL, Fearn R. 2011. The first two nucleotides of the respiratory syncytial virus antigenome RNA replication product can be selected independently of the promoter terminus. *RNA* 17:1895–1906. <http://dx.doi.org/10.1261/rna.2813411>.
4. Fearn R, Collins PL, Peeples ME. 2000. Functional analysis of the genomic and antigenomic promoters of human respiratory syncytial virus. *J Virol* 74: 6006–6014. <http://dx.doi.org/10.1128/JVI.74.13.6006-6014.2000>.
5. Fearn R, Peeples ME, Collins PL. 2002. Mapping the transcription and replication promoters of respiratory syncytial virus. *J Virol* 76:1663–1672. <http://dx.doi.org/10.1128/JVI.76.4.1663-1672.2002>.
6. Rebuffo-Scheer C, Bose M, He J, Khaja S, Ulatowski M, Beck ET, Fan J, Kumar S, Nelson MI, Henrickson KJ. 2011. Whole genome sequencing and evolutionary analysis of human respiratory syncytial virus A and B from Milwaukee, WI 1998–2010. *PLoS One* 6:e25468. <http://dx.doi.org/10.1371/journal.pone.0025468>.
7. Choudhary ML, Wadhwa BS, Jadhav SM, Chadha MS. 2013. Complete genome sequences of two human respiratory syncytial virus genotype A Strains from India, RSV-A/NIV1114046/11 and RSV-A/NIV1114073/11. *Genome Announc* 1(4):e00165-13. <http://dx.doi.org/10.1128/genomeA.00165-13>.
8. Lee WJ, Kim YJ, Kim DW, Lee HS, Lee HY, Kim K. 2012. Complete genome sequence of human respiratory syncytial virus genotype A with a 72-nucleotide duplication in the attachment protein G gene. *J Virol* 86: 13810–13811. <http://dx.doi.org/10.1128/JVI.02571-12>.