

ERRATUM

Open Access



Erratum: Pregnancy complications and birth outcomes among women experiencing nausea only or nausea and vomiting during pregnancy in the Norwegian Mother and Child Cohort Study

Arthur Chortatos^{1,2*}, Margaretha Haugen³, Per Ole Iversen^{2,4}, Åse Vikanes⁵, Malin Eberhard-Gran^{6,7}, Elisabeth Krefting Bjelland^{6,7}, Per Magnus⁵ and Marit B. Veierød^{1,2}

Erratum

After publication of the original article [1] it came to the publisher's attention that there were errors in the formatting of Table five. The first Outcome 'Emergency caesarean delivery' was incorrectly separated by a line break. The second and third Outcomes 'Preterm birth' and 'Birth type (other presentations than normal cephalic)' were formatted on the incorrect lines. These errors have been corrected in Table 5 below.

Author details

¹Oslo Centre for Biostatistics and Epidemiology, Institute of Basic Medical Sciences, University of Oslo, Blindern N-0317, Oslo, Norway. ²Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo, Blindern N-0317, Oslo, Norway. ³Division of Environmental Medicine, Norwegian Institute of Public Health, Nydalen N-0403, Oslo, Norway. ⁴Department of Hematology, Oslo University Hospital, Nydalen N-0424, Oslo, Norway. ⁵Division of Epidemiology, Norwegian Institute of Public Health, Nydalen N-0403, Oslo, Norway. ⁶Department of Psychosomatics and Health Behaviour, Norwegian Institute of Public Health, Nydalen N-0403, Oslo, Norway. ⁷Health Services Research Unit, Akershus University Hospital, Lørenskog N-1478, Norway.

Received: 17 July 2015 Accepted: 17 July 2015

Published online: 13 August 2015

Reference

1. Chortatos A, Haugen M, Iversen PO, Vikanes Å, Eberhard-Gran M, Bjelland EK, et al. Pregnancy complications and birth outcomes among women experiencing nausea only or nausea and vomiting during pregnancy in the Norwegian Mother and Child Cohort Study. *BMC Pregnancy Childbirth*. 2015;15:138. doi:10.1186/s12884-015-0580-6.

* Correspondence: athanasc@medisin.uio.no

¹Oslo Centre for Biostatistics and Epidemiology, Institute of Basic Medical Sciences, University of Oslo, Blindern N-0317, Oslo, Norway

²Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo, Blindern N-0317, Oslo, Norway

Table 5 Odds ratios^a of birth outcomes in relation to group (symptom-free (SF), nausea only (NP), and nausea and vomiting (NVP))

Outcome	Group	n (%) ^b	Crude		Adjusted ^d	
			cOR (95 % CI) ^c	P value	aOR (95 % CI)	P value
Emergency caesarean delivery	SF	1220 (9.4)	1.00	<0.001	1.00	0.07
	NP	1413 (7.6)	0.79 (0.73-0.86)		0.91 (0.84-0.99)	
	NVP	1302 (8.4)	0.88 (0.81-0.96)		0.97 (0.90-1.06)	
Preterm birth	SF	751 (5.4)	1.00	<0.001	1.00	0.02
	NP	879 (4.4)	0.81 (0.73-0.89)		0.86 (0.78-0.96)	
	NVP	800 (4.8)	0.88 (0.80-0.98)		0.91 (0.82-1.01)	
Birth type (other presentations than normal cephalic)	SF	1372 (10.3)	1.00	<0.001	1.00	0.001
	NP	1780 (9.3)	0.89 (0.82-0.96)		0.97 (0.90-1.04)	
	NVP	1363 (8.5)	0.81 (0.75-0.87)		0.87 (0.80-0.94)	
Apgar score after 5 minutes (<7) ^e						
Male infants	SF	124 (1.6)	1.00	<0.001	1.00	0.006
	NP	93 (0.9)	0.56 (0.43-0.73)		0.65 (0.49-0.85)	
	NVP	90 (1.1)	0.69 (0.53-0.90)		0.75 (0.57-0.99)	
Female infants	SF	68 (1.1)	1.00	<0.05	1.00	0.03
	NP	120 (1.2)	1.13 (0.84-1.52)		1.26 (0.93-1.70)	
	NVP	74 (0.9)	0.78 (0.56-1.09)		0.85 (0.61-1.19)	
Low birth weight	SF	339 (2.5)	1.00	<0.001	1.00	0.001
	NP	326 (1.7)	0.66 (0.57-0.77)		0.73 (0.60-0.88) ^f	
	NVP	305 (1.9)	0.74 (0.64-0.87)		0.72 (0.60-0.88) ^f	
Small for gestational age (SGA)	SF	1657 (12.0)	1.00	<0.001	1.00	<0.001
	NP	1908 (9.6)	0.78 (0.73-0.84)		0.78 (0.73-0.84) ^g	
	NVP	1694 (10.2)	0.84 (0.78-0.90)		0.87 (0.81-0.93) ^g	
Birth defects	SF	674 (4.9)	1.00	0.24	1.00	0.15
	NP	1022 (5.1)	1.06 (0.96-1.17)		1.10 (1.00-1.22)	
	NVP	791 (4.8)	0.98 (0.88-1.09)		1.03 (0.93-1.15)	
Gender of infant (female)	SF	6212 (45.0)	1.00	<0.001	1.00	<0.001
	NP	9736 (49.0)	1.18 (1.13-1.23)		1.18 (1.13-1.23) ^h	
	NVP	8625 (52.1)	1.33 (1.27-1.39)		1.34 (1.28-1.40) ^h	

^aOR, odds ratio, logistic regression analysis^bNumber (%) of women with outcome^cCI, confidence interval^dAdjusted for age, body mass index (BMI), smoking during pregnancy, parity, education, and gender^eSignificant interaction between Apgar score after 5 min (<7) and gender of infant ($P = 0.001$)^fAdjusted for age, BMI, smoking during pregnancy, parity, education, gender, gestational length, and energy intake^gAdjusted for age, BMI, smoking during pregnancy, education, and gender^hAdjusted for age, BMI, smoking during pregnancy, parity, and education