

Research

Open Access

Microglial responses to amyloid β peptide opsonization and indomethacin treatment

Ronald Strohmeyer, Carl J Kovelowski, Diego Mastroeni, Brian Leonard, Andrew Grover and Joseph Rogers*

Address: L.J. Roberts Center, Sun Health Research Institute, 10515 West Santa Fe Drive, Sun City, AZ 85351 USA

Email: Ronald Strohmeyer - RWStrohmeyer@NNU.edu; Carl J Kovelowski - cjkovelowski@yahoo.com;

Diego Mastroeni - diego.mastroeni@sunhealth.org; Brian Leonard - brian.leonard@sunhealth.org;

Andrew Grover - andrew.grover@sunhealth.org; Joseph Rogers* - joseph.rogers@sunhealth.org

* Corresponding author

Published: 19 August 2005

Received: 18 June 2005

Journal of Neuroinflammation 2005, 2:18 doi:10.1186/1742-2094-2-18

Accepted: 19 August 2005

This article is available from: <http://www.jneuroinflammation.com/content/2/1/18>

© 2005 Strohmeyer et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Recent studies have suggested that passive or active immunization with anti-amyloid β peptide (A β) antibodies may enhance microglial clearance of A β deposits from the brain. However, in a human clinical trial, several patients developed secondary inflammatory responses in brain that were sufficient to halt the study.

Methods: We have used an in vitro culture system to model the responses of microglia, derived from rapid autopsies of Alzheimer's disease patients, to A β deposits.

Results: Opsonization of the deposits with anti-A β IgG 6E10 enhanced microglial chemotaxis to and phagocytosis of A β , as well as exacerbated microglial secretion of the pro-inflammatory cytokines TNF- α and IL-6. Indomethacin, a common nonsteroidal anti-inflammatory drug (NSAID), had no effect on microglial chemotaxis or phagocytosis, but did significantly inhibit the enhanced production of IL-6 after A β opsonization.

Conclusion: These results are consistent with well known, differential NSAID actions on immune cell functions, and suggest that concurrent NSAID administration might serve as a useful adjunct to A β immunization, permitting unfettered clearance of A β while dampening secondary, inflammation-related adverse events.

Background

Chemotactic and phagocytic responses of microglia to amyloid β peptide (A β) have been inferred from postmortem autopsy evaluations [1-3], animal studies [4,5], and an in vitro model in which cultured rodent microglia were placed directly on Alzheimer's disease (AD) cortical sections [5,6]. Although these valuable experiments confirm that microglia cluster around and may help clear A β deposits, new questions have arisen concerning the effects

of various agents on these microglial interactions with A β . In particular, several studies have indicated that the opsonization of A β deposits with anti-A β antibodies facilitates microglia-mediated A β clearance [6,7]. Here, binding of the antibodies to the A β target presumably enhances microglial recognition of and subsequent responses to the target through Fc receptors expressed by the microglia [6,7]. Based on these results, it has been suggested that microglial responses to A β might represent so

beneficial an inflammatory action that anti-inflammatory drugs might actually be detrimental as a treatment for AD [8]. Alternatively, multiple epidemiologic studies [9,10] have reported decreased risk for AD in persons who take common nonsteroidal anti-inflammatory drugs (NSAIDs).

Over the last decade, our laboratory has developed reliable methods for culturing microglia from rapid (< 4 hour) brain autopsies of AD patients [11,12]. These cultures uniquely match the species, developmental stage, and disease state of AD subjects, and provide the ready experimental manipulability that is helpful in assessing complex physiologic processes such as chemotaxis, phagocytosis, secretory activity, and drug responses. In order to quantitatively assay these processes in the context of microglial interactions with A β , we seeded AD microglial cultures into wells containing pre-aggregated A β 1-42 spots dried down to the well floor. Subsequent experiments measured migration of the cells to the A β spots, phagocytosis of the A β spots, pro-inflammatory cytokine secretion, and the effects on these processes when A β spots were opsonized with an anti-A β antibody or when microglia were treated with a common nonsteroidal anti-inflammatory drug (NSAID), indomethacin. Overall, opsonization with A β antibody enhanced microglial migration to and phagocytosis of A β . Indomethacin had little to no effect on these responses, but did significantly inhibit microglial secretion of IL-6.

Methods

AD microglia cultures

Cultures of microglia from rapid (< 4 hours) autopsies of six antemortem-evaluated, neuropathologically-confirmed AD patients were prepared using our previously published methods [11,12]. By immunoreactivity, these cultures are consistently negative for neuron, astrocyte, oligodendrocyte, and fibroblast markers, consistently positive for multiple markers of activated microglia, and readily maintained at purities of 98% or higher [11,12]. Microglia cultures from all six AD patients were used for biochemical assays. Additional cultures from one of these patients were used for quantitative evaluation of chemotaxis and phagocytosis, and additional cultures from two more of these patients were used for qualitative replication of the chemotaxis and phagocytosis results. At 3–7 days post-plating, the microglia were trypsinized and replated at 50,000 cells/well in 12-well plates. Prior to replating, 2 μ l of a 1 mM solution of A β 1-42 (Bachem) in PBS (pH 7.4) was dried down to the well floor. Each well received two such A β spots, and there were three wells per experimental condition, so that a total of six A β spots were quantified per experimental condition. Serum-free medium was used throughout the experiments. Control

wells containing no A β or no microglia were also prepared.

Treatment with anti-A β antibody

Prior to seeding with microglia, selected wells were pre-treated with vehicle (medium) only or with 10 μ g/ml 6E10 (Signet Laboratories), a mouse monoclonal antibody directed against the first 17 (N-terminal) amino acids in the A β sequence. In some experiments, a 2 μ g/ml concentration of 6E10 was included in order to evaluate effects at a lower dose.

Treatment with indomethacin

Prior to seeding with microglia, selected wells were pre-treated with vehicle (medium) only or with 1.0 μ g/ml indomethacin. Indomethacin, at 1.0 μ g/ml, and vehicle were also replenished at Days 3, 6, and 9 in the course of medium changes. The 1.0 μ g/ml indomethacin concentration is at the upper end of the physiologically normal range achieved in blood after therapeutic doses of the drug [13], and was chosen to insure that any failure of indomethacin to affect chemotaxis to or phagocytosis of A β was not due to inadequate drug dosage. In some experiments, a 0.1 μ g/ml concentration of indomethacin, which is at the lower end of the physiologically normal range achieved in blood after therapeutic doses, was included in order to evaluate effects at a lesser concentration.

Cytochemistry and immunocytochemistry

For qualitative evaluations of microglial responses to A β , microglial cultures were briefly fixed with 4% buffered paraformaldehyde, then immunoreacted overnight with 1:1000 (0.5 μ g/ml) LN3 antibody (MP Biomedical) directed against the major histocompatibility complex type II cell surface glycoprotein, using our previously published methods [11,14,15]. Vectastain ABC kits (Vector Laboratories) were employed using the manufacturer's protocols to detect immunoreactivity with bright field optics. A β spots could be sufficiently resolved under these conditions by their modest opaqueness under bright field optics. To visualize A β spots in phagocytosis experiments, the wells were washed gently in distilled water (3 \times for 5 min each), incubated with 0.1% Thioflavine S (Sigma) for 10 min, washed once in distilled water (5 min), then dehydrated and fixed with 4% buffered paraformaldehyde. In additional experiments, A β immunocytochemistry was applied in selected wells so as to graphically illustrate A β removal and microglial uptake of A β . In these studies, microglial cultures with A β spots were briefly fixed with 4% buffered paraformaldehyde and incubated overnight with 1:1000 (1 μ g/ml) anti-A β antibody 4G8 (Signet Laboratories). Detection of immunoreactivity was accomplished using Vectastain ABC kits (Vector Laboratories) and the manufacturer's suggested protocols.

Microglial migration to A β spots

Microglial cultures were assessed on Day 3 and Day 9 after initial plating. Each A β spot was visualized under phase contrast optics at 100 \times (10 \times objective), and photomontages were made of the spot and surrounding area out to a radius of 2 mm from the spot perimeter. A grid was then placed over the photomontages. The number and percentages of microglia within four 500 μ m \times 500 μ m (0.25 mm²) grid squares centered on the A β spot and within sets of four 500 μ m \times 500 μ m squares at progressively greater distances from the spot were recorded. The distance intervals for the grid squares were 0, 500, 1000, 1500, and 2000 μ m from the A β spot, and each distance interval was measured in quadruplicate (Fig. 1). A total of 141,455 microglia were individually hand-counted in this way. Chemotaxis was evaluated by changes in the distributions of microglia relative to the A β spots over time, with relatively flat distributions indicative of little or no chemotaxis, and increasingly negative slopes to the distributions indicative of migration toward the A β spots (Fig. 1). Slopes of the distributions (m) were operationally defined as the "chemotactic index" [15] for each condition, and the statistical reliability of the measures was assessed with Pearson's Product Momentum (R) statistic and with analysis of variance (ANOVA) techniques. The simplest ANOVAs assessed, for each treatment condition, significant differences in the distributions of microglia over the progressive distance intervals from the A β spot, with percentage of microglia at a particular distance (grid square) as the dependent variable and distance from the A β spot (0, 500, 1000, 1500, and 2000 μ m) as the single factor. Pearson's R Statistic was then run to confirm that the alterations in microglial distributions were consistent with chemotaxis (i.e., showed a significant negative correlation with distance from A β) rather than some other response pattern. Dose dependence was evaluated using two-way ANOVAs, with percentage of microglia as the dependent variable, distance from the A β spot as the first factor, and drug dose as the second factor. Significant interactions of distance with drug dose thereby provided statistical evidence that the different drug doses differentially affected microglial distributions. A similar approach was taken for comparisons of different treatment conditions (e.g., anti-A β antibody exposure \pm indomethacin treatment). All data collection was by a technician blind to experimental condition.

Tests of microglial proliferation

BrdU staining kits (Zymed/Invitrogen) were applied to selected wells in order to assess whether shifts in microglial distributions over time might be due to differential proliferation of microglia relative to A β spots as opposed to migration of the cells. Staining with BrdU followed the manufacturer's recommended directions.

Microglial phagocytosis of A β spots

At Day 12 postplating, selected wells were histochemically reacted with Thioflavine S, as described earlier, and visualized at 100 \times (10 \times objective) with a confocal microscope. Using the ability of the confocal microscope to optically section an object at precise distances, the number of 10 μ m optical slices from the well floor to the top of the remaining A β spot was recorded by an investigator blinded to the experimental conditions imposed in each well. The data were then assessed statistically using 2-way ANOVAs, with spot thickness as the outcome measure, antibody treatment (vehicle only, 2 μ g/ml anti-A β IgG, or 10 μ g/ml anti-A β IgG) as the first factor, and NSAID treatment (vehicle only, 0.1 μ g/ml indomethacin, or 1.0 μ g/ml indomethacin) as the second factor.

Microglial secretion of cytokines

To assess the effects of opsonization with anti-A β antibodies, microglial cultures were preincubated with vehicle or 10 μ g/ml anti-A β monoclonal 6E10 followed by 4 hours exposure to 0 or 10 μ M preaggregated A β 1-42 (Bachem). Conditioned medium was then subjected to TNF- α ELISA (R&D Systems) using the manufacturer's protocols. To confirm the results with another pro-inflammatory cytokine, and to evaluate the interaction of indomethacin with antibody opsonization, microglial cultures were preincubated with vehicle or 10 μ g/ml 6E10, as before, but in the presence or absence of 1 μ g/ml indomethacin. After incubation for 4 hours with 0 or 10 μ M A β 1-42, the conditioned medium was subjected to IL-6 ELISA (R&D Systems) using the manufacturer's protocols.

Results

Microglial migration to A β spots

Overall and within each treatment condition there were shifts in microglial distributions, consistent with chemotaxis, that were both visually apparent (Figs. 2A, 2C) and statistically significant (Figs. 2B, 2D). By Day 3, the greatest concentrations of microglia were midway between the most distal and proximal points from the A β spots ($F_{\text{Distance}} = 40.1$, $P = 0.000$; $R = -.17$, $P = 0.000$; $m = -.016$) (Fig. 2B). By Day 9, the greatest concentrations of microglia were at or adjacent to the spots ($F_{\text{Distance}} = 99.2$, $P = 0.000$; $R = -.41$, $P = 0.000$; $m = -.041$) (Fig. 2D). Microglia seeded into wells without A β spots essentially remained randomly distributed throughout these time periods. Opsonization with anti-A β antibodies significantly enhanced chemotaxis-like shifts in microglial distributions, an effect that was especially prominent at Day 9 (Table 1) (Fig. 3). Indomethacin had no significant or obvious effect on changes in microglial distributions over time under any of the A β antibody treatment conditions. Indeed, the largest chemotactic index (slope) observed in the study occurred at the highest dose of indomethacin (1.0 μ g/ml indomethacin plus 10 μ g/ml anti-A β) (F_{Distance}




Figure 1

Paradigm for estimation of microglial chemotaxis to Aβ. Upper left panel shows a hypothetical example at Day 1, when microglia (black dots) are uniformly distributed relative to Aβ spots (gray circle). A plot of microglial density within 500 μm × 500 μm grid squares at increasing proximity to the spot (lower left) is therefore relatively flat, with a slope near 0, indicative of little or no migratory activity at this early time point. After 9 days (right panels), microglia are clustered over and around the Aβ spot, yielding a pronounced slope to the plot, consistent with chemotaxis to the Aβ. Previous studies have referred to such slopes as "chemotactic indices" [c.f., 15].



= 38.9, $P = 0.000$; $R = 0.69$, $P = 0.000$; $m = -.073$), and the second largest chemotactic index occurred at the second highest dose of indomethacin (0.1 μg/ml indomethacin plus 10 μg/ml anti-Aβ) ($F_{\text{Distance}} = 12.9$, $P = 0.000$; $R = -.53$, $P = 0.000$; $m = -.060$ (Fig. 3).

Differential proliferation versus chemotaxis

Proliferation of microglia more proximal to the Aβ spots, rather than true chemotaxis, did not explain the shifts in microglial distributions that were exhibited over time under the various treatment conditions. There was little to

no BrdU staining under any condition (not shown) and, in fact, there was a slight but significant decrease in microglial numbers in all treatment conditions and overall from Day 3 (mean microglial density/0.25 mm² grid square = 40.8 ± 0.3) to Day 9 (mean microglial density/0.25 mm² grid square = 37.8 ± 0.4) ($F_{\text{Overall}} = 34.5$, $P = 0.000$). Consistent with our previous experience, AD microglia stimulated with M-CSF as a positive control showed little to no evidence of proliferation. However, M-CSF-stimulated THP-1 cells (a monocyte line often used as a surrogate for microglia) that were run in parallel did show clear

DAY 3



DAY 9





Figure 2
Typical responses of cultured AD microglia to pre-aggregated Aβ1-42 spots dried down to the well floor. A) Micrograph of Aβ spot (light brown stain) and LN3 immunoreactive microglia (blue stain) 3 days postplating (vehicle control) (4 × objective). B) Graphic summary of microglial distributions at 3 days postplating (pooled data over all conditions). C) Parallel well 9 days postplating (vehicle control) (4 × objective). Wells seeded with microglia but without Aβ spots exhibited only random distributions of cells (not shown). D) Graphic summary of microglial distributions at Day 9 (pooled data over all conditions). Similar and highly significant shifts over time were observed in all treatment conditions when Aβ spots were present (see text).

Table 1: Effects of opsonization with anti-A β antibody 6E10 on chemotaxis-like changes in microglia distributions

	ANOVA		PEARSON'S		SLOPE m
	F	P	R	P	
Day 3					
0 μ g/ml anti-A β	3.7	0.007	-0.26	0.005	-0.022
2 μ g/ml anti-A β	2.5	0.040	-0.14	NS	-0.023
10 μ g/ml anti-A β	5.5	0.000	-0.27	0.003	-0.027
Dose dependence*	3.6	0.008			
Day 9					
0 μ g/ml anti-A β	5.6	0.000	-0.37	0.000	-0.040
2 μ g/ml anti-A β	11.2	0.000	-0.050	0.000	-0.051
10 μ g/ml anti-A β	16.4	0.000	-0.57	0.000	-0.056
Dose dependence*	2.3	0.050			

*Dose \times distance interaction term

proliferation under the same BrdU assay conditions (data not shown).

Microglial phagocytosis of A β

After incubation with microglia under the various experimental conditions, visible degradation of A β spots was apparent (Fig. 4A), whereas A β spots in wells not containing microglia remained visibly intact over the same time periods (Fig. 4B). Concurrent with degradation of the A β spots, microglia in contact with the spots became A β immunoreactive (Fig. 4A), whereas they exhibited little to no A β immunoreactivity prior to their being seeded into the wells (Fig. 4C). Opsonization of A β spots with 2 μ g/ml anti-A β antibody 6E10 ($F = 28.7$, $P = 0.006$) or 10 μ g/ml anti-A β antibody 6E10 ($F = 35.3$, $P = 0.004$) resulted in significantly smaller (thinner) A β spots compared to the vehicle control condition (Fig. 4D). These effects were not significantly or materially inhibited by indomethacin even at the highest, 1.0 μ g/ml indomethacin concentration (for 2 μ g/ml anti-A β \pm 1.0 μ g/ml indomethacin: $F = 0.3$, $P = 0.639$) (for 10 μ g/ml anti-A β plus \pm 1.0 μ g/ml indomethacin: $F = 0.9$, $P = 0.402$) (Fig. 4D).

Microglial secretion of cytokines

Consistent with our previous studies covering a wide range of cytokines, chemokines, and inflammatory toxins [12], exposure of microglia to A β significantly enhanced secretion of TNF- α (Fig. 5A) and IL-6 (Fig. 5B) compared to cultures that were not exposed to A β . Opsonization with 10 μ g/ml anti-A β antibody 6E10 significantly enhanced A β -induced TNF- α (Fig. 5A) and IL-6 secretion (Fig. 5B). Enhancement of IL-6 expression, however, was significantly decreased by indomethacin treatment (Fig. 5B). Cytokine secretion is typically a fairly rapid response that wanes over time. Presumably, cytokine receptive cells then undergo more long-lasting responses such as

enhanced chemotactic or phagocytic behaviors. Consistent with this, we observed significant changes in TNF- α and IL-6 levels 4 hours after exposure of microglia to A β , but not 3, 6, or 9 days after exposure to A β (data not shown).

Discussion

The present study found that AD microglia in vitro migrate toward A β aggregates, attempt to phagocytose the aggregates, and increase their secretion of TNF- α and IL-6 in the process. Opsonization of A β aggregates with anti-A β antibody 6E10 significantly enhanced these processes. By contrast, the common NSAID indomethacin had no material or statistical effect on microglial migration or phagocytosis, but significantly inhibited the increased IL-6 secretion observed with anti-A β opsonization.

The shifts in microglial distributions relative to A β spots over time are most parsimoniously explained by chemotactic responses to A β . Proliferation of microglia more proximal to A β aggregates was not observed and, in fact, BrdU reactivity, a common marker for cell proliferation, was negligible at all distances from the aggregates. Chemokinesis, enhanced but undirected movement of cells, also did not appear to explain the results, since microglial migration exhibited the gradient characteristics of chemotaxis, with progressive increases in the density of microglia at distances more proximal to A β aggregates. In addition, microglia are now well established to express receptors that can mediate chemotactic behaviors and that appear to have A β as a ligand. These include the macrophage scavenger receptor [16-18], the receptor for advanced glycation endproducts (RAGE) [15], the formyl peptide receptor [19], and others [20,21]. RAGE, in particular, has been shown to help mediate microglial migration to A β spots in an in vitro paradigm similar to that used here, and this migration could be inhibited by anti-RAGE Fab fragments [15].

AD microglia in vitro also exhibited behaviors consistent with phagocytosis of A β aggregates. Entering the paradigm, the microglia showed little or no A β immunoreactivity. After 12 days incubation with A β spots, the microglia were highly immunoreactive for A β and the spots decreased in size. A β spots without microglia remained essentially intact over the same time period. Previous ultrastructural and other studies [3,22,23] have also identified A β filaments within microglia in the vicinity of A β deposits in AD cortex. Although it remains possible that the intracellular A β within microglia in the AD brain may have been produced by the cells [24] rather than phagocytosed from an extracellular deposit, this is clearly not the process observed in the present in vitro studies. We conclude, therefore, that AD microglia in vitro do phagocytose aggregated A β deposits. Given the





Figure 3
Microglial distributions after 9 days incubation with Aβ spots. A) Treatment with 2 μg/ml anti-Aβ antibody plus (yellow) or minus (green) 1 μg/ml indomethacin (INDO). B) Treatment with vehicle control (red) or 10 μg/ml anti-Aβ antibody plus (yellow) or minus (green) 1 μg/ml indomethacin. C) Representative phase contrast image (4 × objective) of microglia and an Aβ spot when treated with vehicle only. D) Representative phase contrast image (4 × objective) of microglia and an Aβ spot when treated with 10 μg/ml anti-Aβ antibody plus 1 μg/ml indomethacin.

**Figure 4**

Evidence for phagocytosis of A β by AD microglia in vitro under the various experimental conditions. A) Twelve days after plating AD microglia with A β spots, diminution of the spots was visually apparent and microglia concurrently had become immunoreactive for A β even under vehicle control conditions, as shown here (anti-A β antibody 4G8 immunocytochemistry). B) In the absence of microglia, the A β spots remained visibly intact (phase contrast). C) Likewise, prior to exposure to A β spots the microglia exhibited little or no immunoreactivity for A β (anti-A β antibody 4G8 immunocytochemistry). D) Summary data illustrating the effects of indomethacin and 6E10 opsonization on A β spot thickness. Microglia in this model system carpet the top of A β spots (c.f., Fig. 2C) and therefore appear to clear the A β from the top down, resulting in progressive thinning of the spot, as measured here. With prolonged exposure, cracks and holes in the spot appear, as shown in Fig. 4A.




Figure 5
Effects on microglial TNF- α (A) and IL-6 (B) secretion into the medium in the presence or absence of A β , as well as after pretreatment of A β with 10 μ g/ml anti-A β antibody 6E10. Opsonization with 6E10 significantly enhanced ($P < 0.05$) (*) TNF- α and IL-6 levels compared to A β alone. IL-6 experiments also measured the effect of 1 μ g/ml indomethacin on 6E10 exacerbation of cytokine secretion. Indomethacin significantly reduced this effect ($P < 0.05$) (#).

experimental accessibility of the model, it will be of interest in future to evaluate the molecular fate of phagocytosed A β in cultured AD microglia.

Exposure to aggregated A β also induced significant increases in TNF- α and IL-6 secretion, confirming our previous experiments [12] and those of others [25-27] with TNF- α , IL-6, and a broad range of chemokines, cytokines, and inflammatory toxins such as reactive oxygen/nitrogen species. Pathways for enhancing TNF- α and IL-6 secretion have been demonstrated, including NF- κ B and C/EBP transcriptional mechanisms, both of which are enhanced

in pathologically-vulnerable regions of the AD brain [28,29].

Opsonization of A β spots with anti-A β antibody 6E10 significantly enhanced microglial migration to the spots, phagocytosis of the spots, and cytokine secretion. Similar effects of opsonization on microglial migration and phagocytosis have also been reported using anti-A β antibodies and an in vitro preparation in which cultured rodent microglia were seeded onto postmortem AD cortex sections laden with A β deposits [6]. Soluble Fab fragments containing the Fc region ligand for Fc receptor binding inhibited A β removal in this paradigm. These effects are consistent with the classic mechanisms of antibody opsonization of immune targets by antibodies specific to epitopes on the target. Scavenger cells that express receptors to the Fc region of the antibodies are then directed to or become focused at the site where the antibody-bound target resides. Fc receptor activation, in addition, activates scavenger cells, promoting attack and phagocytosis. Recently, scientists at Elan Pharmaceuticals have attempted to harness these mechanisms to enhance A β clearance, using immunization with A β to drive production of anti-A β antibodies for subsequent A β opsonization [6,30]. Although there is controversy about the exact site of action of the antibodies (e.g., brain versus peripheral circulation) [6,30,31], this approach does clearly result in significant and sometimes dramatic reductions of A β burden in transgenic mouse models [6], as well as the in vitro model tested here, and may also have been effective in human patients receiving A β immunization [30].

Unfortunately, however, inflammatory responses are often a two-edged sword. Fc receptor binding is known to enhance the activation and pro-inflammatory secretory responses of scavenger cells that bear Fc receptors, and microglia do express these receptors [6,32]. The increased TNF- α and IL-6 secretion observed in the present experiments after opsonization of A β aggregates with a specific anti-A β antibody, 6E10, is therefore not unexpected. On activation, microglial cells are, in fact, well established to secrete a wide range of inflammatory mediators that could not only cause damage to neurons and neurites locally, but also, if sufficiently activated, provide signalling to peripheral immune cells to provoke a more generalized and severe response such as that reported in several A β -immunized patients who experienced lethal adverse reactions [30].

The vast majority of NSAIDs in use today are based on the principle of cyclooxygenase inhibition, and cyclooxygenase inhibition, in turn, is well established to downregulate a wide range of acute phase reactants. Interestingly, however, mechanisms for chemotaxis to and phagocytosis of

an inflammatory target are not necessarily cyclooxygenase dependent. In a survey, for example, of the first 100 publications retrieved from PubMed using the search phrase "indomethacin AND chemotaxis", the majority of studies found no effect of indomethacin on chemotaxis, and some of the papers actually reported enhanced chemotaxis after indomethacin exposure. Such findings have been suggested to explain why physicians commonly prescribe NSAIDs to control fever and other secondary inflammatory responses without being unduly concerned about hampering immune-mediated removal of the fever-inducing agent. Similarly, in the present experiments indomethacin had no material or statistically significant effect on microglial chemotaxis to or phagocytosis of A β aggregates, but did significantly inhibit the exacerbated IL-6 response under opsonized conditions. Although it is never certain that in vitro results will fully apply to the in vivo state, these results suggest that indomethacin or an NSAID like it might be a useful adjunct to A β immunization strategies.

Competing interests

JR is a co-inventor on an issued United States patent covering use of nonsteroidal anti-inflammatory drugs as a treatment for Alzheimer's disease. All other authors declare that they have no competing interests.

Authors' contributions

JR conceived and designed the experiments, performed all data analysis, and wrote the manuscript. RS supervised and took part in all experiments. CJK performed the chemotaxis, phagocytosis, and cytokine experiments. DM, BL, and AG prepared cultures and performed histochemistry and immunocytochemistry.

Acknowledgements

This research was directly supported by NIA AGO7367. Institutional support for Alzheimer's research was provided by the Arizona Alzheimer's Disease Core Center (P30 AG019610) (NIA) and the Arizona Alzheimer's Consortium (State of Arizona). We thank Kyle Mueller, Gita Seetharaman, and Leyla Descheny for technical assistance, and Dr. Emily Lue and Dr. Douglas Walker for technical advice.

References

- Luber-Narod J, Rogers J: **Immune system associated antigens expressed by cells of the human central nervous system.** *Neurosci Lett* 1988, **94**:17-22.
- Rogers J, Luber-Narod J, Styren SD, Civin WH: **Expression of immune system-associated antigens by cells of the human central nervous system: relationship to the pathology of Alzheimer's disease.** *Neurobiol Aging* 1988, **9**:339-349.
- Terry RD, Wisniewski HM: **Structural and chemical changes of the aged human brain.** *Psychopharmacol Bull* 1975, **11**:46-63.
- Weldon DT, Rogers SD, Ghilardi JR, Finke MP, Cleary JP, O'Hare E, Esler WP, Maggio JE, Mantyh PW: **Fibrillar beta-amyloid induces microglial phagocytosis, expression of inducible nitric oxide synthase, and loss of a select population of neurons in the rat CNS in vivo.** *J Neurosci* 1998, **18**:2161-2173.
- Ard MD, Cole GM, Wei J, Mehrle AP, Fratkin JD: **Scavenging of Alzheimer's amyloid beta-protein by microglia in culture.** *J Neurosci Res* 1996, **43**:190-202.
- Bard F, Cannon C, Barbour R, Burke RL, Games D, Grajeda H, Guido T, Hu K, Huang J, Johnson-Wood K, Khan K, Kholodenko D, Lee M, Lieberburg I, Motter R, Nguyen M, Soriano F, Vasquez N, Weiss K, Welch B, Seubert P, Schenk D, Yednock T: **Peripherally administered antibodies against amyloid beta-peptide enter the central nervous system and reduce pathology in a mouse model of Alzheimer disease.** *Nat Med* 2000, **6**:916-919.
- Bard F, Barbour R, Cannon C, Carretto R, Fox M, Games D, Guido T, Hoenow K, Hu K, Johnson-Wood K, Khan K, Kholodenko D, Lee C, Lee M, Motter R, Nguyen M, Reed A, Schenk D, Tang P, Vasquez N, Seubert P, Yednock T: **Epitope and isotype specificities of antibodies to beta - amyloid peptide for protection against Alzheimer's disease-like neuropathology.** *Proc Natl Acad Sci U S A* 2003, **100**:2023-2028.
- Schenk DB, Yednock T: **The role of microglia in Alzheimer's disease: friend or foe?** *Neurobiol Aging* 2002, **23**:677-679.
- McGeer EG, McGeer PL: **Clinically tested drugs for Alzheimer's disease.** *Expert Opin Investig Drugs* 2003, **12**:1143-1151.
- McGeer EG, McGeer PL: **Brain inflammation in Alzheimer disease and the therapeutic implications.** *Curr Pharm Des* 1999, **5**:821-836.
- Lue LF, Brachova L, Walker DG, Rogers J: **Characterization of glial cultures from rapid autopsies of Alzheimer's and control patients.** *Neurobiol Aging* 1996, **17**:421-429.
- Lue LF, Rydel R, Brigham EF, Yang LB, Hampel H, Murphy GM Jr, Brachova L, Yan SD, Walker DG, Shen Y, Rogers J: **Inflammatory repertoire of Alzheimer's disease and nondemented elderly microglia in vitro.** *Glia* 2001, **35**:72-79.
- Bruguerolle B, Barbeau G, Belanger PM, Labrecque G: **Pharmacokinetics of a sustained-release product of indomethacin in the elderly.** *Gerontology* 1986, **32**:277-285.
- Lue LF, Brachova L, Civin WH, Rogers J: **Inflammation, A beta deposition, and neurofibrillary tangle formation as correlates of Alzheimer's disease neurodegeneration.** *J Neuropathol Exp Neurol* 1996, **55**:1083-1088.
- Lue LF, Walker DG, Brachova L, Beach TG, Rogers J, Schmidt AM, Stern DM, Yan SD: **Involvement of microglial receptor for advanced glycation endproducts (RAGE) in Alzheimer's disease: identification of a cellular activation mechanism.** *Exp Neurol* 2001, **171**:29-45.
- Honda M, Akiyama H, Yamada Y, Kondo H, Kawabe Y, Takeya M, Takahashi K, Suzuki H, Doi T, Sakamoto A, Ookawara S, Mato M, Gough PJ, Greaves DR, Gordon S, Kodama T, Matsushita M: **Immunohistochemical evidence for a macrophage scavenger receptor in Mato cells and reactive microglia of ischemia and Alzheimer's disease.** *Biochem Biophys Res Commun* 1998, **245**:734-740.
- Christie RH, Freeman M, Hyman BT: **Expression of the macrophage scavenger receptor, a multifunctional lipoprotein receptor, in microglia associated with senile plaques in Alzheimer's disease.** *Am J Pathol* 1996, **148**:399-403.
- El Khoury J, Hickman SE, Thomas CA, Cao L, Silverstein SC, Loike JD: **Scavenger receptor-mediated adhesion of microglia to beta-amyloid fibrils.** *Nature* 1996, **382**:716-719.
- Lorton D, Schaller J, Lala A, DeNardin E: **Chemotactic-like receptors and Abeta peptide induced responses in Alzheimer's disease.** *Neurobiol Aging* 2000, **21**:463-473.
- Koenigsnecht J, Landreth G: **Microglial phagocytosis of fibrillar beta-amyloid through a beta1 integrin-dependent mechanism.** *J Neurosci* 2004, **24**:9838-9846.
- Bourin M, Ripoll N, Dailly E: **Nicotinic receptors and Alzheimer's disease.** *Curr Med Res Opin* 2003, **19**:169-177.
- Cole GM, Beech W, Frautschy SA, Sigel J, Glasgow C, Ard MD: **Lipoprotein effects on Abeta accumulation and degradation by microglia in vitro.** *J Neurosci Res* 1999, **57**:504-520.
- Wisniewski HM, Terry RD: **Morphology of the aging brain, human and animal.** *Prog Brain Res* 1973, **40**:167-186.
- Banati RB, Gehrman J, Czech C, Monning U, Jones LL, Konig G, Beyreuther K, Kreutzberg GW: **Early and rapid de novo synthesis of Alzheimer beta A4-amyloid precursor protein (APP) in activated microglia.** *Glia* 1993, **9**:199-210.
- Ito S, Sawada M, Haneda M, Fujii S, Oh-Hashi K, Kiuchi K, Takahashi M, Isobe K: **Amyloid-beta peptides induce cell proliferation and macrophage colony-stimulating factor expression via the P13-kinase/Akt pathway in cultured Ra2 microglial cells.** *FEBS Lett* 2005, **579**:1995-2000.

26. Floden AM, Li S, Combs CK: **Beta-amyloid-stimulated microglia induce neuron death via synergistic stimulation of tumor necrosis factor alpha and NMDA receptors.** *J Neurosci* 2005, **25**:2566-2575.
27. Mrak RE, Griffin WS: **Glia and their cytokines in progression of neurodegeneration.** *Neurobiol Aging* 2005, **26**:349-354.
28. Li R, Strohmeyer R, Liang Z, Lue LF, Rogers J: **CCAAT/enhancer binding protein delta (C/EBP delta) expression and elevation in Alzheimer's disease.** *Neurobiol Aging* 2004, **25**:991-999.
29. Bales KR, Du Y, Holtzman D, Cordell B, Paul SM: **Neuroinflammation and Alzheimer's disease: critical roles for cytokine/Abeta-induced glial activation, NF-kappaB, and apolipoprotein E.** *Neurobiol Aging* 2000, **21**:427-430.
30. Masliah E, Hansen L, Bard F, Lee C, Seubert P, Games D, Kirby L, Schenk D: **A β vaccination effects on plaque pathology in the absence of encephalitis in Alzheimer's disease.** *Neurology* 2005, **64**:129-131.
31. DeMattos RB, Bales KR, Cummins DJ, Dodart JC, Paul SM, Holtzman DM: **Peripheral anti-Abeta antibody alters CNS and plasma Abeta clearance and decreases brain Abeta burden in a mouse model of Alzheimer's disease.** *Proc Natl Acad Sci U S A* 2001, **98**:8850-8855.
32. Song X, Tanaka S, Cox D, Lee SC: **Fc γ receptor signaling in primary human microglia: differential roles of PI-3K and Ras/ERK MAPK pathways in phagocytosis and chemokine induction.** *J Leukoc Biol* 2004, **75**:1147-1155.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

