

Cavernous Hemangioma of the Small Bowel: A Case Report and Literature Review

Ceren Durer¹, Seren Durer¹, Mohamad Sharbatji¹, Isin Y. Comba², Ilan Aharoni³, Umair Majeed⁴

1. Internal Medicine, Florida Hospital-Orlando, Orlando, USA 2. Internal Medicine, UCF Internal Medicine Residency Program, ORLANDO, USA 3. Gastroenterology, Florida Hospital, Orlando, USA 4. Internal Medicine Residency, Florida Hospital-Orlando, Placentia, USA

✉ **Corresponding author:** Mohamad Sharbatji, mohamad.sharbatji.md@flhosp.org
Disclosures can be found in Additional Information at the end of the article

Abstract

Hemangiomas of the small intestine are rare and very difficult to diagnose preoperatively. Clinical presentations may include occult or massive gastrointestinal (GI) bleeding, obstruction, intussusception, and perforation. We report a 66-year-old Caucasian male patient with severe anemia secondary to occult GI bleeding from a cavernous hemangioma in the jejunum. A double balloon enteroscopy following capsule endoscopy was performed to obtain biopsy samples, which established the final diagnosis.

Categories: Internal Medicine, Gastroenterology

Keywords: cavernous hemangioma, capsule endoscopy, double balloon enteroscopy, chronic anemia

Introduction

Hemangioma of the small bowel is an uncommon condition, accounting for 5 to 10% of all benign lesions of the small bowel [1]. It usually presents in young people with no sex predilection. Intestinal hemangiomas may cause occult or massive gastrointestinal bleeding, obstruction, intussusception, and small bowel perforation [2]. According to the size of affected vessels, hemangiomas are histologically classified into cavernous, capillary or mixed-type, with the cavernous type being the most common. In the gastrointestinal tract, the most commonly involved site is the jejunum. We report a case of cavernous hemangioma of the jejunum presenting with severe anemia. No remarkable findings were detected in upper and lower endoscopies. Correspondingly, the patient underwent wireless capsule endoscopy and double-balloon enteroscopy, which demonstrated a 25-mm submucosal mass with mild superficial ulceration/erosions of the surface mucosa. The pathology demonstrated that the lesion was consistent with cavernous hemangioma without evidence of malignancy.

Case Presentation

A 66-year-old male presented to the emergency room due to worsening leg pain. His past medical history was significant for peripheral artery disease and iron deficiency anemia. Initial laboratory tests revealed an unexpectedly low hemoglobin level of 5.4 g/dl. He received three units of packed red blood cells and subsequently the hemoglobin level increased to 6.9 g/dl. Our gastroenterology department was consulted for evaluation of occult gastrointestinal bleeding. There was no hematochezia, melena, hematemesis, fatigue, or abdominal pain. The patient had been taking oral iron supplementation for the last five years for iron deficiency anemia. Previous upper and lower endoscopies were negative. On physical examination, the patient had pale conjunctivae. The abdomen was noted to be soft and non-tender. No masses,

Received 07/27/2018

Review began 08/02/2018

Review ended 08/04/2018

Published 08/06/2018

© Copyright 2018

Durer et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 3.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

How to cite this article

Durer C, Durer S, Sharbatji M, et al. (August 06, 2018) Cavernous Hemangioma of the Small Bowel: A Case Report and Literature Review. Cureus 10(8): e3113. DOI 10.7759/cureus.3113

organomegaly or vascular bruits were detectable. The vital signs were stable, and the laboratory investigations were as follows: a hemoglobin (hb) level of 6.9 g/dL, a mean corpuscular volume of 73.5, a hematocrit level of 22.7% with normal white blood cell and platelet counts. The analyses for iron-deficiency anemia showed ferritin levels of 6 ng/mL, serum iron levels of 25 µg/dL, total iron-binding capacity of 535 µg/dL, and transferrin saturation of 5%. Upper and lower endoscopy showed no active bleeding or suspicious lesions. A small bowel capsule endoscopy was performed, which revealed a suspicious lesion over the jejunum with evidence of fresh blood (Figure 1).




FIGURE 1: Capsule endoscopy

For further investigation, double balloon enteroscopy was performed, which revealed a proximal jejunal soft submucosal mass (25 mm) with mild superficial ulceration/erosions of the surface mucosa. Biopsy samples were taken from the mass and the patient was referred for further management and surgical evaluation. The final pathology results revealed a cavernous hemangioma without evidence of malignancy.

Discussion

Cavernous hemangioma is one of the congenital benign vascular lesions and is not a true tumor. It can be solitary, multiple, or associated with various syndromes such as blue rubber bleb nevus syndrome, Klippel-Trenaunay-Weber syndrome, and Maffucci syndrome [3].

Cavernous hemangioma is a soft, compressible, bluish purple lesion and consists of blood-filled spaces (cavern) between dilated vessels within the mucosa and submucosa. Its size may range from a few millimeters to several centimeters. Hemangioma, telangiectasia, angiodysplasia, and phlebectasia are typical forms of small bowel vascular lesions.

Preoperative diagnosis is extremely difficult because small hemangiomas are rarely demonstrable with traditional techniques such as upper and lower endoscopies. However, several imaging modalities including wireless capsule endoscopy, double balloon enteroscopy, multiphase computed tomography enterography (CTE), magnetic resonance enterography (MRE) are currently available options to investigate small bowel lesions [4]. Small bowel capsule endoscopy is a noninvasive imaging test and can be recommended when the source of bleeding remains unidentified after upper and lower endoscopy. On the other hand, double balloon enteroscopy is an invasive and highly sensitive diagnostic tool providing both therapeutic and diagnostic interventions. In the current case, the patient had severe chronic anemia without evidence of gross GI bleeding. Small bowel capsule endoscopy demonstrated suspicious findings in the jejunum and double-balloon enteroscopy helped diagnose the bluish submucosal mass in the small bowel.

A comprehensive search of electronic databases (PubMed, Embase, and the Web of Science databases) was performed to identify studies published from January 1, 2000 until June 30, 2018 using MeSH terms and keywords such as ‘cavernous hemangioma’, ‘wireless capsule endoscopy’, ‘double balloon enteroscopy’, ‘anemia’ and ‘lower gastrointestinal bleeding’. Reference lists were screened to identify additional relevant studies. A standardized form was used for data extraction. There have been about 15 cases of uncomplicated cavernous hemangioma of small intestine diagnosed preoperatively [1, 5-18]. Patient information is summarized in Table 1. Most patients were male with age ranging from 9 to 74 years. The jejunum was the most common site of small intestinal hemangiomas, and the initial clinical presentation was anemia in the majority of the cases. The diameter of the lesions varied from 0.8 cm to 50 cm. Most studies showed suspicious findings on capsule endoscopy and the lesions were resected surgically or endoscopically [1, 5-7, 9-14, 16-18]. Ejtehadi et al. demonstrated a hemangioma on scintigraphy and CT enterography while no positive finding was found on capsule endoscopy [15]; another reported case was diagnosed with contrast-enhanced CT and MRI preoperatively [8]. Cavernous hemangioma can also present with other clinical symptoms and complications. Abdul Aziz et al. reported a six-year-old patient with abdominal pain and distention who developed subacute intestinal obstruction secondary to cavernous hemangioma (15 cm) after blunt trauma to the abdomen [2]. Open surgical excision of the hemangioma along with affected small bowel was performed. In a guideline by the American Gastroenterological Association (AGA) in 2007 about the algorithms for the diagnosis and management of obscure gastrointestinal bleeding, capsule endoscopy is considered as the first-line procedure for the initial examination of small bowel after normal upper and lower endoscopies [19]. When positive findings are obtained on capsule endoscopy, double balloon enteroscopy or push enteroscopy is recommended for further management when there is no evidence of obstruction [19]. The American College of Gastroenterology (ACG) clinical guideline in 2015 also recommends the use of capsule endoscopy followed by deep enteroscopy based on findings on capsule endoscopy [20]. In the presented case, capsule endoscopy and balloon enteroscopy were performed according to these guidelines.

Author, Year	Age (Y)	Sex	Presentation	Preoperative diagnosis study	Hemangioma size (cm)	Hemangioma location	Treatment
Kazama et al. 2000 [8]	64	M	Anemia	Enteroclysis, contrast CT MRI	3.0 and 0.5	Jejunum	Surgery

Magnano et al. 2005 [1]	13	M	Fatigue, weakness anemia	Capsule endoscopy	2.0	Ileum	Laparotomy
Quentin et al. 2007 [9]	32	F	Hematochezia	Capsule endoscopy	2.0	Ileum	Laparotomy
Willert et al. 2008 [10]	19	M	Anemia	Capsule endoscopy+ balloon enteroscopy	0.8, 0.8 and 1.4	Jejunum, ileocecal valve	Endoscopic treatment
Pinho et al. 2008 [5]	9	F	Fatigue, dizziness, Anemia, melena	Capsule endoscopy	2.5	Ileum	Segmental resection
Chen et al. 2009 [7]	23	M	Anemia	Capsule endoscopy	3.0	Ileum	Laparoscopic treatment
Elias et al. 2010 [14]	39	M	Anemia	Capsule endoscopy+ push enteroscopy	NA	Jejunum	Laparotomy
Huber et al. 2012 [6]	23	M	Melena, anemia	Capsule endoscopy+ balloon enteroscopy	3.0	Jejunum	Laparoscopic operation
Pera et al. 2012 [18]	16	M	Anemia, palpitation, fatigue	Capsule endoscopy+ balloon enteroscopy	4.3	Jejunum	Laparoscopic operation
Ersoy et al. 2013 [13]	50	F	Hematemesis, melena	Capsule endoscopy+ balloon enteroscopy	NA	Jejunum	Segmental resection
Fernandes et al. 2014 [12]	56	F	hematochezia syncope, anemia	Capsule endoscopy+ CT enterography	14	Ileum	Laparotomy
Bae et al. 2015 [16]	13	M	Nausea, dizziness, anemia	Capsule endoscopy	5.2	Jejunum	Excision of the mass*
Peng et al. 2016 [11]	47	M	Weakness, melena	Capsule endoscopy+ Contrast-enhanced CT	50	Ileum	Laparotomy
Akazawa et al. 2016 [17]	56	F	Melena	Capsule endoscopy+ balloon enteroscopy	1.3	Jejunum	Laparoscopic operation
Ejtehadi et al. 2017 [15]	40	M	Fatigue, palpitation	Scintigraphy+ CT enterography	2.6	Jejunum	Laparoscopic operation

TABLE 1: Reports on small bowel cavernous hemangioma diagnosed preoperatively between 2000 and 2018

Conclusions

Cavernous hemangioma of the small bowel is a benign vascular lesion that can cause chronic GI bleeding presenting as chronic severe anemia. Clinical suspicion of bleeding originating from the small bowel is essential in the diagnosis of these lesions as they warrant a comprehensive workup. Based on the literature and guideline review, we recommend that capsule endoscopy followed by therapeutic double balloon enteroscopy be considered in those patients.

Additional Information

Disclosures

Human subjects: Consent was obtained by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following:

Payment/services info: All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

- Magnano A, Privitera A, Calogero G, Nanfito L, Basile G, Sanfilippo G: Solitary hemangioma of the small intestine: an unusual cause of bleeding diagnosed at capsule endoscopy. *J Pediatr Surg.* 2005, 40:25-27. [10.1016/j.jpedsurg.2005.06.014](https://doi.org/10.1016/j.jpedsurg.2005.06.014)
- Abdul Aziz DA, Khandasamy Y, Tamba RP, Zaki FM: Bleeding small bowel cavernous haemangioma following blunt trauma to the abdomen presenting as subacute intestinal obstruction in a child. *BMJ Case Rep.* 2011, 2011, [10.1136/bcr.08.2011.4672](https://doi.org/10.1136/bcr.08.2011.4672)
- Wilson CL, Song LM, Chua H, Ferrara M, Devine RM, Dozois RR, Nehra V: Bleeding from cavernous angiomatosis of the rectum in Klippel-Trenaunay syndrome: report of three cases and literature review. *Am J Gastroenterol.* 2001, 96:2783-2788. [10.1111/j.1572-0241.2001.04110.x](https://doi.org/10.1111/j.1572-0241.2001.04110.x)
- Watari I, Oka S, Tanaka S, Nakano M, Aoyama T, Yoshida S, Chayama K: Is occult obscure gastrointestinal bleeding a definite indication for capsule endoscopy? A retrospective analysis of diagnostic yield in patients with occult versus overt bleeding. *Gastroenterol Res Pract.* 2013, 2013:915463. [10.1155/2013/915463](https://doi.org/10.1155/2013/915463)
- Pinho R, Rodrigues A, Proenca L, et al.: Solitary hemangioma of the small bowel disclosed by wireless capsule endoscopy. *Gastroenterol Clin Biol.* 2008, 32:15-18. [10.1016/j.gcb.2007.11.004](https://doi.org/10.1016/j.gcb.2007.11.004)
- Huber A, Abdel Samie A, Kychenko D, Theilmann L: A rare cause of recurrent iron-deficiency anemia: cavernous hemangioma of the small intestine. *J Gastrointest Liver Dis.* 2012, 21:343.
- Chen CH, Jones J, McGowan P: Profound iron deficiency anemia caused by a small-intestinal cavernous hemangioma. *Gastrointest Endosc.* 2009, 69:1392-1393. [10.1016/j.gie.2009.01.049](https://doi.org/10.1016/j.gie.2009.01.049)
- Kazama T, Kurihara Y, Tani I, Takahara T, Nakajima Y, Atari E: MR appearance of the small intestinal cavernous hemangioma. *J Comput Assist Tomogr.* 2000, 24:655-656.
- Quentin V, Lermite E, Lebigot J, Marannes MZ, Arnaud JP, Boyer J: Small bowel cavernous hemangioma: wireless capsule endoscopy diagnosis of a surgical case. *Gastrointest Endosc.* 2007, 65:550-552. [10.1016/j.gie.2006.12.024](https://doi.org/10.1016/j.gie.2006.12.024)
- Willert RP, Chong AK: Multiple cavernous hemangiomas with iron deficiency anemia successfully treated with double-balloon enteroscopy. *Gastrointest Endosc.* 2008, 67:765-767. [10.1016/j.gie.2007.07.044](https://doi.org/10.1016/j.gie.2007.07.044)
- Peng C, Chen H, Li W, Xu R, Zhuang W: A rare cause of recurrent gastrointestinal bleeding: giant diffuse and cavernous intestinal mesentery hemangioma in an adult. *Dig Dis Sci.* 2016, 61:3363-3365. [10.1007/s10620-016-4259-2](https://doi.org/10.1007/s10620-016-4259-2)
- Fernandes D, Dionisio I, Neves S, Duarte P: Cavernous hemangioma of small bowel: a rare

- cause of digestive hemorrhage. *Rev Esp Enferm Dig.* 2014, 106:214-215.
13. Ersoy O, Akin E, Demirezer A, Koseoglu H, Balci S, Kiyak G: Cavernous haemangioma of small intestine mimicking gastrointestinal stromal tumour. *Arab J Gastroenterol.* 2013, 14:139-140. [10.1016/j.ajg.2013.08.008](https://doi.org/10.1016/j.ajg.2013.08.008)
 14. Elias G, Toubia N: Hemangioma of the small intestine presenting with recurrent overt, obscure gastrointestinal bleeding. *Clin Gastroenterol Hepatol.* 2010, 8:18-11. [10.1016/j.cgh.2009.03.036](https://doi.org/10.1016/j.cgh.2009.03.036)
 15. Ejtehad F, Fattahi MR, Safaei A, Safarpour AR, Bananzadeh A: Practical lessons from the small bowel bleeding lesions: a case report on small bowel cavernous hemangioma. *Iran J Med Sci.* 2017, 42:108-110.
 16. Bae SJ, Hwang G, Kang HS, Song HJ, Chang WY, Maeng YH, Kang KS: Single cavernous hemangioma of the small bowel diagnosed by using capsule endoscopy in a child with chronic iron-deficiency anemia. *Clin Endosc.* 2015, 48:340-344. [10.5946/ce.2015.48.4.340](https://doi.org/10.5946/ce.2015.48.4.340)
 17. Akazawa Y, Hiramatsu K, Nosaka T, et al.: Preoperative diagnosis of cavernous hemangioma presenting with melena using wireless capsule endoscopy of the small intestine. *Endosc Int Open.* 2016, 4:249-251. [10.1055/s-0041-111321](https://doi.org/10.1055/s-0041-111321)
 18. Pera M, Marquez L, Dedeu JM, Sanchez J, Garcia M, Ramon JM, Puigvehi M: Solitary cavernous hemangioma of the small intestine as the cause of long-standing iron deficiency anemia. *J Gastrointest Surg.* 2012, 16:2288-2290. [10.1007/s11605-012-1991-6](https://doi.org/10.1007/s11605-012-1991-6)
 19. Raju GS, Gerson L, Das A, Lewis B: American Gastroenterological Association (AGA) Institute technical review on obscure gastrointestinal bleeding. *Gastroenterology.* 2007, 133:1697-1717. [10.1053/j.gastro.2007.06.007](https://doi.org/10.1053/j.gastro.2007.06.007)
 20. Gerson LB, Fidler JL, Cave DR, Leighton JA: ACG clinical guideline: diagnosis and management of small bowel bleeding. *Am J Gastroenterol.* 2015, 110:1265-1287. [10.1038/ajg.2015.246](https://doi.org/10.1038/ajg.2015.246)