

Rat T Cell Responses to Superantigens. II. Allelic Differences in V β 8.2 and V β 8.5 β Chains Determine Responsiveness to Staphylococcal Enterotoxin B and Mouse Mammary Tumor Virus-encoded Products

By Daniel P. Gold,* Charles D. Surh,† Karen S. Sellins,§
Kim Schroder,* Jonathan Sprent,† and Darcy B. Wilson*

From the *San Diego Regional Cancer Center, San Diego, California 92121; the †Department of Immunology, Scripps Clinic and Research Foundation, La Jolla, California 92037; and the §Barbara Davis Center for Childhood Diabetes, Denver, Colorado 80262

Summary

The previous paper in this series demonstrates that rat T cells developing de novo in the presence of mouse mammary tumor virus (Mtv) antigens in rat \rightarrow severe combined immunodeficiency (SCID) mouse xenochimeras display a distinct pattern of V β -restricted deletion; this deletion pattern is remarkably similar to that occurring during thymic development of mouse T cells in Mtv⁺ strains. In addition, T cells developing in the absence of Mtv antigens in these rat \rightarrow mouse xenochimeras are tolerant of host antigens, but show strong primary proliferative responses in cultures stimulated with Mtv-7⁺ (Mls^a) mouse cells; like the mouse, these rat T cell responses are dominated by V β 6 and V β 8 T cells. Here, we continue analysis of rat T cell responses to superantigens; we show that T cells from Lewis and Fischer 344 rats expressing V β 8.2 display an important all-or-nothing difference in their responses to Mtv-7 superantigens. This all-or-none strain difference in the response to Mtv-7 applies also to the response by V β 8.2 and V β 8.5 T cells to the soluble superantigen staphylococcal enterotoxin B. Because these two rat strains express different alleles of these two V β 8 family members, this finding identifies additional, hitherto unreported residues of the T cell receptor β chain important in T cell responses to superantigens.

In the preceding paper (1), evidence is presented that rat T cells closely resemble mouse T cells in possessing strong V β -restricted reactivity for mouse mammary tumor virus (Mtv)¹ superantigens. This evidence came from studies on rat-derived T cells developing in immunodeficient C.B-17 SCID mice reconstituted with fetal liver (FL) cells from Lewis (LEW) rat embryos. Provided that rat FL cells are coinjected with mature mouse B cells, the rat T cells arising in the chimeras display an extensive pattern of V β -restricted clonal deletion in response to various endogenous Mtv antigens of the host. Interestingly, this pattern of V β deletion is remarkably similar to the deletion pattern seen in normal mice. In addition to responding to mouse Mtv antigens in terms of clonal deletion, T cells from rat \rightarrow mouse chimeras mount strong proliferative responses to non-self Mtv antigens, i.e., to the Mtv-7 (Mls^a) antigens of the BALB.D2 and DBA/2 strains.

In this paper we show that proliferative responses of T

cells from LEW FL \rightarrow SCID mouse chimeras to Mtv-7 antigens involve V β 6 and V β 8.2 T cells. Surprisingly, however, this does not apply to chimeras prepared with FL cells from the Fischer 344 (F344) strain. Here the Mtv-7 response involves V β 6 cells but not V β 8.2 T cells. The ability of LEW but not F344 V β 8.2 T cells to respond to mouse Mtv-7 antigens also applies to responses to a soluble superantigen, staphylococcal enterotoxin B (SEB). Genomic sequences of V β 8.2 (and also V β 8.5) indicate that LEW and F344 express different alleles of these two V β 8 family members (Gold, D.P., S. Wiley, and D.B. Wilson, unpublished data). These findings identify residues distant from the defined superantigen combining site (2–5) that have not previously been appreciated to be involved in responses to superantigens.

Materials and Methods

Animals. Female rats of the LEW, F344, and ACI strains were purchased from Harlan Sprague Dawley, Inc. (Indianapolis, IN). Female DA rats were purchased from Bantin and Kingman, Inc. (Fremont, CA). Rats were 3–4 mo of age when used as donors for cell cultures. LEW and F344 \rightarrow C.B-17 SCID chimeras were prepared as described (1).

¹ Abbreviations used in this paper: F344, Fischer 344; FL, fetal liver; LEW, Lewis; Mtv, mammary tumor virus; SEB, staphylococcal enterotoxin B.

Antibodies. mAbs specific for rat T cell markers TCR- α/β (R73 [6]), V β 8.2 (R78) (7), V β 8.5 (B73) (7), CD4 (OX-38) (8), and activated CD4⁺ T cells (OX-40) (9) were obtained from Pharmingen (San Diego, CA). Biotin-modified rat F(ab')₂ anti-mouse IgG (H+L) was purchased from Accurate Chemical & Scientific Corp. (Westbury, NY).

Con A and rSEB Cultures. Lymph node cells (1.5×10^6 /ml) were cultured in RPMI 1640 (6 ml) supplemented with FCS (8% [vol/vol]), 2-ME (5×10^{-5} M), glutamine (2 mM), penicillin/streptomycin (100 μ g/ml), indomethacin (1 μ M; Sigma Chemical Co., St. Louis, MO), and either Con A (3 μ g/ml; Calbiochem, San Diego, CA) or recombinant SEB (rSEB; 2 μ g/ml). The rSEB was generously provided by Dr. John Kappler (National Jewish Hospital, Denver, CO) (10). After 3 d in culture, cells were expanded further in complete medium supplemented with supernatant (10%) from rat spleen cells activated with Con A for 48 h, as a source of IL-2, and methyl α -D-mannopyranoside (50 mM; Sigma Chemical Co.). T cell blasts were harvested 48 h later.

MLR Cultures. MLR cultures were established as described (1). Activated T cell blasts from the MLR cultures were purified by centrifugation through a stepwise Percoll gradient for 30 min at 1,400 g. Blast cells were harvested from the top of the 1.07-density Percoll layer.

Flow Cytometric Analysis. T cell blasts from Con A- and rSEB-stimulated cultures were labeled with the indicated anti-rat TCR mAbs on ice for 30 min, washed to remove unbound antibody, incubated for 20 min on ice with biotin-modified rat F(ab')₂ anti-mouse IgG (H+L), washed again, and then incubated for 20 min with PE-streptavidin (Biomed, Foster City, CA). Labeled cells were analyzed on a FACScan[®] flow cytometer (Becton Dickinson & Co., Sunnyvale, CA). T cell blasts recovered from MLR cultures were doubly stained with the indicated anti-TCR mAbs and anti-CD4, or with OX-38 and OX-40 mAbs as described earlier (1).

Determination of TCR V β Usage. Relative V β utilization among populations of T cells recovered from cultures was determined by a semiquantitative PCR assay (1).

Results

Strain Differences in the V β 8.2 T Cell Response to Mtv-7. In the accompanying paper (1), we report that T cells arising in LEW FL \rightarrow C.B-17 SCID mouse chimeras are tolerant of host-type BALB/c (and C.B-17) stimulators in vitro but give strong proliferative responses to DBA/2 and BALB.D2. Since BALB/c and BALB.D2 are congenic strains that differ

Table 1. PCR Analysis of TCR V β Usage by T Cells from LEW and F344 Rat \rightarrow SCID Mouse Xenochimeras in Response to Mtv-7

V β	LEW \rightarrow SCID			F344 \rightarrow SCID		
	None	BALB.D2	DBA/2	None	BALB.D2	DBA/2
1	1	1	1	2	6	1
2	2	4	7	6	10	2
3	1	0	1	3	1	0
4	7	1	2	3	1	0
5	0	2	1	1	1	0
6	27	51	45	16	61	85
7	0	1	1	5	2	2
8.2	2	10	11	14	3	4
8.5	3	2	3	4	2	2
8.6	3	8	3	4	2	0
9	5	2	2	5	1	1
10	12	3	3	6	2	0
11	1	0	0	4	1	0
12	0	0	0	0	0	0
13	0	1	0	0	0	0
14	13	3	3	3	2	1
15	10	4	4	4	1	0
16	0	0	1	2	0	0
17	2	0	2	3	0	0
18	1	1	3	2	0	0
19	5	1	2	8	1	0
20	3	1	2	4	2	0

Unstimulated rat T cells or T cell blasts stimulated in culture with mouse spleen cells for 5 d were assessed for TCR V β utilization by PCR. Blasts were purified by Percoll gradient. Values expressed are percent of total V β s.

only at the *Mtv-7* locus, the proliferative MLC response of chimera rat T cells to BALB.D2 (and DBA/2) is presumed to be directed to *Mtv-7* antigens. As shown in Table 1, culturing T cells from LEW FL → SCID chimeras with *Mtv-7*⁺ BALB.D2 or DBA/2 stimulators generates T cell blasts that, measured by PCR, are selectively enriched for two *Vβ*s, i.e., *Vβ6* and *Vβ8.2*; the enrichment for these cells correlates with the selective elimination of *Vβ6* and *Vβ8.2* cells in *Mtv-7*⁺ chimeras, i.e., in LEW FL → SCID chimeras coinjected with DBA/2 B cell blasts that express *Mtv-7* antigens (1). Other *Vβ* members, for example, *Vβ2*, sometimes showed modest increases, but this was variable from animal to animal.

Different results apply to F344 FL → SCID chimeras. T cells from these animals give strong MLR responses to DBA/2 and BALB.D2. In this situation, however, the blast response does not involve *Vβ8.2* T cells and is largely restricted to *Vβ6* cells.

These data apply to detection of *Vβ* by PCR. *Vβ* analysis by flow cytometry (limited to *Vβ8.2*, with *Vβ10* as a control) is shown in Fig. 1 *a*. In this experiment, T cells from LEW → SCID chimeras and LEW FL → SCID chimeras injected at birth with BALB/c B cell blasts (1) were cultured in vitro with BALB.D2 or DBA/2 stimulators. With these two *Mtv-7*⁺ stimulators, a high proportion of the OX-40⁺ CD4⁺ blasts in the cultures were *Vβ8.2*⁺ (25–35%). With *Mtv-7*⁻ H-2-different B10.BR stimulators, by contrast, there was no enrichment for *Vβ8.2*⁺ T cells. Control *Vβ10*⁺ T cells were rare with BALB.D2 and DBA/2 stimulators, but were somewhat increased with B10.BR stimulators. These data refer to LEW chimeras. With F344 chimeras (F344 → SCID), stimulation with BALB.D2 and DBA/2 cells also generated large numbers of OX-40⁺ blast cells, but here very few of these blasts were *Vβ8.2*⁺.

The above data indicate, both by PCR and flow cytometry, that the response of *Vβ8.2*⁺ T cells to *Mtv-7* applies only to LEW and not to F344 T cells. As shown in Fig. 1 *b*, this strain difference in the response of rat *Vβ8.2*⁺ cells to *Mtv-7* also applies to primary MLR by normal (nonchimeric) T cells. Thus, in marked contrast to LEW T cells, culturing normal F344 T cells with BALB.D2 stimulators fails to cause a selective enrichment in *Vβ8.2* T cells (relative to BALB/c stimulators).

The data presented below indicate that the inability of F344 *Vβ8*-bearing T cells to respond to *Mtv-7* also applies to responses to SEB.

Strain Differences in the *Vβ8.2* and *Vβ8.5* T Cell Response to rSEB. In a previous study we demonstrated by PCR analysis that T cells from LEW rats preferentially use *Vβ7*, 8.2, 8.5, and 14 in the proliferative response to rSEB in culture (11). During the course of those studies, we noted that whole T cell populations from the DA strain responded well to rSEB, but T cells of the *Vβ8.2* subset did not. With the recent description of two mAbs specific for *Vβ8.2* and *Vβ8.5* T cells of the rat (7), we were able to confirm and extend these findings.

Fig. 2 presents the results of a flow cytometric analysis




Figure 1. Percent rat *Vβ8.2*⁺ and *Vβ10*⁺ cells among activated (OX-40⁺) CD4 T blasts derived by FACS[®] analysis of MLC stimulated with third-party (*B10.BR*) or *Mtv-7*⁺ (*BALB.D2* and *DBA/2*) spleen cells. Responders are LN cells from (a) LEW FL → SCID (with or without BALB/c B cell blasts) and F344 FL → SCID xenochimeras, or (b) normal LEW and F344 donors. *Vβ8.2* T cells from LEW and LEW → SCID chimera donors respond to *Mtv-7*, but *Vβ8.2* T cells from F344 and F344 → SCID donors do not.

showing the percent TCR- α/β ⁺, *Vβ8.2*⁺, and *Vβ8.5*⁺ T cell blasts present after stimulation in culture with Con A or with rSEB. Blasts were gated on the basis of forward and orthogonal light scatter criteria. These results indicate that *Vβ8.2* and *Vβ8.5* T cells from LEW donors, normally present at the level of 5–7% of the peripheral T cell pool, make a substantial contribution to the blast population stimulated with rSEB, but T cells of these two subsets from F344 donors do not.

Table 2 presents a summary of comparisons of *Vβ8.2* and *Vβ8.5* T cell responses to SEB involved several different rat strains. The results show that the frequency of LEW *Vβ8.2* and *Vβ8.5* T cells responsive to rSEB is two- to threefold higher than the response to Con A, while for the other strains, it is about the same or less. The *Vβ8.2* and *Vβ8.5* T cell responses to Con A were similar among the three strains and approximated the level found among normal, nonstimulated T cells.

Predicted Amino Acid Sequences of the TCR *Vβ8.2* and *Vβ8.5* Chains. Genomic clones encoding for both the *Vβ8.2* and




Figure 2. Flow cytometric analysis of rat TCR- α/β , V β 8.2, and V β 8.5 expression among T cell blasts from cultures stimulated with Con A or rSEB. V β 8.2 and V β 8.5 T cells from LEW donors respond to rSEB, but F344 T cells bearing these V β 8 family members do not.

V β 8.5 gene segments were sequenced from liver DNA of F344, DA, and ACI rats (Gold, D. P., S. Wiley, and D. B. Wilson, manuscript in preparation); the predicted amino acid sequences are shown in Table 3. Sequences for the V β 8.2 gene segments from F344, DA, and ACI animals are identical for these three rat strains, as are the sequences for the V β 8.5 chain. Both gene segments encode functional genes, but of particular interest, both display allelic differences from the published predicted sequences for these β chain segments in LEW animals (12, 13).

Discussion

This study explores responses by V β 8.2⁺ and V β 8.5⁺ T cells from different rat strains to two different superantigens.

The general finding here is that T cells from some rat strains bearing these TCR β chain members respond to these superantigens while T cells from other strains do not. This all-or-nothing difference in response to these two superantigens correlates with polymorphic sequence differences in the V β 8.2 and V β 8.5 TCR β chains known to exist in these strains.

When stimulated with the Mtv-7 superantigen, Mls^a, V β 8.2 T cells from normal LEW donors respond in xenogeneic rat-mouse MLC cultures to Mtv-7⁺ BALB.D2 stimulator cells, but contribute almost no cells to the population responding to H-2-identical Mtv-7⁻ stimulators. V β 8.2 T cells from F344 donors give no response to Mtv-7 (Fig. 1 b). Similarly, LEW and F344 T cells made tolerant of mouse H-2^d MHC antigens as a consequence of thymic development in rat \rightarrow SCID mouse chimeras show a major differ-

Table 2. Flow Cytometric Analysis of Strain-specific Differences in Responses to rSEB by V β 8.2 and V β 8.5 T Cells

Strain	Stimulus	Percent of blast cells			
		Control	R73 TCR- α/β	R78 V β 8.2	B73 V β 8.5
LEW	Con A	2.6	99	6.3	9.1
	rSEB	1.2	83	21	24
F344	Con A	1.6	98	7.7	7.5
	rSEB	2.4	86	3.0	8.2
DA	Con A	2.9	99	9.0	7.3
	rSEB	0.9	83	3.3	3.8
ACI	Con A	1.1	98	7.7	6.2
	rSEB	4.9	70	3.4	5.3

Lymph node cells from the indicated strains were incubated with either Con A or rSEB for 3 d. Surviving cells were isolated and further expanded for 2 d in medium supplemented with Con A supernatant. Results are given as percent of blast cells expressing TCR- α/β , V β 8.2, or V β 8.5. Blasts were gated on the basis of forward and orthogonal light scatter criteria.

Table 3. Predicted Amino Acid Sequences for the LEW and F344 TCR V β 8.2 and V β 8.5 Chains

Rat	V β	1	10	20	30	40	50	60	70	80	90
LEW	8.2*	EAAVTQSPRNKVTLLKGGKVTLSCKQNNHNNMYWYRQDMGHGLRLIHYSYDVNSTEKGDVDPNGYKYSRPSQGDFFLTLLESASPSQTSVYFCASS									
F344	8.2†S.....V.....D.....									
LEW	8.5§	EAAVTQSPRNKVTVTGKNVTFNCHQTDNHNMYWYRQDMGHGLRLIHYSYSGSGSFENGDIPEGYKYSRPNQENIFFLTLLESASPSQTSVYFCASS									
F344	8.5‡K.....S.....									

* Smith et al. (12).
 † Gold, D.P., S. Wiley, and D.B. Wilson (manuscript in preparation).
 ‡ Hashim et al. (13). This sequence corresponds to V β 8.3 published by Smith et al. (12).

ence in the response of V β 8.2 T cells to Mtv-7 antigens; LEW T cells contribute ~30% of blasts to the responding population, while F344 T cells contribute <5% (Fig. 1 a).

For the SEB superantigen, T cells from LEW, F344, DA, and ACI rats all show strong proliferative responses; exposure to this superantigen for a few days stimulates the T cell population to the extent that it consists mainly (70–100%) of blast cells, the magnitude of the response being similar to that occurring in cultures stimulated with Con A. But, unlike the response to Con A, there appears to be a highly selective, all-or-none difference with respect to whether T cells expressing the V β 8.2 and V β 8.5 gene segments contribute to the SEB-responsive population. LEW T cells expressing these two β chain members selectively contribute 20–25% to the SEB-stimulated blast populations, a proportion three- to four-fold greater than that contributed to Con A-stimulated blasts, whereas T cells bearing these two TCR β chain members from F344, DA, and ACI rats show no response to SEB (Fig. 2 and Table 2).

What element(s) might account for the all-or-none response of V β 8.2 and V β 8.5 T cells to these superantigens? The simplest and most direct interpretation is that these response differences stem from differences in the primary amino acid sequence of the V β 8.2 and V β 8.5 alleles (Table 3). Three amino acids distinguish SEB and Mtv-7 responder LEW V β 8.2 chains from nonresponder F344 V β 8.2 chains: these occur at positions 9 (R vs. S), 14 (L vs. V), and 62 (N vs. D). To our knowledge this is the first reported example of allelism being associated with responsiveness, or lack of it, to these two very different kinds of superantigens. Similarly, two amino acids at positions 15 (T vs. K) and 83 (P vs. S) are associated with responsiveness or nonresponsiveness of V β 8.5⁺ T cells to SEB.

Other explanations for the difference in responsiveness of V β 8.2 and V β 8.5 T cells from LEW and F344 donors to superantigens can be considered. For example, might some gene(s) other than those encoding TCR β chain alleles be involved? This seems unlikely since whole populations of T cells from F344 donors are fully responsive to both SEB and Mtv-7; whether or not a given T cell responds depends on a particular allele of a given TCR β chain.

Several structure–function studies have helped to define the precise residues of the TCR β chain thought to be responsible for recognition in responses to superantigens (2–5). These involve sequence comparison of allelic TCR β chain gene products where one allows for T cell responses to a given superantigen and another does not (2, 4). Sequence comparisons have also been conducted on closely related members of a given β chain family (5), and the functional consequences of sequence differences have been confirmed by direct mutagenesis studies (3). Results of these studies focus on the importance of residues surrounding positions 19–26 on the B β -strand and 68–76 on the loop between the D and E β -strands, using the Chothia alignment where position 23 is cys (14), in the response of murine V β 8.2⁺ T cells to Mtv-7. Certain residues in these two regions of the β chain are also associated with V β 17 responses to some unknown

Table 4. Summary of TCR V β Polymorphisms in Rats and Mice Associated with Responsiveness to SEB and Mtv-7

V β	Animal	SAg response	Amino acid position										Reference
			9	10	14	15	24	30-31	51-53	62	72-73	83	
8.2	LEW	SEB/Mls +	R	N	L	K	K		GSG	N	GD	P	12
	F344	-	S	N	V	K	K		GSG	D	GD	P	This paper
	ER34	Mls +		S	V		D	DY	GAD	D	KE	P	2
	C57Bl/6	-		N	V		N	NN	GAG	D	EN	P	2
8.5	LEW	SEB +	R		V	T	H			E	EN	P	13
	F344	-	R		V	K	H			E	EN	S	This paper
8.1	C57Bl/6	Mls +			V		H		VAD	D	EN	L	2

ligand associated with I-E molecules in the mouse (4), and with the response of human V β 13.1 and 13.2 T cells to the *Staphylococcus aureus* toxin SEC2 (5). These residues, although situated at opposite ends of the primary sequence, have been predicted from computer modeling studies, based on the crystal structure of Ig, to lie juxtaposed on the surface of the TCR molecule in the β -pleated sheets that contribute to the framework of the β chain; this site is accessible to the aqueous interface of the molecule and is located well away from the CDR surface believed to interact with conventional peptide antigen/MHC complexes (14, 15). The region around residues 68-76 has been suggested to be a fourth hypervariable region in Ig molecules (HV4), but it is not known to be involved in antigen binding (14).

The present studies suggest two conclusions. First, it seems clear that other regions of the TCR β variable chain segment are also involved in superantigen responses. The sequence differences associated with responsiveness by rat V β 8.2 T cells to SEB and Mtv-7 reside at positions 9, 14, and 62, and differences associated with V β 8.5 responsiveness to SEB reside at positions 15 and 83. These two regions are thought to lie well away from the site responsible for recognition of MHC/peptide antigen (14, 15), but also they appear to be quite removed from the HV4 region reported to be important in responses to Mls superantigens (3).

Second, it is tempting to consider the possibility that a particular amino acid or combination of them, in a given

TCR β chain, may be directly involved in the TCR binding interaction to a superantigen that leads to activation. Evidence for such a notion would be provided by the finding that a particular amino acid residue at a given position is always associated with responsiveness (or the lack of it) to a superantigen. But, such predictions based on residues at a given position appear not to be possible. Table 4 is a summary that compares the amino acid residues present at various positions of allelic TCR β chains in mice and rats where one allelic product is responsive to a superantigen and the other is not. For example, in the rat changes from R to S, L to V, and N to D at positions 9, 14, and 62 of the V β 8.2 chain are associated with an all-or-none difference in the response to rSEB and Mtv-7. Yet these residues of the responder V β 8.2 β chain are present in the nonresponder V β 8.5 chain; likewise V¹⁴ and D⁶² present in the nonresponder V β 8.2 chain are also present in the responder V β 8.5 chain. This occurs also in mice. Others have focused on the KE to EN substitution at position 72-73 that distinguishes Mls responder and nonresponder V β 8.2 alleles (2), yet EN is expressed on the mouse V β 8.1 chain that is responsive to Mtv-7. This situation suggests that the role of a given amino acid residue in the response to a superantigen might be an indirect one, for example, causing a conformational difference at a distant site, and that predictions of superantigen responsiveness based on amino acid residues at a given position on the TCR β chain can be considered only in the context of the total sequence.

We appreciate the technical help of Deanna Mueller and Sandra Wiley.

This study was supported by United States Public Health grants AI-21487, AI-22519, AI-31451, AI-33011, CA-25803, and CA-38355. C.D. Surh is a Special Fellow of the Leukemia Society of America.

Address correspondence to Daniel P. Gold or Darcy B. Wilson, San Diego Regional Cancer Center, 3099 Science Park Road, San Diego, CA 92121.

Received for publication 2 July 1993.

References

1. Surh, C.D., D.P. Gold, S. Wiley, D.B. Wilson, and J. Sprent. 1993. Rat T cell responses to superantigens. I. V β -restricted clonal deletion of rat T cells differentiating in rat \rightarrow mouse chimeras. *J. Exp. Med.* 179:27.
2. Pullen, A.M., W. Potts, E.K. Wakeland, J. Kappler, and P. Marrack. 1990. Surprisingly uneven distribution of the T cell receptor V β repertoire in wild mice. *J. Exp. Med.* 171:49.
3. Pullen, A.M., T. Wade, P. Marrack, and J.W. Kappler. 1990. Identification of the region of T cell receptor β chain that interacts with the self-superantigen Mls-1^a. *Cell.* 61:1365.
4. Cazenave, P.-A., P.N. Marche, E. Jouvin-Marche, D. Voegtle, F. Bonhomme, A. Bandeira, and A. Coutinho. 1990. V β 17 gene polymorphism in wild-derived mouse strains: amino acid substitutions in the V β 17 region greatly alter T cell receptor specificity. *Cell.* 63:717.
5. Choi, Y., A. Herman, D. DiGiusto, T. Wade, P. Marrack, and J. Kappler. 1990. Residues of the variable region of the T-cell-receptor β -chain that interact with *S. aureus* toxin superantigens. *Nature (Lond.)*. 346:471.
6. Hünig, T., H.-J. Wallny, J.K. Hartley, A. Lawetzky, and G. Tiefenthaler. 1989. A monoclonal antibody to a constant determinant of the rat T cell antigen receptor that induces T cell activation. Differential reactivity with subsets of immature and mature T lymphocytes. *J. Exp. Med.* 169:73.
7. Torres-Nagel, N.E., D.P. Gold, and T. Hünig. 1993. Identification of rat Tcr V β 8.2, 8.5 and 10 gene products by monoclonal antibodies. *Immunogenetics.* 37:305.
8. Jefferies, W.A., J.R. Green, and A.F. Williams. 1985. Authentic T helper CD4 (W3/25) antigen on rat peritoneal macrophages. *J. Exp. Med.* 162:117.
9. Patterson, D.J., W.A. Jefferies, J.R. Green, M.R. Brandon, P. Cortes, M. Puklavec, and A. Williams. 1987. Antigens of activated rat T lymphocytes including a molecule of 50,000Mr detected only on CD4 positive T blasts. *Mol. Immunol.* 24:1281.
10. Kappler, J.W., A. Herman, J. Clements, and P. Marrack. 1992. Mutations defining functional regions of the superantigen staphylococcal enterotoxin B. *J. Exp. Med.* 175:387.
11. Sellins, K.S., D. Bellgrau, and D.P. Gold. 1992. Specificity of rat T cell receptor V β chain usage in proliferative responses to staphylococcal enterotoxin B. *Eur. J. Immunol.* 22:1931.
12. Smith, L.R., D.H. Kono, and A.N. Theofilopoulos. 1991. Complexity and sequence identification of 24 rat V β genes. *J. Immunol.* 147:375.
13. Hashim, G., A.A. Vandenbark, D.P. Gold, T. Diamanduros, and H. Offner. 1991. T cell lines specific for an immunodominant epitope of human basic protein define an encephalitogenic determinant for experimental autoimmune encephalomyelitis-resistant LOU/M rats. *J. Immunol.* 146:515.
14. Chothia, C., D.R. Boswell, and A.M. Lesk. 1988. The outline structure of the T-cell $\alpha\beta$ receptor. *EMBO (Eur. Mol. Biol. Organ.) J.* 7:3745.
15. Davis, M.M., and P.J. Bjorkman. 1988. T-cell antigen receptor genes and T-cell recognition. *Nature (Lond.)*. 334:395.