


Client perspectives on creating supportive sexual health environments for people with persistent anxiety: a qualitative study

Stéphanie Black ¹, Sarah Watt,¹ Mark Gilbert ^{2,3}, Heather Nicole Pedersen,³ Aidan Ablona ³, Hsiu-Ju Chang,³ Travis Salway^{1,4}

To cite: Black S, Watt S, Gilbert M, *et al.* Client perspectives on creating supportive sexual health environments for people with persistent anxiety: a qualitative study. *BMJ Public Health* 2024;**2**:e000625. doi:10.1136/bmjph-2023-000625

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjph-2023-000625>).

Received 4 October 2023
Accepted 28 June 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. Published by BMJ.

¹Faculty of Health Sciences, Simon Fraser University, Burnaby, British Columbia, Canada

²School of Population and Public Health, The University of British Columbia, Vancouver, British Columbia, Canada

³Clinical Prevention Services, BC Centre for Disease Control, Vancouver, British Columbia, Canada

⁴The University of British Columbia Centre for Gender and Sexual Health Equity, Vancouver, British Columbia, Canada

Correspondence to
Dr Stéphanie Black;
stephanie_black@sfu.ca

ABSTRACT

Objectives We sought to explore sexual health service clients' perspectives, preferences and suggestions for how to better address or alleviate anxiety among clients of sexual health services (eg, sexually transmitted infection testing) regarding both online and in-person sexual health services among those who experience persistent anxiety.

Methods We conducted qualitative interviews with 27 sexual health service clients with persistent experiences of anxiety in British Columbia, Canada. Participants were recruited from respondents to a COVID-19-related sexual health survey who consented to follow up. Interviews were conducted via Zoom, recorded and transcribed. Transcripts were coded and analysed using thematic analysis by searching for themes in order to summarise the experiences and needs of participants.

Results The median participant age was 34. 16 participants identified as women, 10 as men and 1 each as non-binary and gender fluid. Participants described anxiety related to sexual health service access and experiences due to sexual health-related stigma and privacy concerns; provider judgement and lack of communication or information regarding test results. They suggested that routinely integrating discussions with providers about mental health in sexually transmitted and bloodborne infection (STBBI) testing appointments may help clients feel safer and could connect them to support. They highlighted the need for personal and genuine interactions with providers (eg, making appointments feel less perfunctory, asking clients how they are feeling about why they are there) and sufficient time with providers (eg, not feeling rushed through the appointment, time to discuss resources, testing and advice).

Conclusions Disease-specialised health services may not adequately address the multifaceted and inter-related mental health needs of people accessing services. In STBBI testing service settings, more personalised appointments, additional communication with providers and easier access to results can help improve both the service experiences of people experiencing persistent anxiety and their connections to appropriate mental health support.

INTRODUCTION

Existing research suggests that sexual health-care settings can unintentionally contribute

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Existing research suggests that sexual health testing settings may induce or exacerbate feelings of anxiety.

WHAT THIS STUDY ADDS

⇒ Little is known about sexual health clients' mental health experiences or needs. This study describes experiences of anxiety among clients and identifies opportunities for improved sexual and mental health service integration.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Findings will help to inform provider approaches to addressing anxiety within a sexual health testing setting.

to feelings of anxiety.^{1–4} Anxious feelings may stem from the possibility of a positive sexually transmitted and bloodborne infection (STBBI) result, the need for partner notification, anticipated provider judgement, embarrassment, stigma and privacy concerns.^{1–3 5 6} Previous research has demonstrated a high prevalence of general anxiety among STBBI testing clients (including in person and online),^{6 7} suggesting that acute experiences of anxiety may interact with more chronic experiences of anxiety.⁴ Further, in light of pervasive barriers to mental health services (eg, high costs, provider availability, long wait times, stigma and the logistics of accessing care^{7–9}), low-barrier service settings such as sexual health services may offer opportunities to strengthen connections to care.

It is not uncommon for people to access mental healthcare (eg, seeking care for depression, anxiety, other mental health conditions) in other low barrier, more accessible locations like primary care and sexual health clinics.^{4 9 10} Indeed, previous research highlights the potential benefits of greater service integration and accessibility.^{10 11} It

Table 1 Characteristics of participants in a qualitative study regarding anxiety among STBBI testing service clients (B.C., summer 2020)

	n	%
Age		
20–29	8	29.6
30–39	15	55.6
40–59	4	14.8
Gender*†		
Woman	16	59.3
Man	10	37.0
Non-binary	1	3.7
Genderfluid	1	3.7
Sexuality*†		
Heterosexual	12	44.4
Bisexual	6	22.2
Gay	6	22.2
Pansexual	1	3.7
‘Mostly heterosexual’	2	7.4
Ethnicity*†		
White	19	70.4
Chinese	3	11.1
Hispanic	1	3.7
Eastern European	1	3.7
Iranian	1	3.7
Latin	1	3.7
Mixed ancestry	1	3.7
Employment status*		
Employed full time	17	63.0
Employed part time	3	11.1
Self-employed	6	22.2
Student	4	14.8
Retired	1	3.7
Unemployed	1	3.7
Other	1	3.7
Type of STBBI service accessed*		
GetCheckedOnline		
Yes	22	81.5
No	3	11.1
Did not specify	2	7.4
In-person		
Yes	22	81.5
No	0	n/a
Did not specify	5	18.5
Total	27	

*Responses are not mutually exclusive; totals do not equal total sample size.

†Includes open text box responses.

STBBI, sexually transmitted and bloodborne infection.

is important to understand how sexual health settings can better address the needs of people experiencing anxiety—both by helping to mitigate anxiety in the context of sexual health service access and by facilitating connections to mental health supports. Creating supportive sexual health environments for addressing anxiety during STBBI testing encounters may be beneficial to both the client and the broader healthcare system.^{5 10}

However, little is known about how sexual health settings are experienced by people with chronic or persistent experiences of anxiety, nor about their perspectives on anxiety alleviation through improvements to service delivery and connections to mental health. Therefore, we conducted a study to explore sexual health service clients’ perspectives, preferences and suggestions for how to better address or alleviate feelings of anxiety regarding both online and in-person sexual health services among those who experience persistent anxiety. We conducted a qualitative descriptive study and used thematic analysis to analyse the data.^{12 13}

METHODS

Participants

We interviewed 27 participants from British Columbia (B.C.), Canada. They were recruited among respondents of an online COVID-19-related survey about sexual health, administered by the BC Centre for Disease Control to sexual health service clients in July–August 2020.¹⁴ A subset of participants consented to be contacted for interviews following the survey. Interviews took place October–December, 2020. In order to be eligible, prospective participants had to (1) be 16 years of age or older, (2) have accessed sexual health services in B.C. within the past 12 months and (3) have experienced anxiety (eg, nervousness, panic, inability to stop worrying) for 2 weeks or more within the past 12 months. Consent was obtained before proceeding with the interviews.

Data collection

Team members used a semistructured interview guide (online supplemental file 1) to inquire about participants’ experiences of anxiety and other mental health concerns, accessing mental and sexual health supports, and desires and expectations regarding mental and sexual health service integration. Topics included in the interview guide were (A) Participant Motivation; (B) Mental Health Experiences & Access to Services; (C) COVID-19 Experiences; (D) Sexual Health Service Experiences and (E) Interpretations of Mental Health and Substance Use Tools Within Sexual Health Services. We used a semistructured interview guide to allow for natural conversation flow, follow-up questions and dialogue between interviewer and participant. Interviews lasted between 45 and 90 min. They were conducted via Zoom and were recorded, anonymised and transcribed for analysis. A

follow-up survey was sent postinterview to collect demographic and mental health information.

Analysis

Interviews were conducted by SB, SW and patient partner RR. SB and SW are trained in qualitative methodologies and RR was trained by SW and SB to conduct interviews. Interviews were transcribed and accuracy checked by SB, SW and RR. All transcripts were double coded by SB, SW or RR. Team members met regularly to discuss the use of codes and to address any discrepancies. Descriptive coding using NVivo V.12, followed by pattern coding,¹¹ was carried out by two researchers (SB and SW). We inductively searched for higher-level themes in order to summarise the experiences and meanings of participants⁹ while staying close to the data. Themes were defined and iteratively redefined by both analysts to achieve internal consistency.

Patient partner involvement

Team member RR was engaged in the study based on their lived experience working in and accessing sexual health services. They helped develop the interview guide, conducted interviews, supported accuracy-checking of transcripts and cocreated the codebook for analysis.

RESULTS

Characteristics of sample

We conducted 27 interviews with STBBI testing clients (see [table 1](#)) from B.C. All participants self-reported that they had accessed online (n=22) and/or in-clinic (n=22) STBBI testing services within the past 12 months. Most participants accessed testing services routinely while some had only tested periodically or for a particular event.

Almost all participants (n=26) had previously accessed or were currently accessing mental health services (eg, counselling). Mental health experiences varied in duration and intensity. Participants described anxiety related to their sexual health in general, and anxiety when accessing STBBI testing services. Some participants described underlying mental health conditions (eg, generalised anxiety and depression) yet felt comfortable receiving STBBI tests while some reported anxiety specifically related to STBBI testing. Others experienced anxiety both directly related to STBBI testing and various underlying/chronic conditions.

The following sections detail the major themes identified: contributors to anxiety in STBBI testing settings; participant perspectives on integrating mental health supports in sexual health appointments and participant suggestions for 'anxiety-supportive' sexual health services.

Contributors to anxiety in sexual health settings

Regardless of whether they had a diagnosed mental health condition, many participants expressed anxiety related to STBBI testing. This was typically related to provider

judgement, privacy, stigma, the possibility of having an STBBI and communication regarding test results.

Provider judgement

Many participants felt anxious about provider judgement when attending in-person appointments. Some experienced it directly while others anticipated the potential for judgement. Participants felt that accessing testing through online services, without having to speak with a provider allowed them to avoid potentially uncomfortable interactions. One participant said:

[...] I don't need to speak to someone 'cause they're going to give me the same spiel every time. Sometimes there'll be judgments because I'm in an open relationship and we've made that decision to do that. So I don't have to explain all that. (Woman, mid twenties)

Another participant (man, late twenties) worried about the potential for providers to think less of him due to how frequently he sought testing.

Privacy, stigma and testing anxiety

Privacy related to testing concerned several participants. Some felt that clinic spaces did not afford adequate anonymity from others in and around the clinic. A desire for privacy when accessing sexual health services in particular stemmed from the stigmatised nature of sexual health, sexual practices, sexuality and STBBI testing. One participant explained:

[I]t's just like I really don't want to have one [an STBBI]. You know, I think part of just growing up in a world with all those stigmas about it is you're just taught that it's the worst thing in the world to get, and it's like, you know, cancer, but embarrassing. (Man, early twenties)

Similarly, another person identified shame around partner notification when diagnosed with an STBBI. They said: 'it implies that you're slutty or that you aren't safe, um...yeah, it's a very shameful thing to have to do' (Woman, late twenties).

Communication regarding STBBI test results

Many of the services participants accessed only followed up with clients if results were positive. This lack of communication from providers troubled several people who felt anxious waiting for results. One participant (man, early twenties), described the 'no news is good news' approach to test results as 'frustrating', and wished providers would call to deliver results either way. Participants using an online testing service appreciated having access to all results within the service portal. The desire for concrete information was important to participants. Being able to look up their test results through e-health systems offered people something to return to in moments of anxiety. One participant explained:

In those moments I can't just talk myself out of it or do something to distract myself. I have to go back, look at the

written thing that said ‘negative,’ and that’s the only thing that kind of tends to calm it [anxiety] down. (Woman, mid twenties)

Participant perspectives on integrating mental health supports in sexual health appointments

Participants expressed a range of perspectives when asked how and whether mental health supports should be integrated into STBBI testing services. Most participants were supportive of integration, provided that it was optional and sensitive to clients’ needs. They suggested that doing so may enhance the testing experience and help strengthen connections to mental health supports for those who were otherwise disconnected. They desired more connections to mental health services as well as more emotionally supportive interactions with providers.

Participants indicated that routinely asking about mental health in sexual health settings to strengthen pathways to mental health supports could enable a more tailored approach and help to normalise mental health discussions. Several participants emphasised that, given the stigmatised nature of mental health, care must be given to how and when this conversation is raised. They emphasised the importance of transparency from providers about why they are inquiring about mental health so that clients do not feel caught off guard:

I might just be like: ‘Well, why are you asking this, what are you picking up on?’ Whereas if it’s more front-loaded with a: ‘As part of this we’d like to do general screenings for sorts of mental health supports to see if there’s other things that could support you if we noticed you might benefit from them.’ I’d be more open to that idea[.] (Woman, mid twenties)

Conversely, some participants were sceptical of integrating sexual and mental health services. They indicated that the stigmatised nature of sexual health and mental health may cause clients to feel reluctant to discuss the latter. One participant explained that clients may already feel intimidated by the process of accessing STBBI testing services; to add a discussion of mental health may be overwhelming:

I think at a sexual health clinic when you know you’re going to be focused on the swabs and the blood tests and stuff it’s almost like enough to be kind of managing, and then the one-on-one questions about anxiety might just be, I don’t know, like you’d already be anxious because of what was about to happen, and it would just feel a little too personal when you’re already doing something personal. (Woman, late thirties)

For others, the two fields of practice were perceived as too unrelated. For these participants, discussions of mental health would be out of place or even intrusive in sexual health settings. One person likened receiving sexual and mental health services together to ‘getting your Christmas ham at the car dealership’ (woman, mid-thirties), highlighting their disparate nature.

Participant suggestions for ‘anxiety-supportive’ sexual health services

Participants offered a number of suggestions for supporting people with anxiety through their testing experiences (table 2). Suggestions highlighted opportunities to improve communication with providers, access to information and access to support. These are summarised in a tabular format.

Table 2 Participant recommendations for reducing anxiety among sexual health service clients

Recommendation	Example	Rationale
Easily accessible online results	GetCheckedOnline, Health Gateway and My Care Compass.	Clients can see and refer back to results at their convenience, thereby alleviating anxiety about results.
Providing reputable sources of educational information to clients about STBBIs	A list of educational information or websites about STBBIs (eg, Smart Sex Resource) either automatically generated by an online testing service, or given by providers during appointments.	Can help avoid ‘cyberchondria’, or excessive and often anxiety-exacerbating online information-seeking about the STBBI in question.
More communication opportunities with providers outside appointments	Notifying people about both positive and negative results, asking if they are interested in mental health resources.	Alleviate anxiety and provide more tailored advice or information.
More time with providers during appointments to speak about mental health	Addressing the potential mental health implications of a diagnosis and providing resources (eg, stigmatised diagnoses like herpes or HIV diagnosis)	Helps recognise that the provider cares about their well-being and wants to help them. Supports feelings of safety and comfort and encourages mental health discussions.
Live pop-up chat with a nurse in automated online testing services	Based on the answers generated to screening questions, a live pop-up chat with a nurse opens in a browser.	Since online testing service is automated, clients do not speak to providers. Having a chat option with a nurse could bridge the gap, allowing clients the privacy of generating their own requisitions, but also the option to talk to someone if they wish. Allows people access to have a conversation with a health professional, have their anxiety allayed and receive mental health or sexual health resources.
STBBIs, sexually transmitted and bloodborne infection.		

DISCUSSION

This study explored sexual health service clients' suggestions, perspectives and preferences for how to better address or alleviate anxiety concerns within STBBI testing service settings. Participants reported different levels of anxiety, a range of perspectives regarding the integration of mental health and STBBI testing services and provided suggestions for mitigating anxiety in these settings. Results highlight an important opportunity to integrate more 'anxiety-supportive' resources and methods in sexual health settings. Consistent with previous research, this suggests that improved resource and referral options are needed to better support sexual health service clients experiencing anxiety.¹⁵ For instance, a study about a sexual health information chat service described strategies and opportunities used by nurses to respond to persistent sexual health-related anxiety among clients (eg, providing anxiety management information, offering sexual health education and risk assessment).¹ Effective identification of and response to anxiety in STBBI testing service settings can help to ensure that clients are connected with appropriate resources and supports. Furthermore, providing reputable sources of STBBI information to clients can help them avoid excessive online information-seeking, which can be detrimental to one's mental health.^{1 4} Other research has also shown how low barrier (free, anonymous, inclusive of sexual/gender minorities, self-referred) sexual health settings may be particularly well placed to attend to such needs, given that mental health concerns often intersect and co-occur with sexual health concerns.^{4-6 13 14} It is also known from previous research that primary care and sexual health clinics can be advantageous sites of mental healthcare due to their accessibility, cost-effectiveness for both clients and the healthcare system and increased mental and physical health outcomes.^{15 16} Research has further demonstrated high levels of client desire for integrated or bundled mental health services alongside STBBI testing.¹⁰ Findings from the current study provide insight into clients' perspectives about how connections to mental health may be facilitated within a sexual health setting. Previous research has focused on sexual health provider perspectives about the integration of mental and sexual health services.^{12 14} Providers felt that integrating mental health workers within sexual health settings could help direct people to better resources and ensure clients receive timely care from trained mental health professionals. Our current study allowed us to hear clients' own perspectives on a different kind of mental health integration, focusing on anxiety reduction in sexual health settings. Practical implications for sexual health providers include: Ensuring STBBI test results are easily accessible online for clients to refer to at their convenience; providing reputable sources of educational STBBI information to clients, creating greater communication opportunities with providers to ensure clients feel heard and supported, and the creation of a live pop up chat service in online testing interventions for added support

with a professional. These practical implications are in line with previous research showing a need for anxiety-supportive sexual health environments, and a need for further education of sexual health clients.¹ For instance, one study showed that clients preferred one-on-one results notification to Short Message Service (SMS) notification because they had more time with a provider and cited anxiety about test results as a reason to refuse SMS notification of results.¹⁶

Findings from this study also contribute to the sparse literature about how clients with existing anxiety navigate health systems, and how best to improve mental and sexual health pathways for this population. Our findings are relevant to providers seeking to create environments that minimise barriers to sexual healthcare for people experiencing anxiety. More research is needed about the merits of integrating mental health and sexual health services, and about ways to implement such an integration. We also identified several limitations related to participant sampling and recruitment. First, in order to be eligible, prospective participants had to have experienced anxiety for 2 weeks or more within the preceding 12 months. However, recruitment occurred during the second wave of the COVID-19 pandemic. Given the profound mental health effects of the pandemic, some self-reported anxiety may have been situational rather than ongoing. Second, participants were recruited among respondents of a sexual health survey. The majority of respondents to this survey had experience accessing online sexual health services. This could have resulted in a sample with a high degree of digital health literacy, and disproportionately favourable views about, and comfort with, online testing service access. In addition, because we were specifically interested in experiences of anxiety within sexual health service settings, all participants had accessed sexual health services. Future studies should explore the needs of those whose experiences of anxiety have prevented them from accessing services. Existing research indicates that experiences accessing sexual health services may be shaped by intersecting experiences of stigma and discrimination (eg, related to gender, age).^{3 17} However, the homogeneity of the sample (eg, with regard to ethnic identity, gender), impeded our ability to explore these intersections. Future studies may build on existing research^{3 17} by exploring the unique experiences of anxiety in the context of sexual health services based on age, gender or other aspects of identity.

CONCLUSION

Disease-specialised health services may not adequately address the multifaceted and inter-related mental health needs of people accessing services. In STBBI testing service settings, more personalised appointments, additional communication with providers and easier access to results can help improve both the service experiences of people experiencing anxiety and their connections to appropriate mental health support. Providers

should consider ways to alleviate anxiety in sexual health settings, as outlined by the participants in this study. There is a need to review best practices to allow more time in appointments to help those experiencing anxiety feel comfortable with the providers and to ask any questions they have. Furthermore, ensuring that results are available promptly online in all cases (not just in the case of online testing services) would improve the experiences of all clients, including those with anxiety and sexual health anxiety specifically. Developing nurse text chat or phone lines could also be an outlet for those with anxiety to receive comforting and accurate information to alleviate their anxiety.

X Stéphanie Black @n/a

Acknowledgements The research team would like to thank Rowdy Reeves for his contribution to data collection.

Contributors The study was conceived by TS, SW and MG. Data were collected by SB, SW, TS and RR and analysed by SB and SW. The manuscript was originally designed by SB and SW with input from TS. Edits were executed by SB, SW, MG and TS. The paper was also reviewed and edited by AA, H-JC, HNP and MG to ensure accuracy, readability and applicability to practice. The final version was read and approved by all authors. TS is the guarantor and accepts full responsibility for the work.

Funding This study was funded by the Canadian Institutes of Health Research (437703) and Health Research BC, BC SUPPORT Unit (18904).

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and this study underwent a harmonised ethics review (# H20-00698) through the Research Ethics Boards at Simon Fraser University, the University of British Columbia and Fraser Health. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Stéphanie Black <http://orcid.org/0000-0003-2878-0208>

Mark Gilbert <http://orcid.org/0000-0001-5978-6843>

Aidan Ablona <http://orcid.org/0000-0003-1436-6276>

REFERENCES

- 1 Watt S, Salway T, Gómez-Ramírez O, *et al*. Rumination, risk, and response: a qualitative analysis of sexual health anxiety among online sexual health chat service users. *Sex Health* 2022;19:182–91.
- 2 Karamouzian M, Knight R, Davis WM, *et al*. Stigma associated with sexually transmissible infection testing in an online testing environment: examining the perspectives of youth in Vancouver, Canada. *Sex Health* 2018;15:46–53.
- 3 Shoveller JA, Knight R, Johnson J, *et al*. Not the SWAB! young men's experiences with STI testing. *Sociol Health Illn* 2010;32:57–73.
- 4 Tyrer P. Recent advances in the understanding and treatment of health anxiety. *Curr Psychiatry Rep* 2018;20:49.
- 5 Arkell J, Osborn DPJ, Ivens D, *et al*. Factors associated with anxiety in patients attending a sexually transmitted infection clinic: qualitative survey. *Int J STD AIDS* 2006;17:299–303.
- 6 Hottes TS, Farrell J, Bondyra M, *et al*. Internet-based HIV and sexually transmitted infection testing in British Columbia, Canada: opinions and expectations of prospective clients. *J Med Internet Res* 2012;14:e41.
- 7 Gaspar M, Marshall Z, Rodrigues R, *et al*. Mental health and structural harm: a qualitative study of sexual minority men's experiences of mental healthcare in Toronto, Canada. *Cult Health Sex* 2021;23:98–114.
- 8 Slaunwhite AK. The role of gender and income in predicting barriers to mental health care in Canada. *Community Ment Health J* 2015;51:621–7.
- 9 Martin D, Miller AP, Quesnel-Vallée A, *et al*. Canada's universal health-care system: achieving its potential. *The Lancet* 2018;391:1718–35.
- 10 Salway T, Ferlatte O, Shoveller J, *et al*. The need and desire for mental health and substance use-related services among clients of publicly funded sexually transmitted infection clinics in Vancouver, Canada. *J Public Health Manag Pract* 2019;25:E1–10.
- 11 Meiksin R, Melendez-Torres GJ, Falconer J, *et al*. eHealth interventions to address sexual health, substance use, and mental health among men who have sex with men systematic review and synthesis of process evaluations. *J Med Internet Res* 2021;23:e22477.
- 12 Sandelowski M. Whatever happened to qualitative description? *Nurs Health* 2000;23:334–40.
- 13 Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;3:77–101.
- 14 Gilbert M, Chang H-J, Ablona A, *et al*. Accessing needed sexual health services during the COVID-19 pandemic in British Columbia, Canada: a survey of sexual health service clients. *Sex Transm Infect* 2022;98:360–5.
- 15 Ng WZ, Ariffin K, Tay PKC, *et al*. Prevalence and risk factors for alcohol use disorders, substance use disorders, and depression anxiety and stress among users of sexual health services in Singapore: a cross-sectional survey study. *Sex Health* 2023;20:363–5.
- 16 Charron J, Troude P, de La Rochebrochard E, *et al*. Notification of STI test results by text Messaging: Why do patients refuse? cross-sectional study in a Parisian sexual health centre. *Int J STD AIDS* 2022;33:257–64.
- 17 Fallon D. They're gonNA think it now': narratives of shame in the sexual health experiences of young people. *Sociology* 2013;47:318–32.