

# Anti-inflammatory properties of combined aquatic extract of *Ferulago angulata* boiss with aerobic exercise on pro-inflammatory indices in obese males

Iman Zakavi<sup>1</sup>, Shila Nayebifar<sup>2</sup>, Elham Ghasemi<sup>3</sup>, Aliasghar Valipour<sup>1</sup>

<sup>1</sup>Department of Public Health, Abadan Faculty of Medical Sciences, Abadan University of Medical Sciences, Abadan, Iran, <sup>2</sup>Department of Sport Sciences, Faculty of Educational Sciences and Psychology, University of Sistan and Baluchestan, Zahedan, Iran, <sup>3</sup>Department of Sport Sciences, Faculty of Literature and Humanities, University of Zabol, Zabol, Iran

**Background:** The application of supplements, herbal extracts, and exercise training for treatment of diseases and reducing chronic inflammation has been increased progressively among people. Thus, the aim of this investigation was to study the combined aquatic extract of *Ferulago angulata* boiss with aerobic exercise on pro-inflammatory indices in obese males. **Materials and Methods:** In this semi-experimental study, forty young obese men (mean and standard deviation of age  $34.59 \pm 2.24$  years, body mass index (BMI)  $33.14 \pm 2.75$  kg/m<sup>2</sup>) were selected by purposive sampling and were randomly divided into four equal groups ( $n = 10$ ), training, training-supplementation, supplementation, and control. Participants in the supplementary groups received 50 mg/ml *F. angulata* extracts daily for 12 weeks. Aerobic training program included 12 weeks of training, 3 sessions/week, and each session was 20 min at 60%–70% of maximal oxygen consumption. Blood samples were taken from the participants 48 h before and after the intervention in fasting state. Data were analyzed using dependent *t*-test, one-way analysis of variance, and *post hoc* Tukey test at a significant level of  $P < 0.05$ . **Results:** After 12 weeks of exercise and supplementation, levels of interleukin (IL)-6 ( $P = 0.001$ ), IL-18 ( $P = 0.03$ ), IL-1  $\beta$  ( $P = 0.001$ ), tumor necrosis factor alpha ( $P = 0.001$ ), weight ( $P = 0.001$ ), BMI ( $P = 0.001$ ), body fat percent ( $P = 0.001$ ), and waist-hip ratio ( $P = 0.001$ ) decreased significantly and the mean changes of these indicators in training + supplementation group were significantly augmented as compared to the other three groups. **Conclusion:** It appears that aerobic training plus *F. angulata* extract consumption have better effect on improvement of serum inflammatory factors in obese young men.

**Key words:** *Ferulago angulata* extract, interleukins, obesity, tumor necrosis factor alpha, training

**How to cite this article:** Zakavi I, Nayebifar S, Ghasemi E, Valipour A. Anti-inflammatory properties of combined aquatic extract of *Ferulago angulata* boiss with aerobic exercise on pro-inflammatory indices in obese males. J Res Med Sci 2020;25:86.

## INTRODUCTION

Obesity is a multifactorial disease, and medical condition occurs by gene, lifestyle, and environmental factors interactions.<sup>[1]</sup> It is now evident that adipose tissue not only stores excess triglycerols, but also functions as an endocrine organ by releasing adipokines, which play important roles in the regulation of appetite, glucose and lipid metabolism, inflammation, and insulin resistance.<sup>[2]</sup> Research has shown that overproduction of pro-inflammatory cytokines such as interleukin (IL)-6,

IL-8, IL-1  $\beta$ , and tumor necrosis factor alpha (TNF- $\alpha$ ) released from adipose tissue plays an important role in the progression of obesity and serum concentrations of these inflammatory markers elevated in overweight and obese individuals, therefore, obesity especially visceral adiposity is now viewed as a low-grade inflammatory disease.<sup>[3]</sup> Research introduces adipose tissue as one of the main producers of inflammatory cytokines such as IL-6, IL-8, and IL-1  $\beta$ .<sup>[3,4]</sup> IL-6 is produced in a variety of tissues such as adipose tissue and skeletal muscle, and may have either pro- or anti-inflammatory

### Access this article online

Quick Response Code:



Website:  
[www.jmsjournal.net](http://www.jmsjournal.net)

DOI:  
10.4103/jrms.JRMS\_475\_19

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**For reprints contact:** WKHLRPMedknow\_reprints@wolterskluwer.com

**Address for correspondence:** Dr. Aliasghar Valipour, Department of Public Health, Abadan Faculty of Medical Sciences, Abadan University of Medical Sciences, Abadan, Iran. E-mail: aliasgharvalipour698@yahoo.com

**Submitted:** 27-Aug-2019; **Revised:** 02-Feb-2020; **Accepted:** 21-May-2020; **Published:** 30-Sep-2020

effects.<sup>[5]</sup> On the other hand, IL-18, TNF- $\alpha$ , and IL-1 $\beta$  are pro-inflammatory cytokines.<sup>[6-8]</sup> Physical activity and appropriate diet are among the curative and/or preventive measures for obesity-related complications.<sup>[9-12]</sup> According to previous studies, aerobic exercise is capable of promoting a transient anti-inflammatory state in healthy and overweight individuals, and it can aid in the control of chronic low-grade inflammation.<sup>[13,14]</sup>

However, no reduction was reported in the levels of inflammatory cytokines following regular exercise, in some studies.<sup>[15,16]</sup>

In addition, studies regarding the role of herb consumption in weight control and inflammation inhibition have attracted many researchers. *Ferulago angulata* is a plant from *Apiaceae* family, with thick stem, standing tall, and stripped. It has 35 species, while *F. angulata* has phenol and antioxidative properties.<sup>[17]</sup> It is proven that this plant prevents lipid peroxidation and act as anti-inflammatory.<sup>[17,18]</sup> In an animal study, Amirghofran *et al.* found that *F. angulata* extract consumption downregulates the production of pro-inflammatory cytokines, such as IL-1 $\beta$ , significantly.<sup>[19]</sup> Consequently, due to contrasting results about the effect of regular exercise on inflammatory profile in obese people, and the lack of evidence about the effect of *F. angulata* supplementation, the aim of this study was to investigate the effects of aerobic exercise along with water extract of *F. angulata* consumption on serum levels of IL-6, IL-18, IL-1 $\beta$ , and TNF- $\alpha$  in obese men.

## MATERIALS AND METHODS

### Study population

This is randomized placebo-controlled study including four groups containing a pretest, posttest design. The study was approved by the Ethic committee of the medical university of Abadan under the 93U-044 code and the Iranian Registry of Clinical Trials code is www.irct.ir; IRCT20160129026251N4. It is worth noting that the participants were fully informed of any risks and discomforts associated with the experiments before they provided informed written consent for participation. We included forty young, obese men according to MedCalc Statistical Software version 14.8.1 (MedCalc, Ostend, Belgium). This software was used to determine the standard sample size in each group. The sample size was determined based on the main and important variable of the study, "IL-6," based on 80% power, type 1 error is equal to 5%, and number of samples in each group of 10 people and a total of forty people were calculated. All the participants were randomly divided into four groups (training, training + supplementation, supplementation, and control) based on simple randomization procedures.

The inclusion criteria were body mass index (BMI) >30, age range (30–40 years), and gender (men). Exclusion criteria were not taking the researchers' advice, not attending the training sessions regularly, or having a history of special diseases, taking medicine and supplementations, and having a background of physical activity [Figure 1]. Diet monitored by a 24 h questionnaire in the first and last 3 days of the research period. Then, macro and micronutrients and calories in all groups calculated then, macro, micronutrients and calories calculated according to guide to calculate the Nutritional Value Regimes in Iran, book. Then, data were analyzed using analysis of variance (ANOVA) test. There was no significant difference between calories ( $P = 0.07$ ), micronutrients ( $P = 0.12$ ), and macronutrients ( $P = 0.23$ ) in the four studied groups, in pre- and post-test stages.

### Intervention

Before the beginning of the exercise program, the participants' weight was measured using a scale (Seca, Germany), following some seconds of staying motionless and their heights were assessed by meter with an accuracy of <0.5 cm. While participants were standing bare feet, their height and BMI were calculated by dividing body weight (kg) to the height<sup>2</sup> (m). Body fat percent was measured using a Caliper (SAEHAN-SH 5020, England), applying the Jackson and Pollock formula 3-site (chest, abdomen, and thigh) and after the 8–10 h of fasting.<sup>[20]</sup> Waist circumference was measured using a tape at halfway between lowest rib and the top of hipbone, roughly in line with belly button while breathe out normally. At the end of exercise program, all measurements were repeated under identical situations. Serum levels of IL-6, IL-18, IL-1 $\beta$ , and TNF- $\alpha$  were measured using highly sensitive Bender Medsystem kit (USA) and ELIZA method. Inter- and intra-coefficients of variation of the variable measurements were 5%–10%. Blood samples were collected before the first exercise session and 48 h after the last exercise session in fasting state between 8 and 10 am. Participants were asked not to have any strenuous physical activity 48-h before blood sampling. Five milliliters of blood was taken after 12 h fasting from brachial vein, and blood samples were collected in sterilized tubes and then incubated for 10 min at the room temperature. The samples were then centrifuged (at 2000 rpm for 10 min). Serum was separated and stored at -70°C for further analysis.

### Study protocol

Maximal oxygen uptake was measured using Bruce Test,<sup>[21]</sup> and the participants started to exercise after 72 h. Heart rates (HRs) of participants in 1 min were measured using a treadmill,<sup>[21]</sup> and the following formula was applied in order to calculate the maximal oxygen consumption ( $VO_{2max}$ ):<sup>[21]</sup>




Figure 1: Consort flow diagram of the study

$$VO_{2\max}: 14.76 - 1.379 (\text{time}) + 0.451 (\text{time})^2 - 0.12 (\text{time})^3$$

The target HR of participants were calculated using Karvonen formula:<sup>[21]</sup>

$$\text{Target HR: } (\text{maximum HR} - \text{resting HR}) \times (\text{intensity}) + \text{resting HR}$$

Maximal HR was calculated using the following formula: 220-age (years)

The aerobic training consisted of running on a treadmill at 60%–70% of  $VO_{2\max}$ . Each training session included a 5–10 min of warm-up and cool down. After 4 weeks, maximal HR was measured once more, and the training intensity (60%–70%  $VO_{2\max}$ ) was adjusted accordingly.<sup>[21]</sup>

### Supplement consumption

Leaves from *F. angulata* were grounded using an electric mill, and the powder was then bolted to separate the big particles. To make the *F. angulata* extract, 8 g of the powder was wrapped in a piece of two-layer tiffany cloth and then was put in a flask containing 100 mm of distilled, sterilized water. The container was then put in a shaker for 24 h at room temperature. The extract was then passed through a filter under the laboratory’s hood. The produced extract was then stored at 4°C before consumption.<sup>[22]</sup> According to the literature, the daily supplementation dose for each participant in the supplementation and the combination groups was 50 mg/kg body weight.<sup>[19]</sup> The researchers did not get any side effects of this supplement on human, up to now.<sup>[23]</sup>

Participants in the supplementation and the control groups did not take part in any exercise activities during this period and were only involved in their daily routine activities. At the end of intervention, the adherence to the supplementation was assessed 85% according to the self-reported information, while it was about 90% for adherence to exercise according to the sessions participants took part.

### Statistical analysis

Data were analyzed using Statistical Package for the Social Sciences version 19 software (SPSS Inc., Chicago, IL, USA). Data were expressed as mean ± standard deviation percentiles. After assessing the normality and nonnormality of data by Shapiro-Wilk test, differences between the baseline means of three groups in all variables were analyzed using oneway ANOVA. Within-group analysis assessed by paired *t*-test and between-group differences were examined using ANOVA; for pair-wise comparisons, Tukey *post hoc* test was used. A two-sided  $P \leq 0.05$  was considered statistically significant.

## RESULTS

Results from one-way ANOVA showed no significant statistical differences in pretest values of variables ( $P > 0.05$ ).

### Effects of 12 weeks of aerobic training and supplementation on inflammatory cytokines levels

Results from *t*-test shows that following 12 weeks of aerobic exercises and supplementation mean values for IL-6 ( $P = 0.003, P = 0.01, P = 0.001$ ), IL-18 ( $P = 0.03, P = 0.02,$

$P = 0.001$ ),  $\text{TNF-}\alpha$  ( $P = 0.03$ ,  $P = 0.02$ ,  $P = 0.001$ ), and  $\text{IL-1}\beta$  ( $P = 0.03$ ,  $P = 0.02$ ,  $P = 0.001$ ) decreased significantly in the three groups of aerobic training, supplementation, and training-supplementation, as compared to the pretest levels [Table 1].

Results of one-way ANOVA also revealed significant differences in  $\text{IL-6}$  ( $P=0.0001$ ),  $\text{IL-18}$  ( $0.03$ ),  $\text{IL-1}\beta$  ( $P=0.001$ ), and  $\text{TNF-}\alpha$  ( $P = 0.001$ ) mean changes [Table 1]. *Post hoc* Tukey test showed that in training + supplementation

group, mean changes of  $\text{IL-6}$  ( $P = 0.01$ ,  $P = 0.01$ ,  $P = 0.02$  respectively),  $\text{IL-18}$  ( $P = 0.01$ ,  $0.001$ ,  $P = 0.001$ ),  $\text{TNF-}\alpha$  ( $P = 0.001$ ,  $P = 0.001$ ,  $P = 0.01$  respectively), and  $\text{IL-1}\beta$  ( $P=0.03$ ,  $P=0.01$ ,  $P=0.001$ ) were significantly higher as compared to other three groups of aerobic training, supplementation, and control [Figures 2-4].

### Effects of 12 weeks of aerobic training and supplementation on body composition indices

*T*-test results revealed that in exercise + supplementation, exercise, and supplementation groups; body fat percent ( $P=0.01$ ,  $P=0.008$ ,  $P=0.01$ , respectively) and waist-hip ratio (WHR) ( $P = 0.001$ ,  $P = 0.003$ ,  $P = 0.004$ , respectively) decreased significantly following 12 weeks of intervention. Kg decreased significantly ( $P=0.04$ ,  $P=0.04$ ) as well as their BMI ( $P = 0.01$ ,  $P = 0.02$ ) in exercise + supplementation and exercise groups, respectively [Table 2].


According to Table 1 and one-way ANOVA results, between-group differences were significant in the mean changes of kg, body fat percent, and WHR ( $P=0.001$ ) [Table 2]. Results from Tukey's test revealed a pronounced impact of exercise-supplementation group rather than the three groups of aerobic training, supplementation and control in weight ( $P = 0.01$ ,  $P = 0.01$ , and  $P = 0.001$ , respectively), BMI ( $P = 0.02$ ,  $P = 0.01$ , and  $P = 0.03$ , respectively), body fat percentage ( $P = 0.04$ ,  $P = 0.04$ , and  $P = 0.01$ , respectively), and WHR ( $P = 0.01$ ,  $P = 0.01$ , and  $P = 0.001$ , respectively) mean changes.

### Comparison of calorie and macronutrients intake between groups


The mean and standard deviation of participants' calorie and macronutrients intakes are shown in Table 3. Between-group comparison is also summarized in Table 3.

## DISCUSSION


In relation to our first hypothesis, the main finding of the present investigation was a significant improvement in  $\text{IL-6}$ ,  $\text{IL-18}$ ,  $\text{IL-1}\beta$ ,  $\text{TNF-}\alpha$ , body fat percent, body weight, WHR, and BMI levels after 12 weeks of aerobic training and *F. angulata* supplementation consumption. In accordance to the present study, Mohamadzadeh Salamat *et al.* reported significant reductions in  $\text{IL-6}$ ,  $\text{IL-18}$ , and  $\text{IL-1}\beta$  levels following 8 weeks of endurance training at 75%–80% of reserved HR in forty overweighted men.<sup>[24]</sup> In Ordonez *et al.*'s study, significant reduction in  $\text{IL-6}$  and  $\text{TNF-}\alpha$  concentration has been shown after 10 weeks of aerobic training, three times a week, for 40 min, at 55%–65% of HR in obese women.<sup>[25]</sup> There are also some contradictory investigations as well. For instance, Stensvold *et al.* reported no significant differences were observed in the serum levels of  $\text{IL-6}$ ,  $\text{IL-1}\beta$ ,  $\text{TNF-}\alpha$  and body composition in 31 syndrome metabolic



**Figure 2:** Between-group differences for interleukin-1. \*Significant difference with control group ( $P < 0.05$ ). \*\*Significant difference with training-placebo ( $P < 0.05$ ). #Significant difference with training-Ferulago group ( $P < 0.05$ )



**Figure 3:** Between-group differences for interleukin-6. \*Significant difference with control group ( $P < 0.05$ ). #Significant difference with training-Ferulago group ( $P < 0.05$ )



**Figure 4:** Between-group differences for interleukin-18. \*Significant difference with control group ( $P < 0.05$ ). #Significant difference with training-Ferulago group ( $P < 0.05$ )

**Table 1: Pre- and post-test changes in levels of interleukin-6, interleukin-18, interleukin-1 $\beta$ , and tumor necrosis factor-alpha**

Variables	Groups	Mean $\pm$ SD		Significant level of t-test	Significant level of ANOVA
		Pretest	Post-test		
IL-6 (pg/mL)	Control	74.74 $\pm$ 9.89	74.09 $\pm$ 9.43	0.31	0.001*
	Ferulago extract	71.89 $\pm$ 9.48	69.26 $\pm$ 8.04	0.03*	
	Training-placebo	74.23 $\pm$ 6.65	67.41 $\pm$ 7.12	0.01*	
	Training-ferulago	77.76 $\pm$ 8.81	64.33 $\pm$ 7.12	0.001*	
IL-18 (pg/mL)	Control	97.37 $\pm$ 5.82	97.14 $\pm$ 7.21	0.35	0.03*
	Ferulago extract	107.92 $\pm$ 11.89	104.84 $\pm$ 9.17	0.03*	
	Training-placebo	107.27 $\pm$ 11.46	101.02 $\pm$ 9.23	0.02*	
	Training-ferulago	106.12 $\pm$ 12.12	96.90 $\pm$ 9.15	0.001*	
TNF- $\alpha$ (pg/mL)	Control	189.96 $\pm$ 23.40	190.70 $\pm$ 19.95	0.18	0.001*
	Ferulago extract	187.98 $\pm$ 23.58	185.66 $\pm$ 18.28	0.03*	
	Training-placebo	196.54 $\pm$ 25.07	189.61 $\pm$ 24.12	0.02*	
	Training-Ferulago	202.01 $\pm$ 19.12	188.73 $\pm$ 18.65	0.001*	
IL-1 (pg/mL)	Control	3.41 $\pm$ 0.40	3.52 $\pm$ 1.95	0.18	0.001*
	Ferulago extract	5.66 $\pm$ 2.58	4.88 $\pm$ 2.28	0.03*	
	Training-placebo	5.64 $\pm$ 2.07	3.14 $\pm$ 2.12	0.02*	
	Training-ferulago	6.41 $\pm$ 2.12	3.22 $\pm$ 1.65	0.001*	

\*Significant difference in  $P < 0.05$ . SD=Standard deviation; IL=Interleukin; TNF- $\alpha$ =Tumor necrosis factor alpha; ANOVA=Analysis of variance

**Table 2: Pre-posttest changes in weight, body mass index, fat percentage, and waist-to-hip ratio**

Variables	Groups	Mean $\pm$ SD		Significant Level of t-test	Significant Level of ANOVA
		Pretest	Posttest		
Weight (kg)	Control	82.61 $\pm$ 4.33	83.15 $\pm$ 4.92	0.31	0.001*
	Ferulago extract	83.90 $\pm$ 6.85	83.51 $\pm$ 6.85	0.35	
	Training-placebo	80.90 $\pm$ 5.61	79.58 $\pm$ 5.06	0.04*	
	Training-ferulago	84.88 $\pm$ 13.86	82.06 $\pm$ 13.14	0.04*	
BMI (kg/m <sup>2</sup> )	Control	32.37 $\pm$ 2.37	32.7 $\pm$ 2.01	0.29	0.001*
	Ferulago extract	32.82 $\pm$ 2.47	32.65 $\pm$ 2.27	0.34	
	Training-placebo	32.91 $\pm$ 2.55	32.77 $\pm$ 2.23	0.02*	
	Training-ferulago	34.56 $\pm$ 4.94	33.38 $\pm$ 4.39	0.01*	
Fat percentage (%)	Control	41.74 $\pm$ 3.11	42.77 $\pm$ 2.30	0.444	0.001*
	Ferulago extract	41.22 $\pm$ 3.52	39.91 $\pm$ 3.64	0.01*	
	Training-placebo	41.9 $\pm$ 4.48	40.47 $\pm$ 4.78	0.008*	
	Training-ferulago	34.56 $\pm$ 4.94	33.38 $\pm$ 4.39	0.01*	
WHR	Control	0.98 $\pm$ 0.03	0.98 $\pm$ 0.02	0.22	0.001*
	Ferulago extract	0.99 $\pm$ 0.04	0.96 $\pm$ 0.02	0.004*	
	Training-placebo	0.99 $\pm$ 0.04	0.96 $\pm$ 0.04	0.003*	
	Training-ferulago	1.01 $\pm$ 0.07	0.98 $\pm$ 0.06	0.001*	

\*Significant difference in  $P < 0.05$ . SD=Standard deviation; ANOVA=Analysis of variance, BMI=Body mass index; WHR=Waist-to-hip ratio

sedentary participants, following 3 months performing endurance training (80%–90%  $VO_{2max}$ ).<sup>[26]</sup> This discrepancy could be due to high-intensity nature of exercise although their body composition did not change after 12-week intervention. Results from a study by Lee *et al.* also revealed no significant changes in IL-6, IL-18, IL-1 $\beta$ , and TNF-content after 12 weeks of aerobic training (50%–60%  $VO_{2max}$ ) in obese, diabetic teenagers.<sup>[27]</sup> The observed differences might be due to varieties in factors such as participants' basal healthy status, resting levels of inflammatory factors, kind of training protocol, and the participants' ages. It seems that the type of activity affects the amount of observed changes. In addition, different training durations and intensities bring

about different changes in cytokines' levels.<sup>[25,26]</sup> There are contradictory findings about the optimal training intensity for attenuating inflammatory factors.<sup>[25-27]</sup> As compared to lower intensities, high training intensities with long or medium duration exert more effects on inflammatory factors, reducing their levels to a greater extent. Fischer *et al.* showed that exercise duration was the most stimulator factor in increasing postexercise serum levels of cytokines.<sup>[28]</sup> In fact, it was claimed that 50% of cytokines' changes during exercise could be explained solely through exercise duration.

As mentioned earlier, obesity is one of the factors which augment basic levels of IL-6, IL-18, IL-1 $\beta$ , and TNF- $\alpha$

**Table 3: Pre- and post-test changes in calorie and macronutrients' intake between groups**

	Groups	Mean±SD		F	P ANOVA
		Pretest	Posttest		
Energy (Kcal)	Train+ supplement	1925.5±100.32	1920.5±99.32	0.32	0.76
	Train	1933.5±12.34	1910.5±34.34		
	Supplement	1890.5±19.05	1897.5±10.54		
	Control	1854.5±14.76	1900.5±10.01		
Carbohydrate (g)	Train+ supplement	12.45±156.5	11.03±136.31	0.06	0.55
	Train	12.32±144.3	9.54±140.55		
	Supplement	15.17±165.65	11.43±150.6		
	Control	11.43±155.03	7.05±165.00		
Protein (g)	Train+ supplement	3.05±76.05	1.14±70.54	0.12	0.21
	Train	2.02±64.66	4.76±63.06		
	Supplement	2.23±77.76	2.09±69.06		
	Control	1.87±66.88	1.11±69.08		
Total fat (g)	Train+ supplement	1.05±67.87	1.32±67.09	0.20	0.43
	Train	2.43±63.5	2.76±60.06		
	Supplement	3.22±67.07	2.86±59.99		
	Control	2.21±67.66	2.92±69.09		

SD=Standard deviation

in human through stimulating oxidative stress and inflammation. Weight loss improves serum levels of these factors through reduction in oxidative stress and inflammation status. Previous studies revealed a strong, negative correlation between IL-6, IL-18, IL-1 $\beta$ , and TNF- $\alpha$  concentration with body fat percent and body composition indices.<sup>[13,26]</sup> The exact reason for this negative correlation has not been fully clarified, but the effect of physical activity on adipose tissue probably is an effective factor.<sup>[28]</sup> The amount of adipose tissue has the highest correlation with circulating inflammatory indices. Lower levels of inflammation in active people relate to their less amount of net visceral fat.<sup>[13,28]</sup> Indeed, as adipose tissue is one of the main producers of IL-6, IL-18, IL-1 $\beta$ , and TNF- $\alpha$ , its reduction reduces serum levels of mentioned cytokines. In the present study, weight, BMI, WHR, and body fat percent decreased significantly following 12 weeks of exercise training. Our findings also suggested that body composition indices (body fat percent, BMI, and WHR) are highly correlated with IL-6, IL-18, IL-1 $\beta$ , and TNF- $\alpha$ , respectively. This improvement in participants' body composition could probably correlate with reduction in mentioned cytokines' levels, free radicals, and oxidative stress in obese men.

Here, we showed daily consumption of 50 mg/kg *F. angulata* for 12 weeks leads to significantly reduction in IL-6, IL-18, IL-1  $\beta$ , TNF- $\alpha$ , body fat percent, and WHR levels. The antioxidant and antilipoid effects of *F. angulata* plant have been investigated in previous studies.<sup>[17]</sup> It was also reported that *F. angulata* extract has a positive effect on immune indices and activity.<sup>[18]</sup> Rafieian-Kopaei *et al.* reported low detected serum levels of total cholesterol and triglyceride after taking *F. angulata* extract caused inhibition of per

oxidation of lipids.<sup>[29]</sup> Amirghofran *et al.* did an investigation on 8-week-old male mice and described that *F. angulata* extract (50 mg/ml) significantly decreased production of nitrite oxide after 24 h. In follow-up, 48 h after consumption of this extract level of inflammatory blood laboratory cytokines such as (IL-1 $\beta$ ) decreased significantly.<sup>[19]</sup>

Niu *et al.* believed that flavonoids existing in herbs (gene activator transcription factors such as PGC 1 $\alpha$ ) may decline differentiation and multiplication of adipocytes and reduces the expression of genes responsible for lipogenesis. In this way, not only IL-6, IL-8, and TNF- $\alpha$  levels diminish but also adiponectin levels raise and blocks catechol-*o*-methyltransferase enzyme which eventually linked to decreases fat and body weight.<sup>[30]</sup> After 12 weeks of aerobic training with *F. angulata* supplementation, the present researcher showed body fat percent, BMI, WHR, body weight and as a result levels of IL-6, IL18, IL-1 $\beta$ , and TNF- $\alpha$  decreased significantly. As the observed improvement was greater in exercise-supplementation group compared to other intervention groups, *F. angulata* can be considered as a strong, anti-inflammatory and anti-oxidative supplement, which could be considered effective in obesity severity along with aerobic training. It seems physical activity might affect cytokines' production through different mechanisms such as changes in circulating blood factors (e.g. lactate, catecholamine, and growth factors) stimulation of lymphatic system nodes, making NK cells move faster than T-cells.<sup>[26]</sup>

In addition, as lymphatic system gets more stimulated, more cytokines are released from the adipose tissue and it has been proven that exercise reduces sympathetic stimulation. It seems that an increase in expression of

muscles' pro-inflammatory IL receptors in people with high  $VO_{2\max}$ , which is a consequence of exercising, could justify the decrease in serum levels of these indices.<sup>[27,29]</sup> In addition, flavonoids in *F. angulata* improve body's anti-oxidation system through increasing inter-cellular anti-oxidative levels such as glutathione, uric acid, and bilirubin, improved intercellular anti-oxidative enzymes capacity such as glutathione peroxides, and catalase. They also prevent lipogenesis and improve fat oxidation through blocking phospholipase A2, and inhibition of transcription factors of free fatty acid synthesis and blocking acetyl-CoA carboxylase.<sup>[30]</sup> The effects of this supplementation would be more vigorous potentially if they are combined with moderate-to-intense physical activity.<sup>[6,28]</sup>

## CONCLUSION

Although consumption of *F. angulata* leads to an improvement in pro-inflammatory indices in obese young men, it seems that *F. angulata* consumption along with a 12-week aerobic exercise has profound effect on these indices, and we can recommend it as a secondary outcome that nonathlete obese people can take *F. angulata* and perform aerobic exercise to improve body composition and inflammatory system. We suggest further studies with more sample size and precise assessment to conclude whether the positive effect of training with *F. angulata* is permanent or not. In addition, while double blinding condition was not performed in present study, it should be considered in future probes.

## Acknowledgments

The authors would like to thank all the participants in this study. The study was approved by the Ethic committee of the Medical University of Abadan under the 93U-044 code and the Iranian Registry of Clinical Trials code is (www.IRCT.IR; IRCT20160129026251N4).

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Hovsepian S, Haghjooy Javanmard SH, Mansourian M, Hashemipour M, Tajadini M, Kelishadi R. Lipid regulatory genes polymorphism in children with and without obesity and cardiometabolic risk factors: The CASPIANIII study. *J Res Med Sci* 2018;23:11.
- Ellulu MS, Patimah I, Khaza'ai H, Rahmat A, Abed Y. Obesity and inflammation: The linking mechanism and the complications. *Arch Med Sci* 2017;13:851-63.
- Kreuter R, Wankell M, Ahlenstiel G, Hebbard L. The role of obesity in inflammatory bowel disease. *Biochim Biophys Acta Mol Basis Dis* 2019;1865:63-72.
- Flores A, Burstein E, Cipher DJ, Feagins LA. Obesity in inflammatory bowel disease: A marker of less severe disease. *Dig Dis Sci* 2015;60:2436-45.
- Donges CE, Duffield R, Drinkwater EJ. Effects of resistance or aerobic exercise training on interleukin-6, C-reactive protein, and body composition. *Med Sci Sports Exerc* 2010;42:304-13.
- Dorneles GP, Haddad DO, Fagundes VO, Vargas BK, Kloecker A, Romão PR, et al. High intensity interval exercise decreases IL-8 and enhances the immunomodulatory cytokine interleukin-10 in lean and overweight-obese individuals. *Cytokine* 2016;77:1-9.
- Martín-Cordero L, García JJ, Hinchado MD, Bote E, Manso R, Ortega E. Habitual physical exercise improves macrophage IL-6 and TNF- $\alpha$  deregulated release in the obese Zucker rat model of the metabolic syndrome. *Neuroimmunomodulation* 2011;18:123-30.
- Bishop NC. Exercise, the risk of infection and immune function in special populations. In: Gleeson M, editor. *Immune Function in Sport and Exercise*. 2<sup>nd</sup> ed. Tehran: Hatmi; 2015. p. 398-422.
- Dow CA, Stauffer BL, Brunjes DL, Greiner JJ, DeSouza CA. Regular aerobic exercise reduces endothelin-1-mediated vasoconstrictor tone in overweight and obese adults. *Exp Physiol* 2017;102:1133-42.
- Niemen D. Exercise, immunology and nutrition. *World Rev Nutr Diet* 2009;90:89-101.
- Beavers KM, Brinkley TE, Nicklas BJ. Effect of exercise training on chronic inflammation. *Clin Chim Acta* 2010;411:785-93.
- Chen CW, Chen CC, Jian CY, Lin PH, Chou JC, Teng HS, et al. Attenuation of exercise effect on inflammatory responses via novel role of TLR4/PI3K/Akt signaling in rat splenocytes. *J Appl Physiol* (1985) 2016;121:870-7.
- Sloan RP, Shapiro PA, McKinley PS, Bartels M, Shimbo D, Lauriola V, et al. Aerobic exercise training and inducible inflammation: results of a randomized controlled trial in healthy, young adults. *J Am Heart Assoc* 2018;7:e010201.
- Auerbach P, Nordby P, Bendtsen LQ, Mehlsen JL, Basnet SK, Vestergaard H, et al. Differential effects of endurance training and weight loss on plasma adiponectin multimers and adipose tissue macrophages in younger, moderately overweight men. *Am J Physiol Regul Integr Comp Physiol* 2013;305:R490-8.
- Silva CM, Vieira-Junior RC, Trombeta JC, Lima TR, Fraga GA, Sena MS, et al. Effects of aerobic and resistance training of long duration on pro- and anti-inflammatory cytokines in rats. *Revista Andaluza De Med Dep* 2017;10:170-5.
- Libardi CA, De Souza GV, Cavaglieri CR, Madruga VA, Chacon-Mikahil MP. Effect of resistance, endurance, and concurrent training on TNF- $\alpha$ , IL-6, and CRP. *Med Sci Sports Exerc* 2012;44:50-6.
- Ghasemi Pirbalouti A, Izadi A, Malek Poor F, Hamedi B. Chemical composition, antioxidant and antibacterial activities of essential oils from *Ferulago angulata*. *Pharm Biol* 2016;54:2515-20.
- Valipour P, Heidarian E, Khoshdel A, Gholami-Arjenaki M. Protective effects of hydroalcoholic extract of *Ferulago angulata* against gentamicin-induced nephrotoxicity in rats. *Iran J Kidney Dis* 2016;10:189-96.
- Amirghofran Z, Malek-Hosseini S, Golmoghaddam H, Kalantar F, Shabani M. Inhibition of nitric oxide production and proinflammatory cytokines by several medicinal plants. *Iran J Immunol* 2011;8:159-69.
- Jackson AS, Pollock ML, Ward A. Generalized equations for predicting body density of women. *Med Sci Sports Exerc* 1980;12:175-81.
- Wilmore JH, Costill DL. *Physiology of Sport and Exercise*. 6<sup>th</sup> ed. Champaign, IL: Human Kinetics; 2005.

22. Lin D, Tsuzuki E. Effect of methanol extracts from *Ophiopogon japonicus* on rice blast fungus. *Pest Sci Manag* 2003;28:27-8.
23. Mahmoudi R, Rafieirad M, Goudarzi S. Effect of hydroalcoholic extract of *Ferulago angulata* (Schlecht) Boiss on motor and memory disorders in animal model of Parkinson disease. *Qom Univ Med Sci J* 2018;12:36-47.
24. Mohamadzadeh Salamat K, Azarbayjani ML, Ashril Yusof C, Dehghan F. The response of pre-inflammatory cytokines factors to different exercises (endurance, resistance, concurrent) in overweight men. *Alexandria J Med* 2016;52:367-70.
25. Ordonez FJ, Rosety MA, Camacho A, Rosety I, Diaz AJ, Fornieles G, *et al.* Aerobic training improved low-grade inflammation in obese women with intellectual disability. *J Intellect Disabil Res* 2014;58:583-90.
26. Stensvold D, Slørdahl SA, Wisløff U. Effect of exercise training on inflammation status among people with metabolic syndrome. *Metab Syndr Relat Disord* 2012;10:267-72.
27. Lee SS, Yoo JH, Kang S, Woo JH, Shin KO, Kim KB, *et al.* The effects of 12 weeks regular aerobic exercise on brain-derived neurotrophic factor and inflammatory factors in juvenile obesity and type 2 diabetes mellitus. *J Phys Ther Sci* 2014;26:1199-204.
28. Fischer CP, Berntsen A, Perstrup LB, Eskildsen P, Pedersen BK. Plasma levels of interleukin-6 and C-reactive protein are associated with physical inactivity independent of obesity. *Scand J Med Sci Sports* 2007;17:580-7.
29. Rafieian-Kopaei M, Shahinfard N, Rouhi-Boroujeni H, Gharipour M, Darvishzadeh-Boroujeni P. Effects of *Ferulago angulata* extract on serum lipids and lipid peroxidation. *Evid Based Complement Alternat Med* 2014;2014:1-4.
30. Niu Y, Na L, Feng R, Gong L, Zhao Y, Li Q, *et al.* The phytochemical, EGCG, extends lifespan by reducing liver and kidney function damage and improving age-associated inflammation and oxidative stress in healthy rats. *Aging Cell* 2013;12:1041-9.