



Published in final edited form as:

Pediatr Res. 2022 March ; 91(4): 955–961. doi:10.1038/s41390-021-01542-2.

Seizure burden in preterm infants and smaller brain volume at term equivalent age

Zachary Vesoulis¹, Dimitrios Alexopoulos², Cynthia Rogers^{1,3}, Jeffrey Neil^{1,2,4}, Christopher Smyser^{1,2,4}

¹Department of Pediatrics, Division of Newborn Medicine, Washington University

²Department of Neurology, Division of Child Neurology, Washington University

³Department of Psychiatry, Washington University

⁴Department of Radiology, Washington University

Abstract

Background: Seizures are underrecognized in preterm infants, and little is known about their impact on brain growth. We aimed to define the association between early seizures and subsequent brain growth.

Methods: Infants <30 weeks' gestation underwent 72h of prospective aEEG monitoring, term-equivalent age (TEA) MRI, and 2y neurodevelopmental testing.

Seizures were defined as trains of sharp waves >10 seconds, evolving in frequency/amplitude/morphology, and identified using automated algorithms with manual review.

Using T2-weighted images, cortical surface area (CSA) and gyrification index (GI) were calculated and volumes were segmented into five tissue classes: CSF, gray matter, white matter (WM), deep nuclear gray matter, and cerebellum. Correlations between total seizure burden and tissue-specific volumes were evaluated, controlling for clinical variables of interest.

Results: Ninety-nine infants underwent aEEG/MRI assessments (mean GA=26.3 weeks, birthweight=899g). Seizure incidence was 55% with a median of two events; median length=66s and mean burden=285s.

Greater seizure burden was associated with smaller CSA and volumes across all tissue types, most prominently in WM ($R^2=-0.603$, $p<.01$), even after controlling for confounders. There was no association with GI.

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Corresponding Author: Zachary A. Vesoulis MD MSCI, Postal address: 1 Children's Place, Box 8116, St. Louis, MO USA 63110, Phone: 314-286-1524, vesoulis_z@wustl.edu.

Contributor Statement: ZAV developed the study idea, interpreted, and collected aEEG data, performed statistical analysis, generated figures, and wrote the initial manuscript draft. DA collected and analyzed MRI data, generated figures, and critically reviewed the manuscript. CR, JN, and CS participated in study development, and critically reviewed the manuscript. All authors reviewed and approved of the final manuscript.

Disclosure statement: No authors have financial ties or potential/perceived conflicts to disclose.

Category of Study: Clinical Research

Consent: Informed written consent was obtained for all participants prior to any study procedures.

Conclusion: Seizures in preterm infants are common and associated with smaller TEA brain volumes. This relationship was strongest for WM and independent of clinical factors.

INTRODUCTION

Previous investigations of seizures in preterm infants are heterogeneous in cohort composition and study design, and estimates of seizure incidence vary. Nevertheless, preterm infants are known to have more seizures than term infants (1-3), and a significant proportion is subclinical (4). Importantly, there is growing evidence that seizures may be associated with adverse neurologic sequelae in this population, including brain injury, intellectual impairment, and cerebral palsy (5-8). Although there is significant overlap between seizures and preterm brain injury, most notably intraventricular hemorrhage (IVH) (6,9-11), a pathophysiological link has not been established. It is possible that the adverse outcomes associated with seizures in infants with IVH are a direct result of the seizures, but they may also be a consequence of associated brain injury unrelated to seizures and occurring during a critical developmental window (12) when the brain is undergoing rapid changes in physical size (13-15), gyrification (16,17), and synaptic density (18). While the relative contributions of these two mechanisms remain unclear, seizures and/or IVH may permanently alter the trajectory of brain growth.

Evaluating potential mechanistic connections between preterm seizures and subsequent adverse neurologic consequences is a high priority, as it carries important implications for EEG monitoring and seizure treatment in the Neonatal Intensive Care Unit (NICU). If seizures are indeed an independent risk factor for adverse brain growth, approaches for targeted evaluation and effective management must be developed and implemented. In this manuscript, we describe the results from an investigation which aimed to define the association between seizure burden during the first 72 hours of life and brain volume, cortical surface area, and cortical folding at term equivalent age (TEA) in a cohort of very preterm infants (born prior to 30 weeks' gestation). We also examined the additional discriminatory value EEG and volumetric measures bring to the prediction of adverse neurodevelopmental outcomes. We hypothesized that early seizures would be associated with smaller brain volumes, cortical surface area, and folding at TEA, with varying effects according to tissue type.

METHODS

Cohort

Very preterm infants born before 30 completed weeks' gestation were prospectively recruited for a longitudinal cerebral monitoring study between 2008 and 2016. All infants were hospitalized in the Neonatal Intensive Care Unit (NICU) at St. Louis Children's Hospital, a level IV NICU serving urban, suburban, and rural populations. Infants were excluded if there was a known congenital or genetic anomaly or if the infant did not survive to TEA MRI. Detailed clinical and demographic characteristics were collected from the medical record, including gestational age, sex, birth weight, Apgar scores at 1 and 5 minutes, CRIB-II score (19), antenatal steroid exposure, bronchopulmonary dysplasia (BPD, defined as supplemental oxygen beyond 36 weeks), postnatal steroid exposure, sepsis

(defined as culture positive sepsis), and necrotizing enterocolitis (NEC, defined as Bell stage IIA or greater).

aEEG monitoring and seizure classification

All infants underwent continuous two-channel amplitude-integrated EEG (aEEG) recording from the time of recruitment through 72 hours of age. All monitoring was performed with the BRM3 aEEG monitor (Natus Medical, San Carlos, CA) using hydrogel electrodes arranged in the typical C3-P3, C4-P4 montage. As routine EEG/aEEG monitoring is not standard of care in this population, and the risk/benefit of seizure treatment has not yet been established, the clinical team was blinded to the monitor, and all recordings were archived and analyzed after the infant was discharged.

Detection of seizures on a limited-channel montage was enhanced by use of a multimodal approach: utilizing an automated seizure detection algorithm, identifying changes in the aEEG baseline, and confirmation with the raw EEG. Each recording was initially screened by assessing for characteristic changes in the aEEG baseline (sudden change in the upper and lower margin of the time-compressed tracing or a “saw tooth” pattern (20)) or for marks placed by an automatic seizure detection algorithm. Seizures were confirmed or excluded by inspection of the raw EEG at the identified time points. Seizure-like artifactual events, such those induced by high-frequency oscillatory ventilation (21), EKG artifact, or muscle artifact, were excluded (22). As brief seizures are not well detected by automated algorithms or aEEG baseline changes (23,24), the entire raw trace was also manually inspected for seizure activity. All EEG studies were scored by a single reviewer (Z.A.V.) with extensive experience in reading aEEG recordings of preterm infants.

Seizures were defined in the manner described by Scher et al. (25,26); namely, a series of sharp waves at least 10 seconds in length which evolve in frequency, amplitude, and morphology and are clearly distinguishable from the background and artifact. Periodic epileptiform discharges (5) were not classified as seizures. Cumulative seizure burden and average seizure length were recorded.

Neuroimaging

All surviving infants underwent non-contrast brain MRI at TEA. All scans were performed using a 3T TIM Trio system (Siemens, Erlangen, Germany) without sedating medications using an infant-specific head coil. T2-weighted turbo-spin echo sequence parameters included TR 8600 ms, TE 160 ms, and voxel size 1 mm×1 mm×1 mm.

Brain volumes were delineated on native space T2-weighted images. Total and regional brain volumes were estimated using the Advanced Normalization Tools (ANTs) pipeline, which utilizes registration, bias correction, and tissue segmentation tools to develop probabilistic maps of tissue-specific regional volumes, followed by manual editing by a single highly experienced rater (D.A.) using ITK-Snap software tools (27,28). Calculated volumes consisted of cerebrospinal fluid (CSF), cortical gray matter, white matter (WM), deep nuclear gray matter, and cerebellum. Examples of segmented and labeled images are shown in Figure 1. Regional brain volumes were evaluated both uncorrected and corrected for total intracranial volume (ICV).

Cortical surface area (CSA) and gyrification index (GI), a measure of cortical folding, were generated. The cortex was reconstructed to provide mid-thickness, inflated, flat, spherical, and cerebral hull surfaces (29). The gyrification index was calculated as the ratio of the area of the mid-thickness cortical surface to the area of the cerebral hull (30). Both measures were generated for each hemisphere using the surfaces in the individual's native space after alignment along the anterior commissure–posterior commissure axis. For both CSA and GI, the average of the left and right hemispheres was used for analysis.

Qualitative assessment of brain injury was performed, including assessment of IVH, white matter injury (WMI), and cerebellar hemorrhage (CH). IVH was assessed using screening cranial ultrasounds performed in the first month of life as part of routine clinical care and classified based on highest grade of injury on either side using the four-grade Papile scale (31). WMI was assessed on TEA MRI and qualitatively graded as none, punctate, or cystic. CH was also identified as present or absent on TEA MRI. As MRI has superior sensitivity for WMI and CH, ultrasound data were not used for identification of either form of injury (32-34).

Neurodevelopmental outcome

All surviving infants underwent comprehensive neurodevelopmental testing between 18 and 24 months' corrected age using the Bayley Scales of Infant Development, Third Edition (35). Composite scores in the cognitive, language, and motor domains were obtained, and infants were classified as having moderate impairment if the score in any one of the three domains was less than 85 (–1 SD) or severe impairment if the score in any one domain was less than 70 (–2 SD).

Statistical approach

Univariate analysis of relevant demographic and clinical variables was performed using Fisher's exact test for categorical variables and the Mann-Whitney U-test for continuous variables. Univariate correlation was assessed with Pearson's correlation coefficient, and multivariate linear regression was used to assess the strength of relationship between cumulative seizure burden, CSA, GI, and tissue-specific regional brain volume (corrected and uncorrected for ICV), controlling for key clinical factors which may independently alter brain volumes and/or seizure incidence (gestational age, antenatal steroids, grade of IVH, grade of WMI, CH, culture positive sepsis, NEC, BPD, and postnatal steroids). Given the potential for non-independence imparted by inclusion of infants of multiple gestations, linear mixed-effect modeling was used with family as a random variable, an approach previously used in neonatal studies (36,37).

Our group and others have previously examined the relationship between neurodevelopmental outcome and preterm seizure burden (3,6) or tissue volumes (15,38,39). However, it is likely the each of these measures (seizure burden, volumetric data, clinical data) provide non-overlapping information about neurodevelopmental outcome. The additive value of regional volumetric data, CSA, and cumulative seizure burden to predict adverse neurodevelopmental outcome was assessed using binary logistic regression and area under the curve (AUC) analysis.

Statistical tests were considered significant at $p < 0.05$. All statistical testing was performed using R version 4.0.2 (R Foundation for Statistical Computing, Vienna, Austria). The 'nlme' package was used for mixed effects modeling.

RESULTS

Cohort description

A total of 207 infants were initially recruited for the study, and a total of 99 infants met all inclusion criteria, survived to TEA, and had MRI scans of sufficient quality for volumetric analysis. For the 108 excluded infants, 32/108 (30%) died prior to MRI, 7/108 (6%) withdrew from the study, 24/108 (22%) were not available for scan during the term equivalent time window (discharged prior to TEA or were not clinically stable to travel to the scanner), and 42/108 (39%) had MR imaging of insufficient quality for volumetric and surface-based analysis. Excluded infants were more immature (mean GA 25.8 vs. 26.3 weeks, $p < 0.01$), more often male (64 vs. 47%, $p < 0.01$), and died more often (30% vs. 2%, $p < 0.01$). Birthweight and IVH incidence were not statistically different between included and excluded infants.

For the 99 included infants, mean gestational age was 26.3 ± 1.8 weeks, mean birthweight was 899 ± 234 g, and 54% of infants were female. The median one- and five-minute Apgar scores were 4 and 7, respectively. Most infants received antenatal betamethasone (90/99), and approximately half (48/99) received a complete two-dose course prior to delivery. As survival to TEA was an inclusion criterion, death was a rare event, occurring only twice in the cohort. A complete summary of clinical and demographic factors can be found in Table 1.

Brain injury outcomes

All infants included in the study cohort had high-quality (i.e., low motion) MRI scans, with a median postmenstrual age (PMA) at scan of 38 weeks (range 35-42 weeks). The first screening head ultrasound was performed at a median age of 1 day (range 0-12 days) and IVH was first diagnosed at a median age of 3 days (range 0-10 days). The incidence of IVH was comparable to other published studies of VLBW infants, with IVH of any grade in 29% of infants and grade III/IV IVH in 9% of infants. WMI was identified in 24% (24/99) of infants, one-third (7/24) of which was cystic. CH was noted in a similar proportion, with identification in 23% (23/99) of infants.

Seizure outcomes

The median duration of aEEG recording was 70 hours. At least one seizure was detected in 55% of infants and infants had a median of two seizure events. The median length of individual seizures was 66 seconds (range 10-318 seconds) and mean cumulative seizure burden was 285 seconds (range 0-4425 seconds). Of note, no infants were identified clinically as having seizures; they were all identified only by electrographic means. As the recordings were not evaluated until after discharge, no infants received anti-epileptic medications for early seizures. Although none of the infants in this study had additional prospective EEG monitoring, three infants were suspected to have clinical seizures later in

their hospital course. None of these events were confirmed to be electrographic seizures on subsequent EEG monitoring.

Brain injury of any type or severity was associated with an increased mean cumulative seizure burden (408 seconds vs. 173 seconds, $p=0.04$). There was no association between cumulative seizure burden and culture positive sepsis, NEC, or BPD.

Volumetric analysis

The mean total cerebral volume was 280 ± 37 mL and mean CSF volume was 111 ± 22 mL. Tissue-specific mean volumes can be found in Table 2.

Increasing seizure burden was associated with smaller volumes across all tissue types, most prominently in the WM ($R^2 = -0.603$, $p < .01$; Figure 2 and Table 2). In linear mixed-effects modeling, the association between cumulative seizure burden and each tissue type volume remained significant even when controlling for gestational age, antenatal steroids, brain injury, NEC, culture positive sepsis, BPD, postnatal steroid exposure, and multiple gestation ($p < 0.01$). The strength of association was strongest for WM ($t = -6.34$) but was significant for all tissue classes (gray matter $t = -2.89$, deep nuclear gray matter $t = -3.91$, cerebellum $t = -3.41$). Substituting CRIB score for gestational age (to account for severity of illness) did not change the outcomes, likely because of the high degree of collinearity between the two values ($R^2 = -0.91$).

In addition to smaller volumes across tissue types, there was a notable association between greater seizure burden, smaller total intracranial volume, and greater CSF volume. When correcting for total intracranial volume, WM and CSF volumes remained significantly associated with cumulative seizure burden (Table 3).

Cortical surface measures

CSA and GI measures were generated for all 99 infants. The mean CSA was 257 ± 38 cm² and mean GI was 1.79 ± 0.12 , consistent with previously published reports. Smaller mean CSA was associated with increasing cumulative seizure burden, even after correcting for gestational age, antenatal steroids, brain injury, NEC, BPD, culture positive sepsis, postnatal steroid exposure, and multiple gestation ($t = -3.71$, $p < 0.01$). There was no association between mean GI and cumulative seizure burden ($t = -1.46$, $p = 0.14$). There were no notable differences in association between left and right measures of CSA or GI. A summary of all models is shown in Supplemental Table 1.

Neurodevelopmental outcomes

As two infants died after TEA but before neurodevelopmental testing, 97 infants were eligible for follow-up at age 18 to 24 months. Eighty-two of these infants returned for neurodevelopmental testing, for a follow-up rate of 85%. The median cognitive, language, and motor composite scores were 85, 89, and 85, respectively. Most infants were identified as having moderate impairment (52/82, 63%), while a smaller number had severe impairment (12/82, 15%; Supplemental Table 2). A baseline model consisting of standard clinical variables (gestational age, antenatal steroids, presence of brain injury, NEC, BPD,

sepsis, and postnatal steroids) was moderately predictive of severe outcome with an AUC of 0.787. Incorporation of regional MRI volumes and cumulative seizure burden in the model increased the strength of the prediction, reflected as higher AUC values of 0.865 and 0.878, respectively.

DISCUSSION

These results demonstrate an association between TEA brain volumes, average CSA, and cumulative seizure burden as measured by two-channel EEG in the first 72 hours following birth in preterm infants. This association remains significant even when controlling for other clinical factors which might alter seizure incidence or brain development, including gestational age, antenatal steroid exposure, and all three common forms of preterm brain injury (i.e., IVH, WMI, CH). There was no significant association between seizure burden and GI. While cumulative seizure burden was associated with smaller volumes across all tissue types at TEA, it was most strongly linked with smaller WM volumes.

Consistent with other limited-channel EEG studies, we noted a high incidence (55%) of seizures which were exclusively without clinical correlate. Other prospective studies have reported similar frequencies of electrographic seizures identified using aEEG in this population, with rates between 22 and 63% (5,6,9). This high incidence of subclinical seizures across cohorts highlights the necessity of prospective monitoring to measure seizure incidence in this population, and it is clear that reliance on clinical detection alone will lead to significant underreporting (2,40). As is common in investigations of low birthweight infants, rates of brain injury across these studies were also high, with neuroimaging abnormalities detected in approximately 40% of infants. While this injury might independently drive risk for seizures or alter trajectories of brain growth, in this study, relationships between seizure burden and brain volumes remain significant across tissue types even when controlling for these factors.

Brain injury in preterm infants is complex and appears to be caused by a variety of dysmaturational disturbances that affect both white and gray matter, a condition sometimes referred to as encephalopathy of prematurity (4). It is interesting that WM volumes were most strongly associated with seizure burden, as the WM is particularly vulnerable at this developmental stage (41-43). It should also be noted that the distinction between WM and gray matter is somewhat arbitrary, as virtually all axons have their origin and termination in gray matter. Thus, injury to one tissue type would be expected to affect the other. This destruction-development link in gray and white matter is further supported by an autopsy series in which WMI was associated with thalamic and cortical gray matter atrophy (44,45).

The role of seizures in this process is not clear, though seizures have been implicated as a possible cause of brain injury in human and animal models (46). Miller et al. demonstrated abnormal cerebral metabolism and evidence of neuronal injury in proportion to seizure burden in a cohort of infants with perinatal asphyxia (47). Similarly, epilepsy has previously been implicated in microstructural WM changes (48). Further, preclinical studies support a significant deleterious impact of seizures on synaptogenesis (49,50), dendritic proliferation (51), and myelination (52), all processes that are active during the third trimester of

gestation, a period when very preterm infants are confined to the NICU. Thus, though we have shown an association between seizures and brain volumes and not cause and effect, it is plausible that seizures play an active role in causing injury that independently reduces brain volumes.

Importantly, defining seizures in neonates has been a consistent challenge across investigations, even more so in preterm infants who also display electrographic events termed “periodic epileptiform discharges” or PEDs on aEEG. PEDs are characterized by alternating sharp-slow waves with a periodicity of 0.5 to 4 seconds but, unlike classic ictal discharges, do not evolve with respect to time or location (5). PEDs were initially described in adults and older infants and are strongly associated with later epilepsy and poor outcome (53,54). For preterm infants, however, PEDs have not been associated with brain injury or poor cognitive outcome (5). Consistent with previous studies, PEDs were not classified as seizures in this investigation.

As shown in this analysis, models which incorporate information from a variety of modalities provide a more complete and accurate picture of future neurodevelopmental outcome than factors from a single domain. The stressors found in the NICU affect the brain on many scales, from micro- to macrostructural levels. Capturing each element of risk is essential for the most comprehensive view. While clinical variables alone provide a moderate level of accuracy, the incremental improvement in AUC after the addition of regional MRI volumes and cumulative seizure burden indicate that there is additional predictive value added by these variables.

As the association between preterm seizure burden, brain injury, and morbidity becomes increasingly clear (2), there is a developing urgency to understand the role of medications in treating seizures and improving outcomes. Limited existing data suggest that a reduction in seizure burden by treatment with phenobarbital may modify brain injury risk in preterm infants (55). However, caution should be urged at this juncture as the data on antiepileptic medications in the preterm population are limited and primarily retrospective in nature. This concern is further highlighted by the neurodevelopmental risks associated with use of antiepileptic drugs in neonates (56), specifically phenobarbital (57,58), the conventional first-line seizure treatment. While phenobarbital remains the most commonly used treatment (2,59), treatment failure rates as high as 63% have been reported (2). Further, in a retrospective study with limited follow-up and in which seizure burden was not accounted for, phenobarbital was associated with decreased cognitive scores at two years (60). Levetiracetam has also been explored as an alternative agent with fewer neurodevelopmental side effects (61,62), but there is concerning data about lack of efficacy (63). To reach a consensus regarding seizure treatment in preterm infants, randomized controlled trials of the effect of treatment on neurodevelopmental outcome are clearly needed.

There are several limitations of this study. First, although seizures were identified using a robust approach (aEEG baseline evaluation, algorithmic detection, confirmation on raw EEG, manual second-by-second review), seizure detection may have been further enhanced by use of conventional EEG, which provides more complete coverage of the cortex. Second, EEG recordings were made only during the first 72 hours of life in this study. Although a

consensus is beginning to emerge that seizures are more common than previously thought and vastly underrecognized in preterm infants, the true burden of seizures, which can occur over weeks, remains unknown. Third, although each included infant has robust clinical and imaging data, the study design necessarily requires survival to term-corrected age, resulting in the unavoidable exclusion of roughly half of the originally enrolled infants. Fourth, rates of injury in this cohort are significant, although not entirely surprising given the significant degree of prematurity. Finally, each infant underwent a single MRI at TEA. While this provides valuable insight into the final volume of each tissue type, longitudinal data would provide a better understanding of the altered trajectories of brain growth in this cohort.

Conclusions

This study provides indirect evidence of the deleterious impact of seizures on brain development, with a strong association noted between cumulative seizure burden and brain tissue volumes. While IVH, CH, and WMI remain important causes of subsequent neurodevelopmental impairment, seizures may drive deleterious neuropathological effects independent of macro-scale anatomic injury. Improving the quality and accuracy of outcome prediction is a high-priority goal in clinical care. The iterative improvement in model accuracy, as demonstrated by the increase in AUC from 0.783 to 0.878 after the inclusion of volumetric and seizure burden data, suggests that each of these measures provides additive, non-overlapping information about risk for subsequent neurodevelopmental impairment.

This finding has important potential implications for the evaluation and treatment of preterm seizures. Currently, there is a paucity of evidence to support routine prospective EEG monitoring for subclinical seizures in asymptomatic preterm patients or treatment of seizures if detected, particularly given the potential for harmful side effects (64) and unclear efficacy (65) of antiepileptic medication use. Based upon these results, future studies remain necessary which include longitudinal EEG monitoring to evaluate the seizure burden outside of the first 72 hours of life, comparison of the utility of conventional versus amplitude-integrated EEG (specifically if the tradeoff of decreased seizure detection is offset by better tolerance of electrodes), further solidification of seizure definitions (including PEDs) tailored to the preterm population, and, ultimately, a randomized controlled trial of treatment versus no treatment to determine if seizure treatment in very preterm infants improves brain development and long-term neurodevelopmental outcomes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Statement of financial support:

1. NIH Career Development Awards: K23 NS111086 [Vesoulis], K02 NS089852 [Smyser], K23 MH105179 [Rogers]
2. NIH Grants: P30 NS098577, R01 HD061619, R01 HD057098, R01 MH113570, P50 HD103525
3. Washington University Institute of Clinical and Translational Sciences KL2 Training Program (NIH/NCATS KL2 TR000450) [Vesoulis]

4. The Barnes-Jewish Hospital Foundation and the Washington University ICTS Clinical and Translational Funding Program (NIH/NCATS UL1 TR000448)
5. Doris Duke Charitable Foundation [Rogers]
6. Dana Foundation [Smyser]
7. Child Neurology Foundation [Smyser]
8. Cerebral Palsy International Research Foundation [Smyser]

REFERENCES

1. Scher MS. Neonatal seizure classification: A fetal perspective concerning childhood epilepsy. *Epilepsy Res* 2006;70:41–57. [PubMed: 16624524]
2. Glass HC et al. Seizures in Preterm Neonates: A Multicenter Observational Cohort Study. *Pediatr Neurol* 2017;72:19–24. [PubMed: 28558955]
3. Spagnoli C et al. Symptomatic seizures in preterm newborns: a review on clinical features and prognosis. *Ital J Pediatr* 2018;44:115. [PubMed: 30382869]
4. Volpe JJ, Volpe JJ, editors. *Volpe's neurology of the newborn*. Sixth edition. Philadelphia, PA: Elsevier; 2018.
5. Weeke LC et al. Rhythmic EEG patterns in extremely preterm infants: Classification and association with brain injury and outcome. *Clin Neurophysiol* 2017;128:2428–35. [PubMed: 29096216]
6. Vesoulis ZA et al. Early electrographic seizures, brain injury, and neurodevelopmental risk in the very preterm infant. *Pediatr Res* 2014;75:564–9. [PubMed: 24366515]
7. Davis AS et al. Seizures in extremely low birth weight infants are associated with adverse outcome. *J Pediatr* 2010;157:720–725.e1-2. [PubMed: 20542294]
8. Pinchevsky EF, Hahn CD. Outcomes following electrographic seizures and electrographic status epilepticus in the pediatric and neonatal ICUs. *Curr Opin Neurol* 2017;30:156–64. [PubMed: 28118303]
9. Shah DK et al. Electrographic seizures in preterm infants during the first week of life are associated with cerebral injury. *Pediatr Res* 2010;67:102–6. [PubMed: 19745782]
10. Lloyd RO et al. Electrographic Seizures during the Early Postnatal Period in Preterm Infants. *J Pediatr* 2017;187:18–25.e2. [PubMed: 28366355]
11. Strober JB, Bienkowski RS, Maytal J. The incidence of acute and remote seizures in children with intraventricular hemorrhage. *Clin Pediatr (Phila)* 1997;36:643–7. [PubMed: 9391738]
12. Matthews LG et al. Brain growth in the NICU: critical periods of tissue-specific expansion. *Pediatr Res* 2018;83:976–81. [PubMed: 29320484]
13. Boardman JP et al. Early growth in brain volume is preserved in the majority of preterm infants. *Ann Neurol* 2007;62:185–92. [PubMed: 17696128]
14. Padilla N et al. Brain Growth Gains and Losses in Extremely Preterm Infants at Term. *Cereb Cortex* 2015;25:1897–905. [PubMed: 24488941]
15. Kidokoro H et al. Brain injury and altered brain growth in preterm infants: predictors and prognosis. *Pediatrics* 2014;134:e444–453. [PubMed: 25070300]
16. Armstrong E et al. The Ontogeny of Human Gyrfication. *Cereb Cortex* 1995;5:56–63. [PubMed: 7719130]
17. Ajayi-Obe M et al. Reduced development of cerebral cortex in extremely preterm infants. *The Lancet* 2000;356:1162–3.
18. Huttenlocher PR, Dabholkar AS. Regional differences in synaptogenesis in human cerebral cortex. *J Comp Neurol* 1997;387:167–78. [PubMed: 9336221]
19. Parry G, Tucker J, Tarnow-Mordi W, UK Neonatal Staffing Study Collaborative Group. CRIB II: an update of the clinical risk index for babies score. *Lancet Lond Engl* 2003;361:1789–91.
20. Hellström-Westas L, De Vries LS, Rosén I. *Atlas of amplitude-integrated EEGs in the newborn*. London; Boca Raton, FL: Informa Healthcare ; Distributed in North and South America by Taylor & Francis; 2008.

21. Tao JD, Mathur AM. Using amplitude-integrated EEG in neonatal intensive care. *J Perinatol Off J Calif Perinat Assoc* 2010;30 Suppl:S73–81.
22. Sazgar M, Young MG. EEG Artifacts [Internet]. In: *Absolute Epilepsy and EEG Rotation Review*. Cham: Springer International Publishing; 2019 [cited 2020 Sep 11]. p. 149–62. Available from: http://link.springer.com/10.1007/978-3-030-03511-2_8
23. Sun J, Ma D, Lv Y. Detection of seizure patterns with multichannel amplitude-integrated EEG and the color density spectral array in the adult neurology intensive care unit. *Medicine (Baltimore)* 2018;97:e12514. [PubMed: 30235767]
24. Rakshasbhuvankar A et al. Amplitude-integrated EEG for detection of neonatal seizures: a systematic review. *Seizure* 2015;33:90–8. [PubMed: 26456517]
25. Shellhaas RA et al. The American Clinical Neurophysiology Society’s Guideline on Continuous Electroencephalography Monitoring in Neonates. *J Clin Neurophysiol Off Publ Am Electroencephalogr Soc* 2011;28:611–7.
26. Scher MS et al. Ictal and interictal electrographic seizure durations in preterm and term neonates. *Epilepsia* 1993;34:284–8. [PubMed: 8453938]
27. Tustison NJ et al. The ANTs cortical thickness processing pipeline [Internet]. In: Weaver JB, Molthen RC, editors. . Lake Buena Vista (Orlando Area), Florida, USA: 2013 [cited 2020 Sep 11]. p. 86720K. Available from: <http://proceedings.spiedigitallibrary.org/proceeding.aspx?doi=10.1117/12.2007128>
28. Tustison NJ et al. The ANTs Longitudinal Cortical Thickness Pipeline [Internet]. *Neuroscience*; 2017 [cited 2020 Sep 11]. Available from: <http://biorxiv.org/lookup/doi/10.1101/170209>
29. Hill J et al. A surface-based analysis of hemispheric asymmetries and folding of cerebral cortex in term-born human infants. *J Neurosci Off J Soc Neurosci* 2010;30:2268–76.
30. Engelhardt E et al. Regional impairments of cortical folding in premature infants. *Ann Neurol* 2015;77:154–62. [PubMed: 25425403]
31. Papile LA, Burstein J, Burstein R, Koffler H. Incidence and evolution of subependymal and intraventricular hemorrhage: a study of infants with birth weights less than 1,500 gm. *J Pediatr* 1978;92:529–34. [PubMed: 305471]
32. Parodi A et al. Accuracy of ultrasound in assessing cerebellar haemorrhages in very low birthweight babies. *Arch Dis Child - Fetal Neonatal Ed* 2015;100:F289–92. [PubMed: 25637005]
33. Benders MJNL, Kersbergen KJ, de Vries LS. Neuroimaging of White Matter Injury, Intraventricular and Cerebellar Hemorrhage. *Clin Perinatol* 2014;41:69–82. [PubMed: 24524447]
34. de Vries LS, Benders MJNL, Groenendaal F. Imaging the premature brain: ultrasound or MRI? *Neuroradiology* 2013;55 Suppl 2:13–22. [PubMed: 23839652]
35. Bayley N, Psychological Corporation. *Bayley Scales of Infant and Toddler Development: motor scale*. New York: Psychological Corp.; 2006.
36. Burkett KM et al. A Comparison of Statistical Methods for the Discovery of Genetic Risk Factors Using Longitudinal Family Study Designs. *Front Immunol* 2015;6:589. [PubMed: 26635803]
37. Shaffer ML, Kunselman AR, Watterberg KL. Analysis of neonatal clinical trials with twin births. *BMC Med Res Methodol* 2009;9:12. [PubMed: 19245713]
38. Cheong JLY et al. Brain Volumes at Term-Equivalent Age Are Associated with 2-Year Neurodevelopment in Moderate and Late Preterm Children. *J Pediatr* 2016;174:91–97.e1. [PubMed: 27174146]
39. Strahle JM et al. Impaired hippocampal development and outcomes in very preterm infants with perinatal brain injury. *NeuroImage Clin* 2019;22:101787. [PubMed: 30991622]
40. Scher MS et al. Electrographic seizures in preterm and full-term neonates: clinical correlates, associated brain lesions, and risk for neurologic sequelae. *Pediatrics* 1993;91:128–34. [PubMed: 8416475]
41. Khwaja O, Volpe JJ. Pathogenesis of cerebral white matter injury of prematurity. *Arch Dis Child - Fetal Neonatal Ed* 2007;93:F153–61.
42. the PIPARI study group et al. Preterm infants’ early growth and brain white matter maturation at term age. *Pediatr Radiol* 2013;43:1357–64. [PubMed: 23794054]

43. Jensen FE. Developmental factors regulating susceptibility to perinatal brain injury and seizures. *Curr Opin Pediatr* 2006;18:628–33. [PubMed: 17099361]
44. Pierson CR et al. Gray matter injury associated with periventricular leukomalacia in the premature infant. *Acta Neuropathol (Berl)* 2007;114:619–31. [PubMed: 17912538]
45. Nagasunder AC et al. Abnormal microstructure of the atrophic thalamus in preterm survivors with periventricular leukomalacia. *AJNR Am J Neuroradiol* 2011;32:185–91. [PubMed: 20930003]
46. Schmid R, Tandon P, Stafstrom CE, Holmes GL. Effects of neonatal seizures on subsequent seizure-induced brain injury. *Neurology* 1999;53:1754–1754. [PubMed: 10563624]
47. Miller SP et al. Seizure-associated brain injury in term newborns with perinatal asphyxia. *Neurology* 2002;58:542–8. [PubMed: 11865130]
48. Buksakowska I et al. Distinctive Patterns of Seizure-Related White Matter Alterations in Right and Left Temporal Lobe Epilepsy. *Front Neurol* 2019;10:986. [PubMed: 31632330]
49. Zhou C, Lippman JJB, Sun H, Jensen FE. Hypoxia-induced neonatal seizures diminish silent synapses and long-term potentiation in hippocampal CA1 neurons. *J Neurosci Off J Soc Neurosci* 2011;31:18211–22.
50. Katsarou A-M, Moshé SL, Galanopoulou AS. INTERNEURONOPATHIES AND THEIR ROLE IN EARLY LIFE EPILEPSIES AND NEURODEVELOPMENTAL DISORDERS. *Epilepsia Open* 2017;2:284–306. [PubMed: 29062978]
51. Casanova JR, Nishimura M, Swann JW. The effects of early-life seizures on hippocampal dendrite development and later-life learning and memory. *Brain Res Bull* 2014;103:39–48. [PubMed: 24140049]
52. Dwyer BE, Wasterlain CG. Electroconvulsive seizures in the immature rat adversely affect myelin accumulation. *Exp Neurol* 1982;78:616–28. [PubMed: 7173373]
53. Hamano K, Iwasaki N, Takeya T, Takita H. Clinical significance of periodic lateralized epileptiform discharges in children with relation to level of consciousness. *Pediatr Neurol* 1994;11:28–32. [PubMed: 7986289]
54. Chong DJ, Hirsch LJ. Which EEG Patterns Warrant Treatment in the Critically Ill? Reviewing the Evidence for Treatment of Periodic Epileptiform Discharges and Related Patterns: *J Clin Neurophysiol* 2005;22:79–91. [PubMed: 15805807]
55. Mikati MA et al. Phenobarbital modifies seizure-related brain injury in the developing brain. *Ann Neurol* 1994;36:425–33. [PubMed: 8080250]
56. Noguchi KK et al. Brain pathology caused in the neonatal macaque by short and prolonged exposures to anticonvulsant drugs. *Neurobiol Dis* 2020;105245. [PubMed: 33385515]
57. Farwell JR et al. Phenobarbital for febrile seizures--effects on intelligence and on seizure recurrence. *N Engl J Med* 1990;322:364–9. [PubMed: 2242106]
58. Kwan P, Brodie MJ. Phenobarbital for the treatment of epilepsy in the 21st century: a critical review. *Epilepsia* 2004;45:1141–9. [PubMed: 15329080]
59. Glass HC, Kan J, Bonifacio SL, Ferriero DM. Neonatal seizures: treatment practices among term and preterm infants. *Pediatr Neurol* 2012;46:111–5. [PubMed: 22264706]
60. Maitre NL, Smolinsky C, Slaughter JC, Stark AR. Adverse neurodevelopmental outcomes after exposure to phenobarbital and levetiracetam for the treatment of neonatal seizures. *J Perinatol Off J Calif Perinat Assoc* 2013;33:841–6.
61. Han JY et al. Efficacy of levetiracetam for neonatal seizures in preterm infants. *BMC Pediatr* 2018;18:131. [PubMed: 29636029]
62. Khan O et al. Role of Intravenous Levetiracetam for Acute Seizure Management in Preterm Neonates. *Pediatr Neurol* 2013;49:340–3. [PubMed: 23921284]
63. Kurtom W et al. Lack of response to treatment with levetiracetam in extreme preterm infants with seizures. *J Perinatol Off J Calif Perinat Assoc* 2019;39:1480–4.
64. Volpe JJ. *Neurology of the newborn*. 5th ed. Philadelphia: Saunders/Elsevier; 2008.
65. Weeke LC et al. Lidocaine response rate in aEEG-confirmed neonatal seizures: Retrospective study of 413 full-term and preterm infants. *Epilepsia* 2016;57:233–42. [PubMed: 26719344]

Impact:

1. Seizures in preterm infants are common
2. Little is known about the association between early seizures and later brain growth
3. Greater seizure burden is linked with smaller volumes of all brain tissue types, most prominently the white matter
4. This relationship is true even controlling for other factors
5. Additional study is needed to identify the optimal EEG monitoring and seizure treatment strategy for improved brain growth and neurodevelopmental outcomes.




Figure 1: Fully segmented and labeled brains are shown with two slices in the coronal plane. CSF is shown in pink, cortical gray matter in green, white matter in blue, deep nuclear gray matter in yellow, and cerebellum in cyan. Infant A had 14 electrographic seizures, each lasting approximately 100 seconds each. Total white matter volume was 114.6 mL. In comparison, infant B had no detected seizures and had a total white matter volume of 160.2 mL.




Figure 2: Correlation plots for each of the four tissue types in relation to cumulative seizure burden. The line of best fit is shown. Note a significant outlier infant with a very high seizure burden.

Table 1.

Demographic and clinical factors

	Sample (n=99)
Clinical factors	
Gestational age, mean (SD), weeks	26.3 (1.8)
Birth weight, mean (SD), grams	899.2 (234.2)
Multiple gestation, n (%)	23 (23)
Female sex, n (%)	53 (54)
Apgar score, median (range)	
1 minute	4 (0-9)
5 minutes	7 (0-9)
CRIB-II score, median (range)	10 (2-17)
Antenatal steroids, n (%)	
Any doses	90 (91)
Complete course	48 (49)
Postnatal steroids, n (%)	20 (20)
BPD ^a , n (%)	66 (67)
Culture positive sepsis, n (%)	14 (14)
Necrotizing enterocolitis ^b , n (%)	8 (8)
Died, n (%)	2 (2)
Brain injury	
Highest grade of IVH, n (%)	
Any grade	29 (29)
Grade III/IV	9 (9)
White matter injury, n (%)	
None	75 (76)
Punctate	17 (23)
Cystic	7 (7)
Cerebellar hemorrhage, n (%)	23 (23)
Early and late seizure outcomes^c	
Early seizure incidence, n (%)	54 (55)
Number of early seizure events, median (range)	1 (0-28)
Early seizure event length, mean (SD), seconds	82 (79)
Cumulative early seizure burden, mean (SD), seconds	285 (644)
Late clinical seizure incidence, n (%)	3 (3)

^a defined as need for supplemental oxygen after 36 weeks PMA

^b Bell stage IIA or greater

^c early seizures defined as those within 72 hours of birth, late seizures are those past 72 hours.

Table 2.

Descriptive and univariate analysis for regional volumes

Volume	Volume mean (SD), mL	Association with Cumulative Seizure Burden, R ²	p Value
Cerebrospinal fluid	111.0 (21.8)	0.310	<0.01
Gray matter	110.2 (21.1)	-0.274	0.01
White matter	130.1 (16.4)	-0.603	<0.01
Deep nuclear gray matter	22.46 (2.6)	-0.403	<0.01
Cerebellum	17.3 (3.3)	-0.375	<0.01
Total brain	280.0 (36.6)	-0.491	<0.01

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3.

Multivariate modeling of regional volume by cumulative seizure burden

Regional Volume Name	Uncorrected, t Value	Uncorrected, p Value	Corrected, t Value	Corrected, p Value
CSF	-2.51	0.01	3.52	<0.01
Gray Matter	-2.89	<0.01	0.55	0.59
White Matter	-6.34	<0.01	-2.10	0.03
Deep Nuclear Gray Matter	-3.91	<0.01	-0.93	0.38
Cerebellum	-3.41	<0.01	-0.49	0.63

Values provide for volumes uncorrected and corrected for total intracranial volume. All models adjusted for gestational age, antenatal steroids, WMI severity, IVH severity, presence of CH, necrotizing enterocolitis, culture positive sepsis, BPD, and postnatal steroid exposure.