



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib



Data Article

Dataset on species incidence, species richness and forest characteristics in a Danish protected area



Adriano Mazziotta^{a,*}, Jacob Heilmann-Clausen^a,
Hans Henrik Bruun^b, Örjan Fritz^c, Erik Aude^d,
Anders P. Tøttrup^a

^a Center for Macroecology Evolution and Climate, Natural History Museum of Denmark, University of Copenhagen, Copenhagen, Denmark

^b Section for Ecology and Evolution, Department of Biology, University of Copenhagen, Copenhagen, Denmark

^c Naturcentrum AB, Sweden

^d HabitatVision A/S, Denmark

ARTICLE INFO

Article history:

Received 21 September 2016

Received in revised form

19 October 2016

Accepted 25 October 2016

Available online 1 November 2016

Keywords:

Bryophytes

Forest restoration

Lichens

Space-for-time substitution

Vascular plants

Wood-inhabiting fungi

ABSTRACT

The data presented in this article are related to the research article entitled “Restoring hydrology and old-growth structures in a former production forest: Modelling the long-term effects on biodiversity” (A. Mazziotta, J. Heilmann-Clausen, H. H. Bruun, Ö. Fritz, E. Aude, A.P. Tøttrup) [1]. This article describes how the changes induced by restoration actions in forest hydrology and structure alter the biodiversity value of a Danish forest reserve. The field dataset is made publicly available to enable critical or extended analyses.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

DOI of original article: <http://dx.doi.org/10.1016/j.foreco.2016.09.028>

* Corresponding author at: Center for Macroecology Evolution and Climate, Natural History Museum of Denmark, University of Copenhagen, Universitetsparken 15, Copenhagen, Denmark.

E-mail addresses: a_mazziotta@hotmail.com, adriano.mazziotta@snm.ku.dk (A. Mazziotta), jheilmann-clausen@snm.ku.dk (J. Heilmann-Clausen), hbruun@bio.ku.dk (H.H. Bruun), orjan.fritz@naturcentrum.se (Ö. Fritz), eau@habitatvision.dk (E. Aude), aptottrup@snm.ku.dk (A.P. Tøttrup).

<http://dx.doi.org/10.1016/j.dib.2016.10.021>

2352-3409/© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	Biology
More specific subject area	Forest Ecology and Management, Restoration, Conservation Biology
Type of data	Excel file
How data was acquired	Procedures of collection described in Data collection, paragraph 2.3. in Materials and methods of Mazziotta et al. [1].
Data format	Raw, Analyzed
Experimental factors	Information for each variable related with species richness and forest characteristics were aggregated at plot level.
Experimental features	The relationships between the species richness of four ecological groups and their associated habitat characteristics were determined
Data source location	Lille Vildmose nature reserve, Denmark (56°509 N, 10°159 E)
Data accessibility	The data are available within this article

Value of the data

- This data can be used to compare the structures of species communities in other temperate forests.
- This data helps to compare the effect of environmental drivers of species richness in other temperate forests.
- This data can be used as training data in modelling exercises in statistical classes.

1. Data

In this article's data file in [Supplementary material](#), "DiB_A.xls", we stored 5 excel sheets:

- In the matrices of the first four excel sheets (i.e. "bryophytes", "lichens", "fungi", "plants") we recorded species (species Latin name in Column A) incidence (presence–absence) for each ecological group analyzed in the article (i.e. epiphytic bryophytes, epiphytic lichens, vascular plants and wood-inhabiting fungi) in each of the 102 forest plots object of the 2013 monitoring campaign in the Lille Vildmose forest reserve (Denmark) (columns B–CY).
- In the last excel sheet (Table "Richness_Env" in DiB_A.xls) we recorded plot coordinates (columns A–B) and plot number (Column C), along with the forest area they belong to (Column D), the species richness of each of the 4 ecological groups analyzed (columns E–H), the values of the predictor variables used to model species richness: basal area (X_BA) (columns I–M) and deadwood (X_DW) (columns N–P) for each of the tree genera (X); Number of Trees with Rotten Parts (NTRP, column Q), Stand Age (SA, column R), number of tree genera (Tree, column S), measured water level in 2013 (WL2013, column T) and predicted water level by 2050 (WL2050, column U).

2. Experimental design, materials and methods

In the study area of Lille Vildmose we sampled species and forest characteristics in 102 circular sample plots (radius 15 m), settled at a minimum distance of 30 m and with a 15 m minimum buffer from forest edges. In each of the 102 forest plots, we recorded species presence-absence for each ecological group, i.e. vascular plants, wood-inhabiting fungi, and epiphytic bryophytes and lichens with the following procedures during 2013:

- Vascular plants were recorded in a 5 m circular plot concentric with the study plot. Vegetation was monitored in summer;
- Fruit bodies of wood-inhabiting fungi were recorded per each deadwood item in the plot in August and October);
- Epiphytic bryophytes and lichens were recorded on (1) standing live and dead trees with DBH \geq 10 cm (from 0 up to 2 m height) and on (2) stumps taller than 1.5 m; epiphytes were surveyed in April, August and October.

Species identification in all the surveys was carried out in the field if possible, otherwise with the help of a microscope in the lab.

The following variables were measured across sample plots during the species sampling in 2013, mainly following the standard protocols given in [2]: 1) dimensions and tree species for all deadwood items with diameter \geq 10 cm and length \geq 1 m; 2) all living trees with diameter \geq 10 cm identified and their DBH measured; 3) number of trees with rotten parts; 4) water level (WL) averaged from 4 different measures, each taken 5 m from the center of the plot. If WL was above ground, water surface depth was considered as a positive value. Otherwise, a small hole was dug and distance from the ground level to the raised water surface was considered as negative. If no water rose when 40 cm depth was reached, the value was recorded as " > -40 cm". Maximum WLs were measured for all the plots on 24th–25th February 2013. Future WLs by 2050 were estimated by means of hydrological models (described in [3,4]); 5) stand age for each plot was taken from the forestry maps of the area.

Acknowledgements

We thank the Aage V. Jensen Foundation for funding. AM, APT, HHB and JHC acknowledge Danish National Research Foundation for funding the Center for Macroecology, Evolution and Climate (Grant no. DNRF96). We are thankful to Andreas Malmqvist (Naturcentrum AB, Sweden) for participating in epiphytes' surveys. We thank the COWI consulting group (Denmark) for release of measured and projected data on the current and future water level.

Transparency document. Supporting information

Transparency data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2016.10.021>.

Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2016.10.021>.

References

- [1] A. Mazziotta, J. Heilmann-Clausen, H.H. Bruun, Ö. Fritz, E. Aude, A.P. Tøttrup, Restoring hydrology and old-growth structures in a former production forest: modelling the long-term effects on biodiversity, *For. Ecol. Manag.* 381 (2016) 125–133.
- [2] J.R. Fredshavn, R. Ejrnæs, B. Nygaard, Teknisk anvisning til kortlægning af terrestriske naturtyper. -Danmarks Miljøundersøgelser, Teknisk Anvisning TA-N3 ver. 1-04.
- [3] N. Riis, P.F. Møller, B. Aaby, Naturforbedring af Høstemark Skov i Lille Vildmose. Grøn driftsplan med skitseprojekt, Aage V. Jensen Naturfond (2009).
- [4] N. Riis, P.F. Møller, B. Aaby, Naturforbedring af Tofte Skov i Lille Vildmose. Grøn driftsplan med skitseprojekt, Aage V. Jensen Naturfond (2009).