

SCIENTIFIC REPORTS



OPEN

Deregulation of the *OsmiR160* Target Gene *OsARF18* Causes Growth and Developmental Defects with an Alteration of Auxin Signaling in Rice

Jian Huang, Zhiyong Li & Dazhong Zhao

Received: 04 February 2016

Accepted: 27 June 2016

Published: 21 July 2016

MicroRNAs (miRNAs) control gene expression as key negative regulators at the post-transcriptional level. *MiR160* plays a pivotal role in *Arabidopsis* growth and development through repressing expression of its target *AUXIN RESPONSE FACTOR (ARF)* genes; however, the function of *miR160* in monocots remains elusive. In this study, we found that the mature rice *miR160* (*OsmiR160*) was mainly derived from *OsMIR160a* and *OsMIR160b* genes. Among four potential *OsmiR160* target *OsARF* genes, the *OsARF18* transcript was cleaved at the *OsmiR160* target site. Rice transgenic plants (named *mOsARF18*) expressing an *OsmiR160*-resistant version of *OsARF18* exhibited pleiotropic defects in growth and development, including dwarf stature, rolled leaves, and small seeds. *mOsARF18* leaves were abnormal in bulliform cell differentiation and epidermal cell division. Starch accumulation in *mOsARF18* seeds was also reduced. Moreover, auxin induced expression of *OsMIR160a*, *OsMIR160b*, and *OsARF18*, whereas expression of *OsMIR160a* and *OsMIR160b* as well as genes involved in auxin signaling was altered in *mOsARF18* plants. Our results show that negative regulation of *OsARF18* expression by *OsmiR160* is critical for rice growth and development via affecting auxin signaling, which will advance future studies on the molecular mechanism by which *miR160* fine-tunes auxin signaling in plants.

MicroRNAs (miRNAs), which are small (~21 nucleotides) non-coding RNAs, act as critical negative regulators by binding to mRNA complementary sequences for mRNA destabilization and translational inhibition in both plants and animals^{1–5}. In plants, primary miRNAs (pri-miRNA) are transcribed from *MIRNA (MIR)* genes. Stem-loop segments derived from pri-miRNAs are cleaved by RNase III-type endonucleases (also known as Dicers) to produce paired precursor miRNAs (pre-miRNA). After liberation of miRNA duplexes, mature miRNAs approximately 21-nucleotide long direct RNA-induced silencing complexes (RISC) to bind to their target mRNAs by complementary match, leading to cleavage or translational inhibition. In the model species *Arabidopsis*, the general roles of miRNAs are recognized by analysis of mutants that are impaired in miRNA biogenesis, while the functions of specific miRNAs have been established by either expressing miRNA-resistant versions of target genes or overexpressing miRNAs. The roles of miRNAs are also investigated by loss-of-function analyses of *MIR* genes or by interfering with mature miRNAs^{6–8}. Despite identification of a large number of miRNAs from a variety of plant species, the functions of most individual miRNAs remain unclear, particularly in plant species other than *Arabidopsis*.

In rice (*Oryza sativa* L.), miRNAs play a remarkably wide number of roles in growth and development. Rice has six *DICER-LIKE (OsDCL)* genes with *OsDCL1* being the major enzyme for producing mature miRNAs^{9–11}. Strong *OsDCL1* loss-of-function mutants are not viable, while weak lines exhibit pleiotropic defects. The rice *AGO* gene *MEIOSIS ARRESTED AT LEPTOTENE1 (MEL1)* is required for meiosis¹². Loss-of-function studies of *WAF1*, an ortholog of *Arabidopsis* *HEN1*, have shown that *WAF1* is responsible for shoot development via maintaining miRNAs and trans-acting-small interfering RNAs (ta-siRNAs)¹³. Characterization of the *IPA1 (Ideal Plant Architecture 1)* trait locus has revealed that the *OsmiR156*-controlled *OsSPL14* is critical for grain yield¹⁴.

Department of Biological Sciences, University of Wisconsin-Milwaukee, Milwaukee, WI 53211, USA. Correspondence and requests for materials should be addressed to D.Z. (email: dzhao@uwm.edu)

In addition, *OsmiR156* may target *OsSPL16* to control grain size, shape, and quality¹⁵. Furthermore, the gradually increased expression of *OsmiR156* in leaves is important for leaf development¹⁶. Overexpression of *OsmiR172* represses expression of *AP2*-like genes, which consequently causes abnormal floral meristem identity and defects in flower organ and seed development^{17,18}. *OstTIR1* and *OsAFB2* are predicted targets of *OsmiR393*. Transgenic plants overexpressing *OsmiR393a/b* have defects similar to those observed in auxin signaling mutants¹⁹. Although various miRNAs have been found in rice^{20,21}, it is imperative to study functions of individual *OsmiRNAs* in rice growth and development, particularly those of agricultural importance.

MiR160 is essential for plant growth and development⁶. In *Arabidopsis*, miR160 targets *AUXIN RESPONSE FACTOR 10* (*ARF10*), *ARF16*, and *ARF17*. Expression of the miR160-resistant version of *ARF16* (*mARF16*) results in reduced fertility and less lateral roots²². Plants expressing *mARF17* exhibited pleiotropic defects in vegetative and reproductive development²³. Analyses using similar approaches have revealed that the miR160-controlled *ARF10* is essential for seed germination and many post-embryonic growth and developmental processes through the regulation of both auxin and ABA signaling²⁴. Recent studies have shown the dormancy of *mARF10* and *mARF16* seeds is increased²⁵. In tomato, *SlymiR160* and the *SlymiR160a* target *SlyARF10* are required for floral organ and early fruit development^{26,27}. Moreover, in soybean, miR160 is involved in auxin and cytokinin signaling during nodulation²⁸. Although miR160 is conserved in plants^{29,30}, the role of miR160 in monocots is unknown. In this report, we examined expression of six *OsMIR160* genes and found that the mature *OsmiR160* was mainly derived from *OsMIR160a* and *OsMIR160b* genes. In addition, the *OsARF18* transcript was cleaved at the *OsmiR160* target site. We generated transgenic rice plants (named *mOsARF18*) that expressed an *OsmiR160*-resistant version of *OsARF18*. Phenotypic analyses revealed that *mOsARF18* plants showed pleiotropic defects in growth and development. Furthermore, auxin treatment induced expression of *OsMIR160a*, *OsMIR160b*, and *OsARF18*, whereas expression of *OsMIR160a* and *OsMIR160b* as well as other genes involved in auxin signaling was altered in *mOsARF18* plants. Our results support the idea that deregulation of the *OsmiR160* target gene *OsARF18* leads to abnormal growth and development in rice through affecting the auxin signaling.

Results

Expression of *OsMIR160* genes and *OsmiR160* target *OsARF* genes in rice. Rice has six *OsMIR160* (*OsMIR160a* to *OsMIR160f*) genes. Four of six mature *OsmiR160s* (*OsmiR160a* to *OsmiR160d*) have identical sequences, while the other two (*OsmiR160e* and *OsmiR160f*) each differs by a single nucleotide (Fig. S1a). Our RT-PCR results showed that expression of *OsMIR160a* and *OsMIR160b* genes was significantly higher than that of the other four *OsMIR160* genes in leaf (L), young inflorescence (YI), mature inflorescence (MI), and stem (ST) (Fig. S1b). Moreover, quantitative real-time RT-PCR (qRT-PCR) revealed that *OsMIR160a* and *OsMIR160b* genes had higher expression levels in leaf and inflorescences than that in seedling and stem (Fig. 1a,b). Our studies suggest that the mature *OsmiR160* might be mainly derived from *OsMIR160a* and *OsMIR160b* genes.

Our analyses using the psRNATarget web server³¹ showed that *OsARF8* (*Os02g41800*), *OsARF10* (*Os04g43910*), *OsARF18* (*Os06g47150*), and *OsARF22* (*Os10g33940*) were potential *OsmiR160* target genes. Further phylogenetic analysis found that all four targets were similar to *Arabidopsis* *ARF10* and *ARF16*, whereas *OsARF18* had the highest similarity to the *Arabidopsis* *ARF16* (Fig. 1g; Fig. S2). Our qRT-PCR results demonstrated that the expression of *OsARF8* mainly occurred in young and mature inflorescences (Fig. 1c), while *OsARF10* was primarily expressed in seedling and stem (Fig. 1d). *OsARF18* was predominantly expressed in leaf as well as young and mature inflorescences (Fig. 1e). Expression levels of *OsARF22* appeared similar in all examined organs except in leaf (Fig. 1f). A previous study using the “degradome sequencing” approach has shown that cleavage frequencies associated with *OsARF18* and *OsARF22* are the highest, but frequencies for *OsARF8* and *OsARF10* are extremely low³². We further identified that the *OsARF18* transcript was cleaved at the *OsmiR160* target site using a gene-specific 5' RACE (Fig. 1h). Therefore, our results suggest that *OsARF18* (*Os06g47150*) is a promising target gene of *OsmiR160*.

Negative regulation of *OsARF18* by *OsmiR160* is essential for rice growth and development.

To investigate the function of *OsmiR160* in rice, we chose its promising target *OsARF18* to generate transgenic plants expressing an *OsmiR160*-resistant version of *OsARF18* (resulting plants named *mOsARF18*) in the *Oryza sativa* Japonica cv. Nipponbare background (Fig. 2a). Our qRT-PCR results show that the expression of un-cleaved *OsARF18* was significantly increased in leaf, stem, and inflorescence of three examined *mOsARF18* transgenic lines (Fig. 2b).

Eleven out of 19 *mOsARF18* independent transgenic lines exhibited strong but similar defects in both growth and development, while the rest of lines showed mild phenotypes. Compared to wild-type plants, *mOsARF18* lines were dwarf and formed less tillers (Fig. 3a,g,h). *mOsARF18* plants produced short and rolled leaves (Fig. 3b; Fig. S3; Fig. 4a–c). *mOsARF18* plants were also defective in reproduction, as indicated by abnormal flower and seed development (Fig. 3c–f). The lemma and palea did not enclose flowers (Fig. 3c). After fertilization, stamens remained attached to developing seeds, suggesting abnormal senescence of stamens (Fig. 3d). Moreover, *mOsARF18* lines showed reduced seed setting (Fig. 3i). Although no change in seed length (Fig. 3e,f,j), *mOsARF18* seeds had reduced width and weight when compared with wild type (Fig. 3e,f,k,l). Our results suggest that deregulation of *OsARF18* results in abnormal growth and development in rice.

Cell division and differentiation were abnormal in *mOsARF18* leaves. To further examine leaf defects in *mOsARF18* plants, we analyzed the structure of the fifth mature leaf via cross-section. Compared with wild type, *mOsARF18* lines produced rolled leaves (Fig. 4a–c). Bulliform cells are specialized epidermal cells between two vascular bundles on the adaxial blade. Bulliform cells are large and highly vacuolated, which are important for leaf rolling through turgor pressure regulation³³. Wild-type bulliform cells were arranged in groups of approximately 5 cells, among which the middle bulliform cell was larger than others along both sides (Fig. 4d,i).




Figure 1. Expression analysis of *OsMIR160a*, *OsMIR160b*, and *OsMIR160* potential target *OsARFs*. (a,b) Quantitative real time RT-PCR (qRT-PCR) results showing expression of *OsMIR160a* (a) and *OsMIR160b* (b). (c–f) qRT-PCR results showing expression of *OsMIR160* potential target genes: *OsARF8* (c), *OsARF10* (d), *OsARF18* (e), and *OsARF22* (f). S: seedling (7-day old), L: leaf, ST: stem, YI: young inflorescence, and MI: mature inflorescence. Gene expression levels in other organs were normalized based on expression observed in seedling. Stars indicate significant difference ($P < 0.01$). (g) An unrooted phylogenetic tree constructed by the Maximum-Likelihood method showing miR160 target ARFs in *Arabidopsis* and rice. (h) The 5' RACE result showing that *OsARF18* is a target of *OsMIR160*. The arrow points to the *OsMIR160*-directed cleavage site at the *OsARF18* transcript.




Figure 2. Generation of transgenic rice plants expressing an *OsmiR160*-resistant version of *OsARF18* (*mOsARF18*). (a) Domain structure of the *OsARF18* protein. The B3 DNA binding domain (B3) and the C-terminal dimerization domain (CTD) are indicated in black and gray, respectively. The *OsmiR160* complementary (or target) sequence in the *OsARF18* mRNA and the corresponding region of amino acid sequence (GIQGAR) are shown. The silent mutations were created in *mOsARF18* by introducing silent mutations (indicated by*). (b) qRT-PCR results showing that expression levels of un-cleaved *OsARF18* were significantly (* $P < 0.01$) increased in leaf, stem, and inflorescence of three examined independent *mOsARF18* transgenic lines.

However, in *mOsARF18* leaves, we observed reduced number and size of bulliform cells, or bulliform cells were absent (Fig. 4e,f). Further statistical analysis showed that bulliform cell numbers between two vascular bundles in *mOsARF18* leaves were significantly reduced when compared with that of wild type (Fig. 4i). Numbers of total vascular bundles and abaxial epidermal cells between two vascular bundles were also decreased (Fig. 4d,g,h,j,k). In addition, with the exception of the bulliform cells, we did not observe significant size differences in other cells when comparing *mOsARF18* leaves with the wild-type. Thus, our results suggest that repression of *OsARF18* by *OsmiR160* is important for epidermal cell division and bulliform cell differentiation during leaf development.

Starch accumulation during seed development was abnormal in *mOsARF18* plants. To study why *mOsARF18* plants produced small seeds, we examined starch accumulation in developing seeds. Our results demonstrated that endosperm in wild-type developing seeds contained many starch granules that were strongly stained by iodine (Fig. 4l), whereas endosperm in *mOsARF18* developing seeds contained smaller, weakly stained starch granules (Fig. 4m,n). In conclusion, *mOsARF18* plants were defective in starch accumulation during seed development.

The *OsmiR160*-regulated *OsARF18* may control rice growth and development via affecting auxin signaling. ARFs play a primary role in auxin signaling. To test whether auxin signaling was affected in *mOsARF18* transgenic plants, we examined expression of representative genes known to be involved in auxin signaling in rice, including the *AUXIN RESPONSIVE FACTOR* gene *OsARF2*³⁴, the auxin responsive gene *OsGH3-1*³⁵, the *AUX1-LIKE* gene *OsLAX2* (*OsAUX3* or *OsRAU2*; Os03g14080)³⁶, the auxin efflux gene *OsPIN1b*³⁷, and the auxin biosynthesis gene *OsYUCCA2*³⁸ via qRT-PCR. Expression levels of all tested genes were significantly decreased in *mOsARF18* transgenic plant leaves (Fig. 5a–e). Thus, our results suggest that deregulation of *OsARF18* affects auxin signaling, which might cause abnormal growth and development in rice.




Figure 3. *mOsARF18* transgenic plants exhibited pleiotropic defects in growth and development. (a) Forty-day old wild-type (left), *mOsARF18-3* (middle), and *mOsARF18-5* (right) plants. Bar: 10 cm. (b) The fifth leaves of wild-type (left), *mOsARF18-3* (middle), and *mOsARF18-5* (right) plants. Bar: 1 mm. (c,d) Ten-DAP (Days After Pollination) wild-type (left), *mOsARF18-3* (middle), and *mOsARF18-5* (right) developing seeds. Bars: (c,d) 1 mm. Lemma and palea were removed in (d) to show attached stamens. (e,f) Wild-type (e) and *mOsARF18-3* (f) mature seeds. Bars: (e,f) 1 mm. (g,h) Sixty-day old transgenic plants showed significantly (* $P < 0.01$) decreased plant height (g) and tiller numbers (h) than that of wild type plants. (i-l) Mature transgenic plant seeds showed no difference in seed length (j), but significantly (* $P < 0.01$) decreased seed setting (i), width (k) and weight (l).




Figure 4. *mOsARF18* plants were abnormal in leaf cell division and differentiation as well as starch accumulation in seeds. (a–c) Hand cutting of the fifth leaves of wild-type (a), *mOsARF18-3* (b), and *mOsARF18-5* (c) plants. Bars: (a) 500 μm ; (b,c) 200 μm . (d–h) Semi-thin sections of the fifth leaves of wild-type, *mOsARF18-3*, and *mOsARF18-5* plants: (d) wild type; (e,g) *mOsARF18-3*; and (f,h) *mOsARF18-5*. Arrows indicate bulliform cells. Arrowheads indicate the lack of typical bulliform cells. Red brackets indicate regions between two vascular bundles for epidermal cell counting. Bars: (d–h) 50 μm . (i–k) Numbers of bulliform cells (i), total vascular bundles (j), and abaxial epidermal cells between two vascular bundles (k) were significantly ($*P < 0.01$) decreased in leaves of *mOsARF18-3*, *mOsARF18-5*, and *mOsARF18-8* transgenic lines in comparison with wild-type plants. (l) Endosperm in 10-DAP (Days After Pollination) wild-type developing seeds showing strongly stained starch granules by I-KI (iodine-potassium iodide). (m,n) Endosperm in 10-DAP *mOsARF18-3* (b) and *mOsARF18-5* (c) developing seeds showing small and weakly stained starch granules. Bars: (l–n) 20 μm .

We further tested whether auxin affected expression of *OsMIR160a*, *OsMIR160b*, and *OsmiR160* target genes. Our qRT-PCR results showed that expression of *OsMIR160a* was significantly increased after 20 and 40-minute NAA treatment (Fig. 6a), while expression of *OsMIR160b* was significantly induced after 20-minute NAA treatment and then remained normal (Fig. 6b). NAA treatment also significantly induced expression of *OsARF18* after 20 and 40 minutes (Fig. 6e) as well as expression of *OsARF10* after 180 minutes (Fig. 6d). However, expression of *OsARF8* (Fig. 6c) and *OsARF22* (Fig. 6f) remained unchanged after NAA treatment.

Conversely, expression of *OsMIR160a* and *OsMIR160b* was found to be decreased in *mOsARF18* plants (Fig. 6g,h), in which the expression of *OsARF18* was highly increased (Fig. 2b). Our results suggest that auxin up regulates expression of *OsMIR160a*, *OsMIR160b*, and *OsARF18*, whereas *OsARF18* represses expression of




Figure 5. Analysis of auxin signaling gene expression in *mOsARF18* transgenic plants. qRT-PCR results showing that expression of *OsARF2* (a), *OsGH3-1* (b), *OsLAX2* (c), *OsPIN1b* (d), and *OsYUCCA2* (e) was significantly decreased in leaves of *mOsARF18* transgenic plants. Gene expression levels in *mOsARF18-3*, *mOsARF18-5*, and *mOsARF18-8* were normalized based on expression observed in wild type. Stars indicate significant difference ($P < 0.01$).

OsMIR160a and *OsMIR160b*. Therefore, the positive regulation of *OsMIR160a* and *OsMIR160b* expression by auxin and the negative regulation of *OsMIR160a* and *OsMIR160b* by *OsARF18* might be important for *OsmiR160* to fine-tune auxin signaling in a negative feedback loop manner.

To test whether *OsmARF18* seedlings are defective in auxin signaling, we treated *OsmARF18* seedlings hydroponically with 1 μ M of NAA for 7 days. We found that *OsmARF18* seedlings produced significantly shorter primary roots than that of wild type (Fig. 7a,b,e). Primary root growth was significantly inhibited by NAA treatment in both wild type and *OsmARF18* seedlings (Fig. 7c–e); however, the NAA inhibition effect on primary




Figure 6. Auxin up regulated expression of *OsMIR160a*, *OsMIR160b*, and *OsMIR160* target genes, whereas *OsARF18* suppressed the expression of *OsMIR160a*. (a–f) qRT-PCR results showing that expression changes of *OsMIR160a*, *OsMIR160b*, and *OsMIR160* target genes in seedlings (7-day old) treated with NAA for 20, 40, and 180 minutes. Expression levels without NAA treatment (0 minute) were used to normalize expression with treatments. Stars indicate significant difference ($P < 0.01$). Expression of *OsMIR160a* was significantly induced after 20 and 40-minute treatment (a), while expression of *OsMIR160b* was significantly increased only after 20-minute treatment (b). Expression of *OsARF8* (c) and *OsARF22* (f) remained unchanged. Expression of *OsARF10* was significantly induced after 180-minute treatment (d), whereas expression of *OsARF18* was significantly increased after 20 and 40-minute treatment (e). (g,h) qRT-PCR results showing that expression of *OsMIR160a* and *OsMIR160b* was significantly decreased in *mOsARF18* transgenic lines in comparison with wild-type plants. Gene expression levels in #3 (*mOsARF18-3*), #5 (*mOsARF18-5*), and #8 (*mOsARF18-8*) were normalized based on expression observed in wild type. Stars indicate significant difference ($P < 0.01$).




Figure 7. Effect of NAA treatment on lateral root formation. (a,c) Seven-day wild type (WT) seedlings without NAA (a) and with NAA (1 μ M) (c) treatment (inset showing high magnification of roots in (c)). (b,d) Seven-day *OsmARF18* seedlings without NAA (b) and with NAA (1 μ M) (d) treatment (inset showing high magnification of roots in (d)). (e,f) Primary roots in WT 7-day seedlings were significantly (*indicating $P < 0.01$) longer than that of *OsmARF18* seedlings. NAA treatment significantly ($P < 0.01$) inhibited primary root length of both WT and *OsmARF18* seedlings, but the relative primary root length (percentage of root length between that treated and untreated) was similar. (g) Seven-day WT and *OsmARF18* seedlings produced similar numbers of lateral roots. After NAA treatment, lateral root numbers in WT seedlings was significantly ($P < 0.01$) increased, whereas that in *OsmARF18* seedlings was significantly ($P < 0.01$) decreased.

root length was similar in wild type and *OsmARF18* seedlings, because there was no change in relative primary root length (percentage of root length between that treated and untreated) after the NAA treatment (Fig. 7f). Numbers of lateral roots were similar in wild type and *OsmARF18* seedlings (Fig. 7a,b,g). NAA treatment significantly increased the lateral root number in wild type seedlings (Fig. 7a,c,g), conversely, the lateral root number in *OsmARF18* seedlings was significantly decreased with the NAA treatment (Fig. 7b,d,g). Our results suggest that *OsmARF18* might be involved in auxin-regulated lateral root formation in rice.

Discussion

Conserved and diverse roles of miR160 in plant growth and development. MiR160 is conserved throughout the plant kingdom from mosses to higher plants^{29,30}. Sequence similarity of mature miR160s is more than 80% among different species. MiR160 target genes are also conserved, although their numbers vary with species. *Arabidopsis* contains *MIR160a*, *MIR160b*, and *MIR160c* three genes which produce the same mature miR160⁶. MiR160 targets *AUXIN RESPONSE FACTOR 10* (*ARF10*), *ARF16*, and *ARF17* three genes and each target gene has conserved but some distinct functions in *Arabidopsis*. Plants expressing the miR160-resistant version of *ARF10* (*mARF10*) exhibit pleiotropic defects that resemble phenotypes of some ABA and auxin defective mutants²⁴, while *mARF16* plants have reduced fertility and less lateral roots²². *ARF10* and *ARF16* are required for maintaining the expression of *ABI3* gene²⁵, suggesting that *ARF10* and *ARF16* are involved in both auxin and ABA signaling or cross-talk between them. *ARF17* plays a general role in both vegetative and reproductive development via modulating expression of early auxin response genes²³. *SlymiR160* and the *SlymiR160a* target *SlyARF10* control ovary patterning, early fruit development and floral organ abscission in tomato^{26,27}, whereas the soybean miR160 regulates auxin and cytokinin signaling during nodulation²⁸.

Six *OsmIR160* (*OsmIR160a* to *OsmIR160f*) genes are found in rice. The mature *OsmiR160* has *OsmARF8* (*Os02g41800*), *OsmARF10* (*Os04g43910*), *OsmARF18* (*Os06g47150*), and *OsmARF22* (*Os10g33940*) four potential target genes. In this study, we characterized the function of *OsmARF18* (*Os06g47150*), an orthologue of *Arabidopsis* *ARF16*. *Arabidopsis mARF16* and rice *mOsARF18* plants show some similar but also different defects in growth and reproductive development. In *Arabidopsis*, *mARF16* plants produce curved leaves²². Our studies showed that the formation of rolled leaves in *mOsARF18* plants was caused by alterations in shape, size, and number of bulliform cells. Auxin plays a key role in controlling leaf shape³⁹. Phenotypic analysis indicates that alteration of bulliform cells on the adaxial leaf blade surface is tightly linked to the formation of rolled leaves in rice. Disruption of the rice *CONSTITUTIVELY WILTED1* (*OsCOW1*) gene, a member of *OsyUCCA* gene family, results in rupture of the largest bulliform cell and consequently rolled leaves⁴⁰. In addition, alterations in number and size of bulliform cells lead to rolled leaves in the *narrow leaf 7* (*nal7*) mutant, which is an *oscow1* allele⁴¹. Therefore, our data support that the *OsmiR160* target *OsmARF18* controls leaf shape via affecting auxin signaling. Besides rolled leaves, *mOsARF18* plants produced less and smaller seeds than wild type. Moreover, starch accumulation in seeds from *mOsARF18* plants was significantly decreased when compared to wild-type seeds. Collectively, miR160 plays conserved and diverse roles in plant growth and development. It will be interesting to test the loss-of-function of *OsmiR160* and functions of other *OsmiR160* target genes in rice growth and development.

Negative regulation of *OsmARF18* by *OsmiR160* is important for its normal function. MiRNAs negatively regulate gene expression at the post-transcriptional level by binding to mRNA complementary sequences for mRNA destabilization and translational inhibition in both plants and animals^{1–5}. One miRNA normally has multiple target genes. Over/ectopic-expression of normal miRNA-target genes usually does not cause a change in phenotype, because over/ectopic-expressed normal mRNAs can be still targeted by miRNAs for cleavage. Therefore, a primary approach for studying the function of miRNA and its target is to express the miRNA-resistant version of individual target gene. Employing 35S, *Ubi*, and *ACTIN* promoters, previous studies identified functions of many miRNAs and their target genes via expressing miRNA-resistant versions of target genes, such as 35S::*mTCP2*, 35S::*mTCP3*, 35S::*mTCP4*, 35S::*mCUP1*, and 35S::*mCUP2* (*Arabidopsis* miR164 target genes)^{42,43}, 35S::*mAP2* (*Arabidopsis* miR172 target gene)⁴⁴, 35S::*mSlyARF10* (tomato miR160 target gene)²⁶, *UBI:mGRF6* (rice miR396 target gene)⁴⁵, as well as *ACTIN::mOSHB1*, *ACTIN::mOSHB3* and *ACTIN::mOSHB5* (rice miR166 target genes)⁴⁶.

So far, no studies report that over/ectopic-expression of normal versions of miR160 target genes (*ARF10*, *ARF16*, and *ARF17*) causes phenotypes in leaf and flower development, but only plants expressing miR160-resistant versions of miR160 target genes (*mARF10*, *mARF16*, and *mARF17*) exhibit various phenotypes^{22–24,26}. In this study, we used the *Ubiquitin* (*Ubi*) promoter to drive *mOsARF18* expression. To further rule out the possibility that the phenotype of *mOsARF18* was caused by ectopic activity of the *Ubi* promoter, we generated *ARF16*, *mARF16*, *OsmARF18*, and *mOsARF18* *Arabidopsis* transgenic plants under the control of the *Ubi* promoter. Of 30 *ARF16* and 32 *OsmARF18* transgenic plants that we obtained, none of them showed detectable defects in growth and development (Fig. S5a–c,f–h; Fig. S6a–c,f–h,k–m,p); however, *mARF16* (32 out of 40) and *mOsARF18* (20 out of 29) transgenic plants formed narrow and curled leaves, and had short stature (Fig. S5d,e,i,j). In addition, *mARF16* and *mOsARF18* plants produced abnormal flowers and a lower number of smaller seeds compared with wild-type plants (Fig. S6d,e,i,j,n,o,q). Our results showed that *mARF16* and *mOsARF18* caused defects in plant growth and development, but plants ectopically expressing normal *ARF16* and *OsmARF18* were similar to the wild type. Taken together, the results from *Arabidopsis* and rice suggest that the negative regulation of *OsmARF18* and its *Arabidopsis* orthologous gene *ARF16* by rice and *Arabidopsis* miR160, respectively, is critical for their function.

Molecular mechanism of miR160 in fine-tuning auxin signaling. MiRNAs play a pivotal role in auxin signaling by negative regulation of *ARFs*. In *Arabidopsis*, miRNAs are involved in expression regulation of 8 of a total 23 *ARFs* (*ARF2*, 3, 4, 6, 8, 10, 16, and 17). MiR390-derived trans-acting-small interfering RNAs




Figure 8. A hypothetical working model for auxin signaling that is modulated by *OsmiR160* during rice growth and development. Auxin promotes the release of *OsARF* activators or repressors via the auxin receptor SCF E3 ligase. The balance between *OsARF* activators and repressors decides up or down expression of *OsMIR160* genes. Conversely, the change in abundance of mature *OsmiR160* negatively regulates expression of its target *OsARFs*. *OsARF* activators and repressors might positively or negatively regulate expression of each other.

(ta-siRNAs) target *ARF2*, *ARF3*, and *ARF4*^{47,48}. The tasiRNA gradient is important for establishing the normal patterning of *ARF3* protein during leaf development⁴⁹. During growth of lateral roots, miR390 affects production of tasiRNAs, and thus inhibits *ARF2*, *ARF3*, and *ARF4*⁵⁰. Conversely, auxin activates *ARF2*, *ARF3*, and *ARF4*, which consequently influences the formation of miR390. Therefore, the regulatory network modulated by miR390 maintains normal expression of *ARF2*, *ARF3*, and *ARF4*. Negative regulation of *ARF6* and *ARF8* by miR167 is essential for anther and ovule development⁵¹. Moreover, miR160 negatively regulates expression of *ARF10*, *ARF16*, and *ARF17*^{6,22–24}. In rice, *OsmiRNAs* are predicted to regulate 11 of a total of 25 identified *OsARFs*^{52,53}. Functional disruption of *OsDCL4* causes increased expression of *OsmiR165/166* and three *OsARFs*, which are orthologs of *Arabidopsis* *ARF2*, *ARF3*, and *ARF4*⁵⁴. *OsmiR160* is predicted to have four *OsARF* targets, including *OsARF18* that was characterized in this study. *OsmiR167* potentially targets *OsARF6* (*Os02g06910*), *OsARF12* (*Os04g57610*), *OsARF17* (*Os06g46410*), and *OsARF25* (*Os12g41950*), which are highly similar to *Arabidopsis* *ARF6* and *ARF8*⁵⁴. Together, miRNAs target a group of similar *ARF* genes in both monocots and dicots.

Auxin promotes the SCF^{TIR1/AFB} E3 ligase-mediated degradation of Aux/IAA proteins, which sequester ARFs^{55,56}. Upon the perception of auxin, the released ARFs activate or suppress expression of a large set of auxin-responsive genes by binding to auxin response elements (AuxREs) in their regulatory regions. Based on transient expression studies and protein structures, among 23 ARFs in *Arabidopsis*, ARF5–8, and 19 act as activators, while ARF1–4, 9–18, and 20–22 may function as repressors^{57–59}.

Previous studies have implied that the feedback regulation between miRNA and its target *ARF* genes could provide a fine-tuning mechanism to regulate auxin signaling. During adventitious root development in *Arabidopsis*, expression of *ARF6* and *ARF8* is regulated by miR167 and miR160, whereas the abundance of *ARF17* transcripts is controlled by miR160⁶⁰. In addition, *ARF6* and *ARF8* activators as well as *ARF17* repressors positively and negatively regulate the expression of each other. Thus, the delicate balance between miRNAs and *ARFs* is critical for auxin-regulated developmental processes. Our studies show that *OsmiR160* negatively regulates the expression of *OsARF18* by cleaving *OsARF18* transcripts. Auxin induces expression of *OsMIR160a*, *OsMIR160b*, and *OsARF18*, whereas expression of *OsMIR160a* and *OsMIR160b* was suppressed by *OsARF18*. Our analyses revealed that promoters and 3' regions of *OsMIR160a* and *OsMIR160b* had clusters of AuxRE cores and AuxRR cis elements (Fig. S4), which may create *OsARF* binding sites by which *OsARF* activators or repressors could regulate expression of *OsMIR160a* and *OsMIR160b*.

In our hypothetical model, auxin promotes the release of *OsARF* activators and repressors via the SCF^{TIR1/AFB} E3 ligase (Fig. 8). The balance between *OsARF* activators and repressors decides up or down expression of *OsMIR160* genes, which consequently affects the abundance of mature *OsmiR160*. Conversely, *OsmiR160* negatively modulates expression of its target *OsARFs*. *OsARFs* also mutually control their own expression. Future studies should focus on examining the feedback loop regulation between *OsmiR160* and its target *OsARFs*, which might be important for fine-tuning highly dose-sensitive auxin signaling during rice growth and development.

Manipulating miRNAs and their target genes has demonstrated improvement of many important crop traits, including biomass yield, grain yield, fruit yield, nutritional quality, abiotic stress resistance (e.g. drought, salinity, cold, heat, oxidative stress, nutrient deprivation tolerance, and heavy metal detoxification), and biotic stress resistance (e.g. virus, bacteria, fungus, and nematode resistance)^{14,15,61,62}. Our results showed that *OsmiR160* played a pivotal role in rice growth and development via regulating auxin signaling. In particular, *OsmiR160* is essential for leaf and seed development in rice. Leaf shape is important for photosynthesis, respiration, and transpiration. Moderate leaf rolling can enhance photosynthesis and stress responses by inhibiting water loss and radiant heat absorption, which, therefore, increases crop yield. Future studies on the molecular mechanism by which *OsmiR160* modulates auxin signaling will lead to potential applications for improving crop agronomic traits.

Methods

Plant materials and growth condition. Rice (*Oryza sativa* L. Japonica Nipponbare) plants were grown in Metro-Mix 360 soil (Sun-Gro Horticulture, Agawam, MA, USA) supplemented with sand and iron in a walk-in growth chamber under a 12-hour light (28 °C)/12-hour dark (22 °C) photoperiod regime. Transgenic rice plants were generated at the Plant Transformation Facility at Iowa State University. In total, 124 *mOsARF18* plants were obtained from 19 independent transgenic lines. For expression studies, seven-day old wild-type seedlings were treated with 1 μ M of NAA for 20, 40, and 180 min with untreated 7-day old seedlings as control. For studying the effect of exogenous auxin on root development, germinated seeds were hydroponically grown in the 1/2 Kimura B nutrient solution (pH 5.6) containing 1 μ M of NAA for 7 days with continuous light at 25 °C.

Vector construction and rice transformation. PCR reactions (Primers are shown in Table S1) were performed using the Phusion High-Fidelity DNA Polymerase (New England Biolabs, Ipswich, MA, USA). The *OsARF18* cDNA was amplified from rice leaf cDNAs and then cloned into the pCR2.1 vector (Invitrogen, Grand Island, NY, USA), resulting in *pCR2.1-OsARF18*. Point mutations of *OsARF18* were created by overlapping PCR to generate *pCR2.1-mOsARF18*. *mOsARF18* was then subcloned into the modified pCambia1301 binary vector harboring the Gateway cassette sequence and the maize *Ubiquitin (Ubi)* promoter using the Gateway LR recombinase II enzyme mix (Invitrogen, Grand Island, NY, USA).

For rice transformation, the *mOsARF18* construct was introduced into the *Agrobacterium* strain EHA101. The callus induction (from mature embryos of Japonica cv. Nipponbare seeds), *Agrobacterium* infection, co-cultivation, selection of transformed calli, and plant regeneration were performed essentially as described previously⁶³.

RT-PCR and real time qRT-PCR. Total RNAs were isolated from different rice tissues using the RNeasy Plant Mini Kit (Qiagen, Valencia, CA, USA). After determining the RNA quantification by the NanoDrop 2000c (Thermo Scientific, Bannockburn, IL, USA), RNA reverse transcription was conducted using the QuantiTect Reverse Transcription Kit (Qiagen, Valencia, CA, USA). Real-time PCR (DNA Engine Opticon 2 system, Hercules, California, USA) and data analysis were performed as previously described⁶. Expression of *OsMIR160a* through *OsMIR160f* was examined by RT-PCR. Expression of *OsMIR160a*, *OsMIR160b*, *OsARF8*, *OsARF10*, *OsARF18*, and *OsARF22*, as well as other auxin signaling related genes was tested by real-time qRT-PCR. Three biological repeats were conducted and each value indicates the average with the standard error. All primer sequences are shown in Table S1.

5' RACE. Using the GeneRacer™ kit (Invitrogen, Grand Island, NY, USA), a gene-specific 5'-rapid amplification of cDNA ends was conducted as described previously²³. Gene-specific primers for *OsARF18* (Os06g47150) are shown in Table S1.

Semi-thin section analysis. Semi-thin sectioning was performed as described previously^{64,65}. The fifth leaves of 6-week-old rice were fixed in 2.5% of glutaraldehyde and post-fixed with 1% of OsO₄ at room temperature. Samples were dehydrated through a graded acetone series (10% increments) for 60 minutes each. Infiltrated was started with 20% of Spurr's resin and then 40%, 60%, and 80% of Spurr's resin every 3 hours. Following infiltration in three changes of 100% Spurr's resin for 24 hours each, samples were finally embedded in 100% Spurr's resin and polymerized at 60 °C overnight. Semi-thin sections (0.5 μ m) were made using an RMC MT-7 ultramicrotome (Reichert-Jung, Depew, NY, USA) and were stained with 0.25% Toluidine Blue O. Images were photographed by an Olympus BX51 microscope equipped with an Olympus DP 70 digital camera (Olympus, Center Valley, PA, USA).

Histological detection of starch. Ten-DAP (Days After Pollination) seeds were fixed in FAA (50% ethanol, 10% formalin, 5% acetic acid). Following fixation, samples were dehydrated through an ethanol series, embedded in paraffin, and sectioned at 8 μ m with a Spencer 820 microtome. Sections were dewaxed and stained with Lugol's iodine solution (6 mM iodine, 43 mM KI, and 0.2 N HCl) for detection of starch granules. Images were photographed with an Olympus BX51 microscope equipped with an Olympus DP 70 digital camera (Olympus, Center Valley, PA, USA).

References

- Bartel, D. P. MicroRNAs: target recognition and regulatory functions. *Cell* **136**, 215–233 (2009).
- Carrington, J. C. & Ambros, V. Role of microRNAs in plant and animal development. *Science* **301**, 336–338 (2003).
- Jones-Rhoades, M. W., Bartel, D. P. & Bartel, B. MicroRNAs and their regulatory roles in plants. *Annu. Rev. Plant Biol.* **57**, 19–53 (2006).
- Voinnet, O. Origin, biogenesis, and activity of plant microRNAs. *Cell* **136**, 669–687 (2009).
- Rogers, K. & Chen, X. Biogenesis, turnover, and mode of action of plant microRNAs. *Plant Cell* **25**, 2383–2399 (2013).
- Liu, X. *et al.* The role of floral organs in carpels, an *Arabidopsis* loss-of-function mutation in *MicroRNA160a*, in organogenesis and the mechanism regulating its expression. *Plant J.* **62**, 416–428 (2010).
- Todesco, M., Rubio-Somoza, I., Paz-Ares, J. & Weigel, D. A collection of target mimics for comprehensive analysis of microRNA function in *Arabidopsis thaliana*. *Plos Genet.* **6**, e1001031 (2010).
- Yan, J. *et al.* Effective small RNA destruction by the expression of a short tandem target mimic in *Arabidopsis*. *Plant Cell* **24**, 415–427 (2012).
- Liu, B. *et al.* Loss of function of *OsDCL1* affects microRNA accumulation and causes developmental defects in rice. *Plant Physiol.* **139**, 296–305 (2005).
- Song, X. *et al.* Roles of *DCL4* and *DCL3b* in rice phased small RNA biogenesis. *Plant J.* **69**, 462–474 (2012).
- Liu, B. *et al.* *Oryza sativa dicer-like4* reveals a key role for small interfering RNA silencing in plant development. *Plant Cell* **19**, 2705–2718 (2007).
- Nonomura, K. I. *et al.* A germ cell-specific gene of the ARGONAUTE family is essential for the progression of premeiotic mitosis and meiosis during sporogenesis in rice. *Plant Cell* **19**, 2583–2594 (2007).

13. Abe, M. *et al.* WAVY LEAF1, an ortholog of *Arabidopsis* HEN1, regulates shoot development by maintaining MicroRNA and trans-acting small interfering RNA accumulation in rice. *Plant Physiol.* **154**, 1335–1346 (2010).
14. Jiao, Y. *et al.* Regulation of OsSPL14 by OsmiR156 defines ideal plant architecture in rice. *Nature Genet.* **42**, 541–544 (2010).
15. Wang, S. *et al.* Control of grain size, shape and quality by OsSPL16 in rice. *Nature Genet.* **44**, 950–954 (2012).
16. Xie, K. *et al.* Gradual increase of miR156 regulates temporal expression changes of numerous genes during leaf development in rice. *Plant Physiol.* **158**, 1382–1394 (2012).
17. Zhu, Q. H., Upadhyaya, N. M., Gubler, F. & Helliwell, C. A. Over-expression of miR172 causes loss of spikelet determinacy and floral organ abnormalities in rice (*Oryza sativa*). *BMC Plant Biol.* **9**, 149 (2009).
18. Lee, D. Y. & An, G. Two AP2 family genes, *supernumerary bract* (SNB) and *Osindeterminate spikelet 1* (OsIDS1), synergistically control inflorescence architecture and floral meristem establishment in rice. *Plant J* **69**, 445–461 (2012).
19. Bian, H. *et al.* Distinctive expression patterns and roles of the miRNA393/TIR1 homolog module in regulating flag leaf inclination and primary and crown root growth in rice (*Oryza sativa*). *New Phytol.* **196**, 149–161 (2012).
20. Sunkar, R., Girke, T., Jain, P. K. & Zhu, J. K. Cloning and characterization of microRNAs from rice. *Plant Cell* **17**, 1397–1411 (2005).
21. Jeong, D. H. *et al.* Massive analysis of rice small RNAs: mechanistic implications of regulated microRNAs and variants for differential target RNA cleavage. *Plant Cell* **23**, 4185–4207 (2011).
22. Wang, J. W. *et al.* Control of root cap formation by MicroRNA-targeted auxin response factors in *Arabidopsis*. *Plant Cell* **17**, 2204–2216 (2005).
23. Mallory, A. C., Bartel, D. P. & Bartel, B. MicroRNA-directed regulation of *Arabidopsis* AUXIN RESPONSE FACTOR17 is essential for proper development and modulates expression of early auxin response genes. *Plant Cell* **17**, 1360–1375 (2005).
24. Liu, P. P. *et al.* Repression of AUXIN RESPONSE FACTOR10 by microRNA160 is critical for seed germination and post-germination stages. *Plant J.* **52**, 133–146 (2007).
25. Liu, X. *et al.* Auxin controls seed dormancy through stimulation of abscisic acid signaling by inducing ARF-mediated ABI3 activation in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **110**, 15485–15490 (2013).
26. Hendelman, A., Buxdorf, K., Stav, R., Kravchik, M. & Arazi, T. Inhibition of lamina outgrowth following *Solanum lycopersicum* AUXIN RESPONSE FACTOR 10 (SlARF10) derepression. *Plant Mol. Biol.* **78**, 561–576 (2012).
27. Damodharan, S., Zhao, D. & Arazi, T. A common miRNA160-based mechanism regulates ovary patterning, floral organ abscission and lamina outgrowth in tomato. *Plant J.* doi: 10.1111/tpj.13127 (2016).
28. Turner, M. *et al.* Ectopic expression of miR160 results in auxin hypersensitivity, cytokinin hyposensitivity, and inhibition of symbiotic nodule development in soybean. *Plant Physiol.* **162**, 2042–2055 (2013).
29. Axtell, M. J., Snyder, J. A. & Bartel, D. P. Common functions for diverse small RNAs of land plants. *Plant Cell* **19**, 1750–1769 (2007).
30. Luo, Y., Guo, Z. & Li, L. Evolutionary conservation of microRNA regulatory programs in plant flower development. *Dev. Biol.* **380**, 133–144 (2013).
31. Dai, X. & Zhao, P. X. psRNATarget: a plant small RNA target analysis server. *Nucleic Acids. Res.* **39**, W155–159 (2011).
32. Li, Y. F. *et al.* Transcriptome-wide identification of microRNA targets in rice. *Plant J.* **62**, 742–759 (2010).
33. Zhang, G. H., Xu, Q., Zhu, X. D., Qian, Q. & Xue, H. W. SHALLOT-LIKE1 is a KANADI transcription factor that modulates rice leaf rolling by regulating leaf abaxial cell development. *Plant Cell* **21**, 719–735 (2009).
34. Wang, D. *et al.* Genome-wide analysis of the auxin response factors (ARF) gene family in rice (*Oryza sativa*). *Gene* **394**, 13–24 (2007).
35. Zhao, S. Q., Xiang, J. J. & Xue, H. W. Studies on the rice LEAF INCLINATION1 (LC1), an IAA-amido synthetase, reveal the effects of auxin in leaf inclination control. *Mol. Plant* **6**, 174–187 (2013).
36. Chhun, T. *et al.* Saturated humidity accelerates lateral root development in rice (*Oryza sativa* L.) seedlings by increasing phloem-based auxin transport. *J. Exp. Bot.* **58**, 1695–1704 (2007).
37. Li, G. *et al.* Rice actin-binding protein RMD is a key link in the auxin-actin regulatory loop that controls cell growth. *Proc. Natl. Acad. Sci. USA* **111**, 10377–10382 (2014).
38. Yamamoto, Y., Kamiya, N., Morinaka, Y., Matsuoka, M. & Sazuka, T. Auxin biosynthesis by the YUCCA genes in rice. *Plant Physiol.* **143**, 1362–1371 (2007).
39. Vanneste, S. & Friml, J. Auxin: a trigger for change in plant development. *Cell* **136**, 1005–1016 (2009).
40. Woo, Y. M. *et al.* Constitutively wilted 1, a member of the rice YUCCA gene family, is required for maintaining water homeostasis and an appropriate root to shoot ratio. *Plant Mol. Biol.* **65**, 125–136 (2007).
41. Fujino, K. *et al.* NARROW LEAF 7 controls leaf shape mediated by auxin in rice. *Mol. Genet. Genomics* **279**, 499–507 (2008).
42. Baker, C. C., Sieber, P., Wellmer, F. & Meyerowitz, E. M. The *early extra petals1* mutant uncovers a role for MicroRNA miR164c in regulating petal number in *Arabidopsis*. *Curr. Biol.* **15**, 303–315 (2005).
43. Koyama, T., Furutani, M., Tasaka, M. & Ohme-Takagi, M. TCP transcription factors control the morphology of shoot lateral organs via negative regulation of the expression of boundary-specific genes in *Arabidopsis*. *Plant Cell* **19**, 473–484 (2007).
44. Chen, X. A microRNA as a translational repressor of APETALA2 in *Arabidopsis* flower development. *Science* **303**, 2022–2025 (2004).
45. Liu, H. *et al.* OsmiR396d-regulated OsGRFs function in floral organogenesis in rice through binding to their targets OsJM706 and OsCR4. *Plant Physiol.* **165**, 160–174 (2014).
46. Itoh, J., Hibara, K., Sato, Y. & Nagato, Y. Developmental role and auxin responsiveness of Class III homeodomain leucine zipper gene family members in rice. *Plant Physiol.* **147**, 1960–1975 (2008).
47. Williams, L., Carles, C. C., Osmont, K. S. & Fletcher, J. C. A database analysis method identifies an endogenous trans-acting short-interfering RNA that targets the *Arabidopsis* ARF2, ARF3, and ARF4 genes. *Proc. Natl. Acad. Sci. USA* **102**, 9703–9708 (2005).
48. Fahlgren, N. *et al.* Regulation of AUXIN RESPONSE FACTOR3 by TAS3 ta-siRNA affects developmental timing and patterning in *Arabidopsis*. *Curr. Biol.* **16**, 939–944 (2006).
49. Chitwood, D. H. *et al.* Pattern formation via small RNA mobility. *Genes Dev.* **23**, 549–554 (2009).
50. Marin, E. *et al.* miR390, *Arabidopsis* TAS3 tasiRNAs, and their AUXIN RESPONSE FACTOR targets define an autoregulatory network quantitatively regulating lateral root growth. *Plant Cell* **22**, 1104–1117 (2010).
51. Wu, M. F., Tian, Q. & Reed, J. W. *Arabidopsis* microRNA167 controls patterns of ARF6 and ARF8 expression, and regulates both female and male reproduction. *Development* **133**, 4211–4218 (2006).
52. Guilfoyle, T. J. & Hagen, G. Auxin response factors. *Curr. Opin. Plant Biol.* **10**, 453–460 (2007).
53. Shen, C. *et al.* Functional analysis of the structural domain of ARF proteins in rice (*Oryza sativa* L.). *J. Exp. Bot.* **61**, 3971–3981 (2010).
54. Nagasaki, H. *et al.* The small interfering RNA production pathway is required for shoot meristem initiation in rice. *Proc. Natl. Acad. Sci. USA* **104**, 14867–14871 (2007).
55. Dharmasiri, N., Dharmasiri, S. & Estelle, M. The F-box protein TIR1 is an auxin receptor. *Nature* **435**, 441–445 (2005).
56. Tan, X. *et al.* Mechanism of auxin perception by the TIR1 ubiquitin ligase. *Nature* **446**, 640–645 (2007).
57. Ulmasov, T., Hagen, G. & Guilfoyle, T. J. Activation and repression of transcription by auxin-response factors. *Proc. Natl. Acad. Sci. USA* **96**, 5844–5849 (1999).
58. Wang, S., Tiwari, S. B., Hagen, G. & Guilfoyle, T. J. AUXIN RESPONSE FACTOR7 restores the expression of auxin-responsive genes in mutant *Arabidopsis* leaf mesophyll protoplasts. *Plant Cell* **17**, 1979–1993 (2005).

59. Boer, D. R. *et al.* Structural basis for DNA binding specificity by the auxin-dependent ARF transcription factors. *Cell* **156**, 577–589 (2014).
60. Gutierrez, L. *et al.* Phenotypic plasticity of adventitious rooting in *Arabidopsis* is controlled by complex regulation of AUXIN RESPONSE FACTOR transcripts and microRNA abundance. *Plant Cell* **21**, 3119–3132 (2009).
61. Zhang, B. MicroRNA: a new target for improving plant tolerance to abiotic stress. *J. Exp. Bot.* **66**, 1749–1761 (2015).
62. Zhou, M. & Luo, H. MicroRNA-mediated gene regulation: potential applications for plant genetic engineering. *Plant Mol. Biol.* **83**, 59–75 (2013).
63. Hiei, Y., Ohta, S., Komari, T. & Kumashiro, T. Efficient transformation of rice (*Oryza sativa* L.) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. *Plant J.* **6**, 271–282 (1994).
64. Jia, G., Liu, X., Owen, H. A. & Zhao, D. Signaling of cell fate determination by the TPD1 small protein and EMS1 receptor kinase. *Proc. Natl. Acad. Sci. USA* **105**, 2220–2225 (2008).
65. Zhao, D. Z., Wang, G. F., Speal, B. & Ma, H. The *EXCESS MICROSPOROCYTES1* gene encodes a putative leucine-rich repeat receptor protein kinase that controls somatic and reproductive cell fates in the *Arabidopsis* anther. *Genes Dev.* **16**, 2021–2031 (2002).

Acknowledgements

We thank X. Li for his early involvement in this project, H.A. Owen, A. Smith, S. Malik, and J. Skurski for critical reading of the manuscript. We are grateful for K. Wang, whose lab performed rice transformation. We also thank Z. Yang for generously offering us the binary vector for rice transformation, H.A. Owen for the phenotypic analysis, and T. Schuck and J. Gonnering for the plant care. This work was in part supported by the National Science Foundation (NSF IOS-0721192 and IOS-1322796), the Research Growth Initiative (RGI) at the University of Wisconsin-Milwaukee, and the UW-Madison/UW-Milwaukee Intercampus Research Incentive Grants Program. D. Zhao also thankfully acknowledges the support of the Shaw Scientist Award from the Greater Milwaukee Foundation.

Author Contributions

D.Z. and H.J. conceived and designed experiments. H.J. and Z.L. conducted experiments. D.Z. and H.J. wrote the paper.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Huang, J. *et al.* Deregulation of the OsmiR160 Target Gene *OsARF18* Causes Growth and Developmental Defects with an Alteration of Auxin Signaling in Rice. *Sci. Rep.* **6**, 29938; doi: 10.1038/srep29938 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>