

MEETING ABSTRACT

Open Access

Functional evaluation of GUCY1A3 mutations associated with myocardial infarction risk

Jana Wobst^{1*}, Tan An Dang¹, Thorsten Kessler¹, Simon von Ameln¹, Stephanie Tennstedt², Christian Hengstenberg^{1,3}, Jeanette Erdmann^{2,4}, Heribert Schunkert^{1,3}

From 7th International Conference on cGMP Generators, Effectors and Therapeutic Implications Trier, Germany. 19-21 June 2015

Myocardial infarction (MI) is the main complication of coronary artery disease (CAD). Recently, a locus tagging the *GUCY1A3* gene has been shown to be genome-wide significantly associated with CAD [1]. *GUCY1A3* encodes for the α_1 -subunit of the soluble guanylyl cyclase (sGC) which consists of α_1 - and β_1 -subunits and catalyzes the production of cGMP upon stimulation with nitric oxide (NO). cGMP acts a second messenger that mediates diverse cellular functions, e.g. smooth muscle relaxation and inhibition of platelet aggregation. Using whole-exome sequencing, our group also identified nine rare variants in the coding sequence of *GUCY1A3* [2]. Two of these variants were found in two extended families with a high prevalence of premature CAD/MI. Seven further rare variants were found in 252 young MI patients. In this study, we aimed to investigate the

functional implication of these rare variants found in CAD/MI patients (Table 1) regarding protein level, dimerization capability and enzymatic activity.

Two of the investigated α_1 variants exhibited significantly decreased protein levels compared to wild type α_1 . The amount of β_1 correlated with those of α_1 in all cases. All α_1 variants, except for p.Leu163Phefs*24, still dimerized with the β_1 subunit, as shown by co-immunoprecipitation. Using radioimmunoassay three of the rare variants demonstrated significantly decreased cGMP amounts at every time point tested (0.5/1/2 min). The activity only in part correlated with the observed protein levels pointing to an effect of the tested variants on enzymatic activity. As we have shown that loss of function-mutations in *GUCY1A3* may lead to CAD/MI [2], decreased enzymatic activity might also increase risk.

Table 1 Rare variants of sGC α_1 subunit found in MI patients:

Variant	Identified in	Predicted effect on protein function		
		PolyPhen-2	SIFT	SNAP
p.Leu163Phefs*24	MI family	frameshift	-	-
p.Lys53Glu	252 young MI cases	possibly damaging	damaging	non-neutral
p.Thr64Ala	252 young MI cases	benign	tolerated	neutral
p.Thr229Met	252 young MI cases	possibly damaging	tolerated	non-neutral
p.Ser478Gly	252 young MI cases	benign	tolerated	neutral
p.Val587Ile	252 young MI cases	benign	tolerated	neutral
p.Gly573Arg	MI family	probably damaging	affect protein function	non-neutral
p.Cys610Tyr	252 young MI cases	probably damaging	tolerated	neutral
p.Ile571Val	252 young MI cases	possibly damaging	affect protein function	neutral

* Correspondence: jana.wobst@tum.de

¹Klinik für Herz-und Kreislauferkrankungen, Deutsches Herzzentrum München, Technische Universität München, Munich, Germany

Full list of author information is available at the end of the article

Future studies focus on mRNA abundance and protein degradation to uncover the reason for attenuated activity of the respective variants.

Authors' details

¹Klinik für Herz-und Kreislauferkrankungen, Deutsches Herzzentrum München, Technische Universität München, Munich, Germany. ²Institut für Integrative und Experimentelle Genomik, Universität zu Lübeck, Germany. ³Deutsches Zentrum für Herz-Kreislauf-Forschung (DZHK) e.V., partner site Munich Heart Alliance (MHA), Munich, Germany. ⁴Deutsches Zentrum für Herz-Kreislauf-Forschung (DZHK) e.V., partner site Hamburg/Kiel/Lübeck, Lübeck, Germany.

Published: 2 September 2015

References

1. CARDIoGRAMplusC4D Consortium, Deloukas P, Kanoni S, Willenborg, Farrall M, Assimes TL, et al: **Large-scale association analysis identifies new risk loci for coronary artery disease.** *Nat Genet* 2013, **45**(1):25-33.
2. Erdmann J, Stark K, Esslinger UB, Rumpf, Koesling D, de Wit C, et al: **Dysfunctional nitric oxide signalling increases risk of myocardial infarction.** *Nature* 2013, **504**(7480):432-436.

doi:10.1186/2050-6511-16-S1-A100

Cite this article as: Wobst et al.: Functional evaluation of GUCY1A3 mutations associated with myocardial infarction risk. *BMC Pharmacology and Toxicology* 2015 **16**(Suppl 1):A100.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

