



POSTER PRESENTATION

Open Access

Genetic polymorphisms of inflammatory molecules in Tunisian inflammatory bowel diseases

Y Gorgi^{1*}, W Ben Aleya¹, D Khazen¹, I Sfar¹, H Aounallah-Skhiri⁴, L Mouelhi², S Jendoubi-Ayed¹, M Makhlouf¹, T Ben Rhomdhane¹, S Matri³, A Filali³, H Aouadi¹, T Najjar², K Ayed¹, T Ben Abdallah¹

From 5th European Workshop on Immune-Mediated Inflammatory Diseases
Sitges-Barcelona, Spain. 1-3 December 2010

Introduction

As chemokines and adhesion molecules play a major role in the process by which leukocytes are recruited from the bloodstream into sites of inflammation [1], genetic variation in these production or activity molecules may influence susceptibility to inflammatory diseases [2-4].

Aim

To detect a possible association between the functional polymorphisms of these molecules and susceptibility to Crohn's disease (CD) and ulcerative colitis (UC) in Tunisian population.

Materials and methods

We have analysed polymorphisms of CCR5-Δ32, CCR5-59029-A/G, CCR2-V64I, MCP-1 G/A (-2518), ICAM-1 G241R, PECAM-1 V125L, E-selectin L554F and L-selectin F206L in 194 Inflammatory bowel disease (IBD) patients and 169 healthy blood donors using PCR-RFLP and PCR-SSP methods. The patients were classified in 126 patients with CD and 68 patients with UC.

Results

A significant increase in allele frequency of 206L of L-selectin was observed in IBD patients compared with controls (OR: 1.53; 95%CI: [1.05-5.60]; $p=0.02$) but does not constitute factor influencing clinical manifestations. The genotypic and allelic frequencies of chemokine polymorphisms did not reveal significant differences between patients and controls, and among CD and UC

patients. However, analysis of CD patients revealed that those carrying A/A and/or A/G CCR5-59029 genotypes are more frequently in remission compared to those with G/G genotype (OR: 0.4; 95%CI: [0.174-0.928]; $p=0.03$). Also, the frequency of the 64I CCR2 muted allele was statistically higher in CD patients in remission disease than those in active form (OR: 0.267; 95%CI: [0.09-0.78]; $p=0.01$). Adjustment for known covariates (age, gender and immunosuppressive regimen) confirmed these univariate findings and revealed that the A/G CCR5-59029 and V64I CCR2 genotype were associated to remission form of CD (OR: 2.63 ; 95%CI: [1.01-6.80] ; $p=0.047$ and OR: 4.64 ; 95%CI: [1.01-21.31]; $p=0.049$ respectively).

Conclusion

The present study supports the involvement of chemokine receptor (CCR2 and CCR5) polymorphisms on the susceptibility to clinical course of IBD in Tunisian patients. However, further studies are needed to confirm the association of these diseases with allele L206 of L-selectin gene.

Author details

¹Immunology Research Laboratory (LR03SP01), Charles Nicolle Hospital, Tunisia. ²Dept. of Gastroenterology, Charles Nicolle Hospital, Tunisia. ³Dept. of Gastroenterology, La Rabta Hospital, Tunisia. ⁴National Public Health Institute, Tunisia.

Published: 25 November 2010

References

1. Dong VM, McDermott DH, Abdi R: Chemokines and diseases. *Eur J Dermatol* 2003, **13**:224-30.

¹Immunology Research Laboratory (LR03SP01), Charles Nicolle Hospital, Tunisia

Full list of author information is available at the end of the article

2. Palmieri O, Latiano A, Salvatori E, *et al*: The -A2518G Polymorphism of Monocyte Chemoattractant Protein-1 Is Associated With Crohn's Disease. *Am J Gastroenterol.* 2010, **105**(7):1586-94.
3. Herfarth H, Gökem M, Hellerbrand C, *et al*: Polymorphism of monocyte chemoattractant protein 1 in Crohn's disease. *International journal of colorectal disease* 2003, **18**:401-5.
4. Rector A, Vermeire S, Thoelen I, *et al*: Analysis of the CC chemokine receptor 5(CCR5) delta-32 polymorphism in inflammatory bowel disease. *Hum Genet* 2001, **108**:190-193.

doi:10.1186/1479-5876-8-S1-P7

Cite this article as: Gorgi *et al*: Genetic polymorphisms of inflammatory molecules in Tunisian inflammatory bowel diseases. *Journal of Translational Medicine* 2010 **8**(Suppl 1):P7.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

