

RESEARCH ARTICLE

Effects of Acupuncture on Sensory Perception: A Systematic Review and Meta-Analysis

Petra I. Baeumler^{1*}, Johannes Fleckenstein¹, Shin Takayama², Michael Simang^{1,3}, Takashi Seki², Dominik Irnich¹

1. Multidisciplinary Pain Center, Department of Anaesthesiology, University of Munich (LMU), Munich, Germany, 2. Department of Traditional Asian Medicine, Tohoku University, Sendai, Japan, 3. Institute for Medical Information Sciences, Biometry and Epidemiology, University of Munich (LMU), Munich, Germany

*Petra.Baeumler@med.uni-muenchen.de



CrossMark
click for updates

 OPEN ACCESS

Citation: Baeumler PI, Fleckenstein J, Takayama S, Simang M, Seki T, et al. (2014) Effects of Acupuncture on Sensory Perception: A Systematic Review and Meta-Analysis. PLoS ONE 9(12): e113731. doi:10.1371/journal.pone.0113731

Editor: Bing Hou, Beijing Institute of Radiation Medicine, China

Received: June 2, 2014

Accepted: October 29, 2014

Published: December 12, 2014

Copyright: © 2014 Baeumler et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper.

Funding: University of Munich & Tohoku University provided support in the form of salaries for all authors as depicted in the corresponding affiliations, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The specific roles of these authors are articulated in the 'author contributions' section.

Competing Interests: All authors declare that no competing interests exist.

Abstract

Background: The effect of acupuncture on sensory perception has never been systematically reviewed; although, studies on acupuncture mechanisms are frequently based on the idea that changes in sensory thresholds reflect its effect on the nervous system.

Methods: Pubmed, EMBASE and Scopus were screened for studies investigating the effect of acupuncture on thermal or mechanical detection or pain thresholds in humans published in English or German. A meta-analysis of high quality studies was performed.

Results: Out of 3007 identified articles 85 were included. Sixty five studies showed that acupuncture affects at least one sensory threshold. Most studies assessed the pressure pain threshold of which 80% reported an increase after acupuncture. Significant short- and long-term effects on the pressure pain threshold in pain patients were revealed by two meta-analyses including four and two high quality studies, respectively. In over 60% of studies, acupuncture reduced sensitivity to noxious thermal stimuli, but measuring methods might influence results. Few but consistent data indicate that acupuncture reduces pin-prick like pain but not mechanical detection. Results on thermal detection are heterogeneous. Sensory threshold changes were equally frequent reported after manual acupuncture as after electroacupuncture. Among 48 sham-controlled studies, 25 showed stronger effects on sensory thresholds through verum than through sham acupuncture, but in 9 studies significant threshold changes were also observed after sham acupuncture. Overall, there is a lack of high quality acupuncture studies applying comprehensive assessments of sensory perception.

Conclusions: Our findings indicate that acupuncture affects sensory perception. Results are most compelling for the pressure pain threshold, especially in pain conditions associated with tenderness. Sham acupuncture can also cause such effects. Future studies should incorporate comprehensive, standardized assessments of sensory profiles in order to fully characterize its effect on sensory perception and to explore the predictive value of sensory profiles for the effectiveness of acupuncture.

Introduction

Acupuncture is gaining popularity as a non-pharmacological option in pain medicine [1, 2]. There is substantial evidence for acupuncture being effective in the treatment of acute [3–5] and chronic pain [6]. However, for several other pain conditions such as neuropathic pain [7] or fibromyalgia [8], evidence remains inconclusive. In order to specify indications for which acupuncture should be used and to optimize treatment, it is crucial to understand how the effect of acupuncture is mediated.

Various mechanisms underlying the effect of acupuncture have been suggested. Brain imaging studies have shown that acupuncture alters activation patterns in brain areas associated with pain processing [9]. It is postulated that in response to the needle stimulation mechanisms of the endogenous pain modulation such as diffuse noxious inhibitory controls (DNIC), segmental inhibition, and descending pain control pathways lead to a decrease in pain perception [10, 11]. At this, various centrally and/or peripherally acting neuromodulators and neurotransmitters such as endorphins [12], serotonin [13], ATP [14], etc. have been identified to play an important role in the analgesic effect of acupuncture. In summary, one can assume that a modulation of the nervous system forms a central part of the effect of acupuncture although details are far from being understood. In special, effects on afferent nerve fibers which might be critical to the modulation of sensory perception by acupuncture remains unclear.

For investigating how acupuncture operates through the nervous system assessments of sensory threshold changes are essential. Evaluations of sensory detection and pain thresholds is referred to as Quantitative Sensory Testing (QST) and has been recognized an important tool in basic science, clinical trials, and for diagnostic and monitoring purposes [15]. QST is deemed to allow for inferences about the type of nerve fibers and about the structures of the nervous system that are affected by a disease or an intervention, according to which modality of sensory perception is changed and at which body sites these changes occur [16–18].

Despite the extensive use of sensory threshold assessment in acupuncture research, the impact of these data on the understanding of how acupuncture acts on the nervous system has never been systematically analyzed. Yet, there is no

consensus about which modalities of sensory perception (thermal and/or mechanical thresholds, detection and/or pain thresholds) are affected by acupuncture, and whether this effect is influenced by other factors e.g. the measurement tool, the type of stimulation or the target population. The aim of this systematic review, therefore, is to give an overview about data available on the effect of acupuncture on sensory thresholds and to substantiate the respective findings by meta-analyses of high quality studies. Our work provides the first summary of knowledge about how sensory perception is modulated by acupuncture which is crucial to approach a better comprehension of its mechanisms and to improve treatment.

Materials and Methods

The study protocol containing all steps followed for systematically reviewing literature and performing meta-analyses is available from the authors. Reporting was conducted in accordance with the PRISMA statement [19] as depicted in [S1 Table](#).

Literature Search

Pubmed, EMBASE, and Scopus were searched from their respective inception dates (Pubmed 1948, EMBASE1988, Scopus 1823) to the 1st of June 2012 using the following search strategy. 1: acupuncture; 2: perception; 3: sensory; 4: threshold; 5: pressure AND pain; 6: pain AND thermal; 7: heat AND pain; 8: cold AND pain; 9: mechanical AND pain; 10: vibration; 11: experimental AND pain 12: experimentally AND pain; 13: #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12; 14: #1 AND #13

Selection Criteria

We included research articles published in English or German which describe the effect of manual acupuncture (MA; needle insertion with or without manipulation), electroacupuncture (EA; needle insertion with electrical stimulation), or dry needling (DN; needle insertion into myofascial triggerpoints (MTrPs)) on thermal or mechanical detection or pain thresholds in humans. Animal studies, studies using other types of pain paradigms (e.g. electrical or ischemic pain), and studies investigating acupuncture related techniques (e.g. laser stimulation or transcutaneous electrical nerve stimulation) were excluded.

Article Selection

In a first screening step, two reviewers (PIB and JF or PIB and ST respectively) independently assessed the articles retrieved from the literature search for relevance by title and abstract. Full texts of all remaining articles were obtained and screened for eligibility by two independent reviewers (PIB and JF). When the

reviewers disagreed or had doubts, the full-text paper was evaluated by all authors. Remaining disagreements were resolved via discussion and consensus.

Quality Assessment

Two independent researchers (PIB, JF) evaluated the quality of the included studies by assessing the risk of bias by means of the Cochrane Collaboration's tool [20] and the quality of reporting of interventions and assessments of sensory thresholds based on the *Revised Standards for Reporting Interventions in Clinical Trials of Acupuncture* (STRICTA) [21]. In brief, risk of bias was assessed by answering questions about the following features of the studies with 'Yes' (low risk of bias), 'No' (high risk of bias) or 'Unclear' (lack of information or uncertainty over the potential for bias): random sequence generation, allocation concealment, blinding of participants, blinding of outcome assessment, incomplete outcome data, selective reporting and other bias. When sham acupuncture interventions were used as controls we assumed that participants were blinded for group allocation. Blinding of therapists was excluded from the risk of bias assessment since it is not possible to keep the acupuncturist unaware of the point location, stimulation and type of needle. Prior to analysis possible sources of 'other bias' were determined by consensus of the authors. This included bias due to a short washout phase in cross-over studies, questionable outcome assessment, and large baseline differences which were not taken into account in the subsequent analysis. Studies lacking a control group were a priori rated as having a high risk of bias. In addition, we report the total sample size as well as the number of study participants in each group.

Data Extraction

Articles were analyzed by two independent researchers (PIB, JF). Information from the included articles was extracted and tabulated.

Eligible studies are described according to the sensory threshold under investigation, the type of needle stimulation, the characteristics of the study population, and whether the immediate or long term effect of acupuncture was studied. The outcome of the studies is rated as positive or negative according to the authors' conclusion regarding pre-post treatment effects or group differences. Articles with elusive data presentation were rated as unclear. In addition, we compared the effects of verum acupuncture to the effects of inactive or sham-control procedures as well as local to distant needling effects (homo- to heterosegmental, ipsi- to contralateral).

Statistical Analysis - Meta-Analysis

Chi-squared test was used to test whether a positive study outcome (effect of acupuncture on at least one sensory threshold) was independent of the type of needle stimulation (MA or EA).

A meta-analysis was conducted in order to compare verum and sham acupuncture in high quality studies. Studies were eligible for meta-analysis if they fulfilled the following criteria: no rating of 'high risk of bias' in none of the items of the Cochrane risk of bias tool, sham-controlled, blinding of the outcome assessment, and clear reporting of data. All studies fulfilling these criteria were grouped according to reviewers' opinion about clinical homogeneity. We were able to combine studies in which the pressure pain threshold (PPT; kPa) had been assessed before and after a series of acupuncture treatments in patients suffering from musculoskeletal pain. One of the selected studies followed a cross over design [22]. Accordingly, we only included data obtained at baseline and after the first treatment phase. In cases of multiple evaluations of the outcome at one time point, e.g. several measure sites, data were averaged in order to achieve an equal weighting of all studies for the analysis. In order to account for baseline differences, we used delta scores (post-treatment values minus pre-treatment values) for all calculations. A conservative estimator of the delta score variance was obtained according to the variance sum law without correcting for dependent samples. Meta-analytical comparison of effects of verum and sham acupuncture on the PPT was performed by using the package 'metafor' from the R-project (Version 2.15.1, www.metafor-project.org). The standardized mean difference (SMD) was calculated by dividing the delta scores of the verum and sham acupuncture group by the pooled standard deviation of the two groups. Cochran's Q test was applied to evaluate statistical heterogeneity (I^2). We regarded heterogeneity between studies as substantial if Tau^2 was greater than zero and either I^2 was greater than 50% or the Cochran's Q test resulted in a low P value (less than 0.10). An assessment of reporting biases did not appear meaningful due to the small number of studies included in the meta-analyses.

Results

General Aspects

By electronic literature search we identified 3007 citations of which 2922 were excluded; 2830 were screened by title or abstract in a first selection step, and full text was obtained of 177 articles ([fig. 1](#)). Eighty five articles published between 1974 and 2012 met our inclusion criteria ([table 1](#) & [2](#)). More than half of these articles (50 out of 85, 58.8%) were issued after 1999.

Most of the studies were performed in the US and Europe (49 out of 85, 57.6%) and were published in English (81 out of 85, 95.3%). The majority (77 out of 85, 90.6%) describe the effect of acupuncture on a single sensory threshold [[22–98](#)], whereas eight studies (9.4%) assessed acupuncture evoked changes in more than one sensory threshold [[99–106](#)]. About half of the studies (44 out of 85) were conducted with healthy volunteers and evaluated the immediate effect of one acupuncture session ([table 1](#)). Forty-one studies (48.2%) included subjects suffering from pain conditions and assessed either the effect of one single




Fig. 1. Systematic Review Process Flowchart.

doi:10.1371/journal.pone.0113731.g001

acupuncture treatment (13 out of 41; 31.7%) or a series of treatments (29 out of 41; 70.7%; [table 2](#)).

In 65 studies (76.5%) an effect of acupuncture on at least one sensory threshold was observed. A statistically significant effect of acupuncture was found in 60 studies (70.6%). In contrast, 15 studies (17.6%) showed no effect on any sensory

Table 1. Studies Conducted with Healthy Subjects.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
Amand [24]	2011	CPT	MA	Non-acu. point stimulation Period of rest	TH 5, LI 4, SI 4 (dominant side)	Forearm (dominant side)	x verum = sham
Anderson [25]	1974	CPT	EA	Non-acu. point stimulation Period of rest	LI 11, LI 5, SI 5, SI 8 (right)	Hand (bilat.)	x Verum > sham
Ashton	1984	CPT	MA TENS 100 Hz TENS 8 Hz	Placebo pill	PC 7 (non-dom. side)	Hand (non-dom.)	x
Barlas [28]	2006	PPT	EA low intensity EA high intensity	Non-penetrating needle with or without patient blinding	LI 10, TH 5, GB 34, ST 38 (dom. side)	First dorsal interosseous muscle of the hand (bilat.)	x Verum > sham
Benoliel [29]	2011	WDT	MA 4 points MA 6 points	-	1) ST 6, LI 4 (bilat.) 2) ST 6, LI 4, ST 2 (bilat.)	Dermatome of infraorbital nerve & mental nerve (bilat.)	x
Berlin [30]	1975	HPT	EA	Non-acu. point stimulation Period of rest	LI 4, TH 5 (left)	Post lateral aspect of the left forearm	x verum = sham
Brockhaus [31]	1990	HPT	MA LA	Non-acu. point stimulation Sham-laser	LI 4 (bilat.)	Forearm ventral side (left)	x Verum > sham
Chae [32]	2006	HPT	MA	-	LI 4 non-dom. hand	Distal digit of finger of dom. hand	x
Clark [34]	1974	HPT	EA ipsilat. EA contralat.	-	HT 1, SI 9, HA 3 LU 5, TH 3, LI 4 (body side according to study group)	6 sites on volar surface of forearm (bilat.)	x
Croze	1976	HPT	MA	Non-acu. point stimulation • close to measure site • distant to measure site Period of rest	LI 10, ST 36 (right)	Thenar eminence (left)	unclear
Day [36]	1975	HPT	EA	-	LI 4, GB 21, TH 8, LI 14 (bilat.)	Area over thyroid gland, chest (right)	-
Downs [99]	2005	CDT, WDT CPT, HPT	MA	Non-penetrating needle Period of rest	TH 5, LI 11 (right)	Thenar eminence (right)	- verum = sham
Farber [39]	1997	PPT	EA	-	LI 4, 1 needle at 5 mm distance for EA	LI 5, LI 11, 15 mm lateral from LI 5 or LI 11,	x

Table 1. Cont.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
Galloon [42]	1977	HPT	EA	Non-acu. point stimulation	induction (unilat., side not indicated) not indicated	(side not indicated), LI 20 Forehead, throat, chest, forearm, upper leg (right)	- verum = sham
Kitade [55]	1988	HPT	Morphine EA + D-Phenylala.	Placebo pill EA + placebo pill	LI 4, LI 10 (bilat.)	Palmar side of forearm (side not indicated)	unclear
Kitade [53]	1979	HPT	7 different EA regimens	Non-acu. point stimulation (auricular) Period of rest	1) auric. lung (bilat.) 2) auric. sympathetic (bilat.) 3) auric. shen-men (bilat.) 4) auric. kidney (bilat.) 5) auric. neck (bilat.) 6) auric. elbow-arm (bilat.) 7) auric. lung + LI 4 (bilat.)	ST 32, KI 23, BL 11, PC 4, GB 35 (bilat.), EX-HN 3	x (no statistics) Verum > sham
Kitade [54]	1981	HPT	EA + D-Phenylalanine EA + L-Phenylalanine	EA + lactose D-Phenylalanine	LI 4 & ST 36 (bilat.)	Center part of anterior forearm	x (no statistics)
Knox	1977	CPT	EA	Period of rest	LI 4, EX-JUE 2 (right)	Forearm (right)	x
Knox [56]	1981	CPT	EA	Period of rest	LI 4, EX-JUE 2 (right)	Forearm (right)	-
Knox [57]	1979	CPT	Hypnosis EA	Non-acu. point min. stimulation Period of rest	LI 4, TH 5 (unilat., side not indicated)	Hand & forearm (ipsilat.)	x Verum > sham
Knox [59]	1977	CPT	EA	Period of rest	LI 4, EX-JUE 2 (unilat., side not indicated)	Hand & forearm (ipsilat.)	-
Kong [60]	2005	HPT	EA	Non-penetrating needle	LI 4, ST 36, SP 6 (right)	Forearms & legs (bilat.)	x
Kong [61]	2009	HPT	MA EA	Non-penetrating needle	LI 3, LI 4 (right)	Medial aspect of forearm (right)	Verum > sham x
Lang [100]	2010	HPT, CPT, WDT, CDT,	MA	-	SP 6, SP 9, ST 36, GB 39 (left)	Anterolateral skin of lower limb	verum = sham x

Table 1. Cont.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
		TSL, MDT, MPS, MPT, VDT, PPT	EA (80 Hz)			(sensory region of peroneal nerve; bilat.)	
Leung [101]	2005	CDT, WDT, CPT, HPT,	EA (2 Hz) EA	-	SP 1, LR 1 (left)	4 points on calf & thigh along spleen & liver meridian (bilat.)	x
Leung [102]	2008	CTD, WDT, CPT,	EA 5 min	-	SP 1, LR 5 (left)	4 points on calf & thigh along spleen & liver meridian (bilat.)	x
		HPT, MDT	EA 15 min EA 30 min				
Li [65]	2008	PPT	4 different MA regimens	-	1) LI 4 (right) 2) LI 4 (bilat.) 3) LI 11 (right) 4) LI 11 & LI 4 (right)	LI 10, GB 20 (right) ST 36, KI 3, LI 5, non-acu. points on the upper limbs (bilat.)	x
Lim [66]	1977	HPT	5 different EA regimens	Acu. point no stimulation	1) LI 4 (right)	Forearm (bilat.)	x
					2) ST 36, 1) (right) 3) Ulnar nerve (right) 4) Auricular: shenmen, 3) (right) 5) LI 4, 4) (right)		verum = sham
Lin [67]	1981	HPT	2 different MA regimens	-	1) SP 6 (right)	Palm & sole of the foot (bilat.)	x
Lloyd [69]	1976	HPT	EA	-	2) LI 11 (left) 2x near elbow, 1x ventral & 2x dorsal part of forearm, 2x between thumb & first finger (right)	Back of the hand (right)	-
Lundeberg [103]	1989	TSL VDT	MA 2 regimens	Acu. point no stimulation (LI 4)	1) ST 7 (bilat.)	TSL: upper lip	-
			EA 2 Hz 2 regimens	Acu. point no stimulation (ST 7)	2) LI 4 (bilat.)	VDT: tip of index finger, forearm, fore head (side not indicated)	verum = sham
			EA 80 Hz 2 regimens				

Table 1. Cont.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
Lynn [105]	1977	HPT, CPT MPS, MPT	5 different EA regimens	-	1) Auricular: stomach 1) ST 36, SP 6 2) Auricular: stomach, ST 36, SP 6 3) SP 6, SP 9 4) LI 4, PC 6 (each bilat.)	Thigh, abdomen, neck (side not indicated)	x
Moret [71]	1991	CPT	EA + placebo pill Hypnosis + placebo pill	EA + Naloxone Hypnosis + Naloxone	LI 4, LI 11 (right)	Hand (right)	x
Pauser [73]	1975	MPT	EA	Non-acu. point stimulation	PC 6, LI 4, auric. lung (side not indicated)	5 defined points on the neck (side not indicated)	x
Schlessbach [76]	2011	PPT	EA	Non-penetrating needle	LI 4, LI 11 (unilateral, side not indicated)	2nd toes (ipsilat.)	verum = sham x
Schlessbach [98]	2012	PPT	MA	Non-penetrating needle	LI 4 (unilateral, side not indicated)	2nd toes (ipsilat.)	Verum > sham x
Shukla [80]	2011	HPT	EA	Acu. point min. stimulation	LR 1, SP 1 (left)	Medial calf (left)	verum = sham x
Stern	1977	CPT	EA	Non-acu. point stimulation	LI 4, LI 11, LI 14, LI 15 (left)	Length of fingers (left)	Verum > sham x
Stewart [85]	1977	HPT	EA	Non-acu. point stimulation	LI 4, ST 36 (bilat.)	Epigastrium, sternum, upper arm & lower leg (left), forearm & thigh (right)	Verum > sham x
Umino [90]	1984	HPT	EA	Period of rest	LI 4, LI11, ST36 (bilat.)	Forearm (side not indicated)	Verum > sham x
Wang [92]	2009	CPT	EA 20 min	Acu. point no stimulation	ST 36, SP 6 (left)	3 sites on lower leg (medial, right)	x
Yoon [94]	1986	HPT	EA	Non-acu. point stimulation	1 individually chosen acu. point (bilat.)	ST 6, LI 20 (bilat.)	Verum > sham x (no statistics)
Zaslowski [95]	2003	PPT	MA	Acu. point no stimulation	LI 4 (right)	LI 20, PC 6, SI 3, 2R, LI 10, 1R, LI 5, CV 12, 3R, ST 36,	Verum > sham x

Table 1. Cont.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
				Non-acu. point stimulation		Non-acu. points on forearm, wrist, & lower limb (right)	Verum > sham
				Non-acu. point no stimulation			
				Sham-laser			
Zhang [97]	2003	CPT	EA	Acu. point min. stimulation	ST 36, SP 6 (left)	Thenar eminence (left)	x Verum > sham

Sensory thresholds are abbreviated as follows: CDT: cold detection threshold; CPT: cold pain threshold; HPT: heat pain threshold; MDT: mechanical detection threshold; MPS: mechanical pain sensitivity; MPT: mechanical pain threshold; PPT: pressure pain threshold; TSL: thermal sensory limen; VDT: vibration detection threshold; WDT: warm detection threshold; Acupuncture styles are abbreviated as follows: EA: electroacupuncture; MA: manual acupuncture; DN: dry needling; Acupuncture points are abbreviated according to the WHO standard international nomenclature [147].

doi:10.1371/journal.pone.0113731.t001

Table 2. Studies Conducted with Patients.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Treatment Frequency	Subjects	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
Abuaisha [23]	1998	VDT	MA	-	6x within 10w	Diabetic neuropathy	LI 3, SP 6, SP 9, ST 36 (all bilat.)	Great toe (side not indicated)	-
Ahn [106]	2007	WDT	MA (Japan. style)	-	1x/w for 10w	Diabetic neuropathy	Individually chosen according to acupuncture style used	Lower limb (bilat.)	unclear
Bartas [27]	2000	PPT	MA (TCM)	Non-acu. point min. stimul.	1x/d for 5d	Healthy induced DOMS	1) PE 2, LI 11, LU 5, LI 4 (non-dom. side) 2) 4 most tender points, of biceps brachii (non-dom. side)	Biceps brachii muscle at 8 equidistant points (non-dom. side)	verum = sham
Chou [33]	2011	PPT	DN	Non-penetrating needle	1	Chronic muscle pain	TH 5, LI 11 (ipsilat. to active MTrP)	Active MTrP in upper trapezius muscle	x
Deluze [37]	1992	PPT	EA	Non-acu. point min. stimul.	6x within 3w	Fibromyalgia	LI 4, MA 36 (bilat.)	18 tender points	x
Edwards [38]	2003	PPT	DN + Stretch	Stretch	Individually over 3w	Musculoskeletal pain	Active MTrPs	MTrP treated	Verum > sham
Fernandes-	2010	PPT	DN	Acu. point no stimulation	1	Temporomandibular disorder	Most painful MTrP in m. masseter & mandibular condyle	MTrP treated	x
Camero [40]	2007	PPT	DN along muscle	-	1	MTrPs in neck	7–8 cm aside from the most painful MTrP in the neck	MTrP treated	verum > sham
Goddard [43]	2002	PPT	MA	Non-acu. point stimulation	1	Pain in jaw muscle	LI 4, ST 6 (bilat.)	Sensitive area of m. masseter	x
Harris [45]	2008	PPT	MA	Non-penetrating needle	9x within 4w	Fibromyalgia	"traditional acupuncture treatment" (not further indicated)	Thumb nail (left)	verum = sham unclear
Harris [44]	2006	PPT	MA	Acu. point no stimulation	1x/w (week 1–3)	Fibromyalgia	"traditional locations" (not further indicated)	Thumb nail, lateral epicondyle (bilat.)	unclear

Table 2. Cont.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Treatment Frequency	Subjects	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
				Non-acu. point stimulation	2x/w (week 6–8)				
				Non-acu. point no stimul.	3x/w (week 11–13)				
He [46]	2004	PPT	EA + MA + auricular seeds	Non-acu. point no stimul. + auricular seeds	10x within 3–4w	Chronic neck & shoulder pain	EA: 4 Ex-points (not indicated), GB 21 (bilat.), BL 12, DU 14, SI 14, SI 15 (unilat. side not indicated)	28 MTPs bilat. on neck & shoulders	x
							MA: LI 4, LI 11, GB 31 (bilat.), auricular seeds: shenmen, neck, cervical spine, shoulder, shoulder-joint, shoulder-back (unilat.)		Verum > sham
Hübscher [47]	2008	PPT	MA	Non-acu. point min. stimul.	Immediately,	Healthy induced DOMS	GB 34, LU 3, LU 5, LI 11, SP 10,	7 equidistant points along line joining insertion of	-
				Period of rest	24 h and 48h post		ah shi-points (non-dom. side)	biceps brachii on radius & acromion	verum = sham
				DOMS induction				(non-dom. side)	
Ilbuldu [48]	2004	PPT	DN	Sham-laser	DN 1x/w for 4w	MTPs in upper trapezius	3 MTPs in the upper trapezius on both sides	Not indicated	-
			LA		LA/SL 3x/w for 4w	muscle			verum = sham
Irnich [49]	2001	PPT	MA + DN	Sham-laser	5x within 3w	Chronic neck pain	Individualized MA (mostly SI 3, UB 10, UB 60, LR 3, GB 20, GB 34, TH 5, auric. neck), DN at MTPs	Levator scapulae, trapezius descendens, paravertebral of 6th cervical spine (bilat.)	-
			Message						verum = sham
Irnich [50]	2003	PPT	MA	Non-acu. point stimul.	3x within 10d	Epicondylopathy	LI 4, LI 10, SI 3, GB 34, TH 5 (unilat. on affected side)	Insertion of common tendon of m. extensor carpi	x
								radialis at lateral elbow, belly of m. extensor	Verum > sham
								carpi radialis brevis at transit between proximal	

Table 2. Cont.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Treatment Frequency	Subjects	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
Iltho [51]	2011	PPT	MA seg.3 mm MA seg.10 mm	Period of rest	1	Healthy induced DOMS	1) & 2) max. tender point within m. extensor digitorum 3) max tender point on distal third of belly of tibialis anterior	third & distal two thirds of affected forearm 20 mm distal to max. tender point within m. extensor digitorum m. extensor digitorum	x
Karst [52]	2000	PPT	MA heteroseg. 10 mm MA	Non-penetrating needle	2x/w for 5 w	Chronic tension-type headache	Obligatory: LI 4, LR 3, GB 20; Optional: GB 8, GB 14, GB 21, GB 41, UB 2, UB 10, UB 60, LU 7, TH 5, ST 8, ST 36, ST 44, DU 20 (maximum of 15 needles, side not indicated)	Temporal region where palpation had shown anterior part of temporal muscle to be most prominent (bilat.)	x
Kotani [62]	2001	PPT	Press needles at painful points	Press needles at non-painful points	20x 24 h within 4w	Abdominal scar pain	Painful points in scar area	Painful points treated in scar area	x
Kummerddee [63]	2009	PPT	MA	-	5x within 10d	Myofascial back pain	7 acupuncture points, MTRPs (location not indicated)	All MTRPs	x
Li [64]	1983	PPT	EA affected side EA normal side	-	1	Syringo-myelia	LI 4, PC 6 (unilat., side according to study group)	TH 5 (bilat.), forehead	x
List [68]	1993	PPT	MA + EA	Waiting list	6-8x within 6-8w	Temporomandibular disorder	MA:EX-HN 2, ST 7, ST 6, GB 20 EA at LI 4, ST 36	Belly of the right & left masseter muscle	x
Lundeberg [104]	1988	TSL	MA	Acu. point min. stimulation	1	Sinus pain,	LU 7, LI 4, GB 20, ST 6, ST 7, LI 20, EX-HN 3, EX-HN 5	Face (bilat.)	-
		HPT	EA 2 Hz	Placebo TENS		Healthy controls	(painful side)	dorsal aspect of hand on painful side	-

Table 2. Cont.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Treatment Frequency	Subjects	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
Ma [70]	2010	PPT	EA 80 Hz MA + stretching	Stretching	1–2x within 1–2w	Myofascial pain syndrome	MTrPs in upper trapezius	MTrPs (not further indicated)	x
Nabeta [72]	2002	PPT	Miniscalpel + stretching MA	Non-penetrating needle	1x/w for 3w	Fibromyalgia	All tender points (neck, shoulders, back)	All tender points (neck, shoulders, back)	x
Perez-Palomares [74]	2010	PPT	DN	-	1x/w over 3w	Chronic low back pain	8 needles placed bilat. within dermatomes L2 to L5	MTrPs (location not indicated)	x
Price [75]	1984	HPT	EA	-	1	Low back pain	1) BL 24, BL 25, BL 26, BL 27 2) GB 30+1 3) BL 50, BL 51, BL 57 (bilat.; semi-standardized; 3-5 needles)	Lower back & volar forearm (of painful side)	x (no statistics)
Seidel [77]	2002	PPT	MA	Sham-laser	2x/w for 4w	Chronic neck pain	15 acu. points (individually chosen, mostly bilat.)	Not indicated	x
Shen [78]	2007	PPT	LA 7 mWatt LA 30 mWatt MA	Non-penetrating needle	1	Healthy + clenching teeth 2min (DOMS)	LI 4 (left)	M. masseter (right)	x verum = sham
Shen [79]	2009	PPT	MA	Non-penetrating needle	1	Myofascial pain in jaw muscle	LI 4 (left)	M. masseter (right)	x Verum > sham
Singh [81]	2006	PPT	MA	-	2x/w for 8w	Fibromyalgia	Semi-stand. alternating anterior (GB 20, GV 14, GB 21, SI	18 tender points (bilat.)	x Verum > sham

Table 2. Cont.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Treatment Frequency	Subjects	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
Sprott [82]	2000	PPT	MA	-	1x/w for 6 w	Fibromyalgia	12, GB 30, BL 25, BL 23, BL 40) & posterior points (LR 3, GB 34, KI 25, LI 4, ST 36, SI 11, ST 40, SP 6, LR 8)	12 tender points (bilat.)	x
Srbley [83]	2010	PPT	DN	Non-acu. point stimulation	1	MTrP in m. supra-spinatus, m. infra-spinatus & m. gluteus medius	MTrP in m. supra-spinatus (right)	MTrPs in m. infra-spinatus & m. gluteus medius (right)	x
Takeda [86]	1994	PPT	MA	Non-acu. point min. stim.	3x/w for 3w	Osteoarthritis of the knee	EX 31, EX 32, SP 9, ST 35, GB 34 (affected knee)	Four points on the affected knee: medial & lateral joint lines, distal musculo-tendinous junctions of m. vastus medialis & lateralis	x
Targino [87]	2008	PPT	MA + tricyclic antidepressants. + exercise	Tricyclic antidepressants. + exercise	2x/w for 10w	Fibromyalgia	LR 3, LI 4, PC 6, GB 34, SP 6 (bilat.) Ex-HN 3	18 tender points (bilat.)	x
Tong [88]	2010	VDT	MA	Acu. point min. stimulation	1x/d for 15d	Diabetic neuropathy	LI 4, ST 40, LI 11, ST 36, SP 6 (bilat.)	Medial malleolus muscle in lower extremities	x
Ulett [89]	1978	CPT	EA	-	1	Chronic pain	LI 4, LI 11, LI 15, LI 14 (left)	Length of fingers (left)	Verum > sham
Vincent-Barrero [91]	2012	PPT	MA	-	1x/d for 3d,	Temporomandibular disorder	EX-HN 5, TH 21, GB 2, TH 17, ST 6, LI 4, ST 36, TH 5,	Preauricular, m. masseter,	x
			Decompression splints						

Table 2. Cont.

Author et al.	Year	Threshold	Verum Interventions	Control Interventions	Treatment Frequency	Subjects	Acupuncture Points (Verum Treatment)	Measure Site	Threshold Changes
Xue [93]	2004	PPT	EA	Non-acu. point min. stimul.	4 week phase	Tension type headache	Acu. points individually chosen	Frontal, suboccipital, posterior cervical,	x
Zhang [96]	2009	PPT	MA: PC7	-	5x/w for 2w	Plantar fasciitis	PC 7 or LI 4 (contralat. to pain or bilat. when bilat. pain)	m. masseter & temporalis (bilat.)	Verum > sham
Zhu [22]	2002	PPT	MA + EA	Non-acu. point min. stimul.	9x/3w	Chronic neck pain	Individualized: MA at two local points (1 bilat.); 2 distal points & EA at 2 distal points	Medial tubercle of calcaneum of non-painful foot & most painful site on painful foot	verum = sham

Sensory thresholds are abbreviated as follows: CDT: cold detection threshold; CPT: cold pain threshold; HPT: heat pain threshold; MDT: mechanical detection threshold; MPS: mechanical pain sensitivity; MPT: mechanical pain threshold; PPT: pressure pain threshold; TSL: thermal sensory limen; VDT: vibration detection threshold; WDT: warm detection threshold; Acupuncture styles are abbreviated as follows: EA: electroacupuncture; MA: manual acupuncture; DN: dry needling; Acupuncture points are abbreviated according to the WHO standard international nomenclature [147].

doi:10.1371/journal.pone.0113731.t002

Table 3. Overall Study Outcome.

Threshold	Healthy			Patients		
	+	-	?	+	-	?
Heat Pain	18	4	2	1	1	-
Cold Pain	10	6	-	1	-	-
Pressure Pain	7	-	-	27	6	2
Mechanical Pain	3	-	-	-	-	-
Warm Detection	3	2	-	-	-	1
Cold Detection	2	2	-	-	-	1
Thermal Sensory Limen	1	2	-	-	1	-
Mechanical Detection	-	2	-	-	-	-
Vibration Detection	0	2	-	1	1	-

“+” indicates a change of the respective threshold through acupuncture, “-” indicates no effect of acupuncture, “?” indicates an unclear study outcome. Studies in which more than one sensory threshold was assessed are listed several times, respectively.

doi:10.1371/journal.pone.0113731.t003

threshold. In five of the included publications (5.9%), authors report a pronounced threshold change after acupuncture, but no statistical analysis was provided. Still, we classified the outcome of these studies as positive, in terms of an effect of acupuncture, on the basis of individual reasoning. The overall review results were not affected by these studies and the individual reasoning will be stated in the respective chapter [53, 54, 74, 81, 94]. The results of another five studies (5.9%) were rated as unclear due to poor reporting [35, 55, 106], or because only a pooled analysis of all study arms was reported (table 3).

Forty studies (47.1%) investigated the effect of EA, 41 studies (48.2%) the effect of MA. Both were equally frequent shown to be effective in changing sensory thresholds (EA 32 out of 40, 80.0%; MA 30 out of 41, 73.2%; Chi-squared-test n.s.). In three studies effects of EA tended to be larger than those of MA [60, 76, 100]. DN was evaluated only with regard to its effect on the PPT, which was found to be elevated in six out of seven studies (8.2%). In four studies (4.7%) different types of needle stimulation were combined, with two showing a significant effect and two showing no effect. In six studies (7.1%) acupuncture was combined with another intervention. Among these, five resulted in altered sensory thresholds.

Quality of Included Studies

Twenty nine out of the 85 included studies (34.1%) were performed with a group size of 20 subjects or more [22, 23, 31, 37, 41, 48–50, 52, 56, 57, 62, 64, 65, 68, 74, 76, 81–84, 86–89, 93, 96, 98, 100], and ten studies (11.8%) did not include a control group [23, 32, 36, 39, 69, 75, 81, 82, 90, 101] (table 4).

Results of the risk of bias assessment are presented in table 4 and in summary in fig. 2. Only six studies (7.1%) were rated as ‘low risk of bias’ in six or more items of the Cochrane risk of bias tool [33, 37, 40, 47, 49, 62]. A judgment as ‘high risk of bias’ resulted most frequently from methodological shortcomings within the

Table 4. Quality Assessment.

Author	Total Sample Size (Comp. Size)	Risk of Bias Assessment					Details on Reporting										Details on Outcome Assessment			
		Random Sequence Generation	Allocation Concealment	Blinding of Participants	Blinding of Outcome Assessment	Incomplete Outcome Data	Selective Reporting	Other Bias	Number of Studies per Outcome	Points	Depth	Response (Depth/Touch)	Stimulation	Needle in Time	Needle (Thickness or Length/Duration)	Number of Sessions	Frequency/Duration of Treatment	Body Site	Time Point	Tool (Treatment/Default)
Avasthi 1996	44	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Ali 2007	7 (40)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Amend 2011	36 (10/12)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Anderson 1974	30 (10/10)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Anton 1984	464 (10/11/12)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Bales 2005	48 (20/10/15)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Bales 2006	48 (10/10/10)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Beard 2011	20 (10/5)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Bell 1976	30 (10/10)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Brookman 1980	NA (not reported)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Chen 2006	15	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Chen 1996/2011	45 (10/10/10)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Chen 1974	15 (10)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Chen 1976	6 (none used)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Day 1975	4	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
DeLoo 1982	70 (30/10)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Davis 2005	16 (none used)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Edwards 2003	40 (10/10/10)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Faber 1987	8	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Fernando 2010	13 (none used)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Fu 2007	47 (20/20)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Gilman 1977	13 (none used)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Graded 2002	16 (10)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Hahn 2006	65 (10/10)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Hahn 2008	10 (10)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Hu 2004	24 (10/10)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Hübner 2008	22 (10/7)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Itala 2004	60 (20)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Irwin 2011	177 (10/10/10)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Irwin 2003	50 (20/20)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Joh 2011	22 (10/10/10)	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Kell 2000	39 (20/10)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Kobak 1979	8 (none used)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Kobak 1981	3 (none used)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Kobak 1988	150 (10/10/10)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Koo 1977	48 (10/10)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Koo 1977	48 (10/10)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Koo 1979	72 (10/10/10)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Koo 1981	40 (20/20)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y

Table 4. Cont.

Author	Total Sample Size (Group Size)	Risk of Bias Assessment				Details on Reporting										Details on Outcome Assessment				
		Random Sequence Generation	Allocation Concealment	Blinding of Participants	Blinding of Outcome Assessment	Interrater Outcome Data	Selective Reporting	Other Bias	Number of Needles per Session	Points	Depth	Response (e.g. Touch)	Stimulation	Needle-in-Time	Needle (Thickness or Length/ Diameter)	Number of Sessions	Frequency/ Duration of Treatment	Body BA	Time Point	Tool (Treatment, Detailed)
Kong 2005	11 (eyes over)	?	?	+	-	+	?	+	+	+	+	+	+	+	+	+	+	+	+	+
Kong 2009	24 (12/12)	?	?	+	?	+	-	+	+	+	+	+	+	+	+	+	+	+	+	+
Robert 2001	10 (5/5/2/4)	+	+	+	+	+	?	+	+	+	+	+	+	+	+	+	+	+	+	+
Kernotzke 2009	18 (9/9)	?	?	-	+	+	?	+	+	+	+	+	+	+	+	+	+	+	+	+
Lang 2010	24 (eyes over)	?	?	?	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Leung 2005	13	n.c.	n.c.	n.c.	n.c.	+	?	+	+	+	+	+	+	+	+	+	+	+	+	+
Leung 2008	16 (eyes over)	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Li 2008	22 (eyes over)	?	?	+	+	+	?	+	+	+	+	+	+	+	+	+	+	+	+	+
Li 1983	25 (eyes over)	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Lin 1977	60 (15/15/15/15/15)	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Lin 1981	8 (eyes over)	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Liu 1993	15 (20/20/15)	?	?	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Lloyd 1976	8	n.c.	n.c.	n.c.	n.c.	+	?	+	+	+	+	+	+	+	+	+	+	+	+	+
Lundberg 1984	refers to (7/7/7/7) healthy controls 15 (3/3/3/3/3)	?	?	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Lundberg 1989	6 (eyes over)	-	-	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Lynn 1977	24 (4/4/4/4/4)	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Mu 2010	43 (10/15/13)	+	?	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Mori 1991	8 (eyes over)	?	?	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Nahata 2002	24 (12/12)	+	+	+	-	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Phalar 1985	16 (eyes over)	?	?	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Perez-Almaraz 2010	122 (62/60)	+	-	?	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Piro 1984	15	n.c.	n.c.	n.c.	n.c.	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Schneidbach 2011	48 (eyes over)	+	?	+	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Schneidbach 2012	48 (eyes over)	+	?	+	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Sekel 2002	8 (12/12/13/13)	+	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Shen 2007	16 (16/8)	-	-	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Shen 2009	28 (14/12)	+	-	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Shukla 2011	10 (eyes over)	?	?	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Shyu 2006	21	n.c.	n.c.	n.c.	n.c.	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Sprui 2000	20	n.c.	n.c.	n.c.	n.c.	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Stacy 2010	46 (20/23)	+	?	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Steen 1977	20 (eyes over)	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Stewart 1977	12 (eyes over)	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Takemi 1994	40 (20)	?	?	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Tangro 2008	58 (34/24)	+	?	-	+	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Tong 2010	63 (42/21)	+	-	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Ueki 1978	20 (eyes over)	?	?	-	-	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Umeno 1984	10	n.c.	n.c.	n.c.	n.c.	+	?	?	?	?	?	?	?	?	?	?	?	?	?	?




Fig. 2. Risk of Bias Assessment. Results of the risk of bias assessment as depicted in [table 4](#) are summarized. Percentages of studies with 'low risk of bias', 'high risk of bias', or 'unclear risk of bias' are illustrated for each item of the Cochrane Collaboration's tool.

doi:10.1371/journal.pone.0113731.g002

randomization process and the choice of comparison groups not allowing for blinding of participants (8.2% random sequence generation, 14.1% allocation concealment, 11.8% blinding of participants). Blinding of the outcome assessment was assured in less than half of the included studies (42.4%). The overall large proportions of studies with 'unclear risk of bias' reflects the low quality of reporting. In contrary, overall reporting of assessments of sensory thresholds was good and treatment regimens were reported in a mediocre manner ([table 4](#)). In 23 of the included publications (27.1%), information was missing for three or more items of the STRICTA guidelines [21]. About a quarter of all studies (24.7%) met all of the nine treatment related items. Threshold measurements were well described in 62 studies (72.9%), but especially combinations of different threshold assessments lacked standardization. Among the eight studies (9.4%), that evaluated acupuncture evoked changes in more than one modality of sensory perception, there was a single study which applied a comprehensive and standardized test battery for sensory testing [100].

Comparison of Verum Acupuncture to Inert or Sham-Control Procedures

Nineteen studies (22.4%) compared acupuncture to an inert control such as a placebo pill [26, 42], a waiting list [38, 62, 68], or a period of rest [24, 25, 27, 30, 35, 47, 51, 53, 56–59, 85, 99]. In 12 of these studies, acupuncture had a greater effect than the inert control procedure




Fig. 3. Meta-Analysis. Considering the previously defined quality criteria (see [methods](#)) as well clinical homogeneity according to the investigators accordance, five studies on the effect of acupuncture on the PPT conducted with patients were combined in two meta-analyses. The short-term effect on the PPT directly after a series of acupuncture sessions as assessed in four studies (A) was found to be significant but small. Two studies were analyzed regarding long-term effects of acupuncture on the PPT (B) which was also found to be significant but small. Heterogeneity was found to be substantial in both analyses.

doi:10.1371/journal.pone.0113731.g003

[24–26, 30, 38, 51, 53, 57, 58, 62, 68, 85], while results of six studies indicate no or only a minor change of the respective sensory threshold in any of the study groups [27, 42, 47, 56, 59, 99]. Results of one study were unclear. One article reported a significant change of sensory thresholds after an inert control procedure [24]. All four studies evaluating acupuncture as an add-on treatment reported a significant additional effect [38, 46, 70, 87].

In 48 studies (56.5%) one or several types of sham acupuncture treatments were used as controls. In the following, the term “sham” is used according to the description of the respective publication. Sham interventions were needling at non-acupuncture points with stimulation (17 out of 48; 35.4%) [24, 25, 30, 31, 35, 42–44, 50, 53, 62, 73, 83–85, 94, 95], without or with minimal stimulation (10 out of 48; 20.8%) [22, 27, 37, 44, 46, 47, 57, 86, 93, 95], minimal or no intensity needle stimulation at classical acupuncture points (10 out of 48; 20.8%) [40, 44, 66, 80, 88, 92, 95, 97, 103, 104], non-penetrating placebo needles (12 out of 48; 25%) [28, 33, 45, 52, 60, 61, 72, 76, 78, 79, 98, 99], or treatment with an inactive laser pen (5 out of 48; 10.4%) [31, 48, 49, 77, 95]. In 25 of these 48 studies (52.1%) the effect on sensory thresholds was significantly larger in the verum than in the sham acupuncture group [25, 28, 31, 33, 37, 40, 46, 50, 52, 57, 60, 62, 72, 76, 78–80, 83–85, 88, 92, 93, 95, 97]. In two studies (4.2%), it was shown descriptively that verum acupuncture was

superior to sham acupuncture, but no statistical analysis was provided [53, 94]. In nine studies (18.8%) both, verum and sham acupuncture, had a significant effect on the respective sensory threshold [43, 61, 62, 66, 73, 76, 77, 86, 98]. Among these, two studies also found verum acupuncture to cause significantly larger threshold changes than sham interventions [62, 76]. In nine other studies (18.8%) no effect was observed neither after verum nor after sham acupuncture [22, 27, 42, 47–49, 99, 103, 104], and the results of three studies (6.3%) were rated as unclear [35, 44, 45]. None of the twelve studies including both control modalities found significant differences between the sham and the inert control group [24, 25, 27, 30, 35, 42, 47, 53, 57, 62, 85, 99], although in three of these studies, the pre-post comparison indicates larger effects in the sham than in the inert control group [24, 30, 53, 85].

Heat Pain Threshold (HPT)

All 26 studies (30.6%) that assessed whether acupuncture had an effect on the HPT evaluated immediate treatment effects. Nineteen of these studies (73.1%) showed that HPT was elevated through acupuncture. In five articles (19.2%), no statistical analysis was provided. Nevertheless, in three cases we agree with a positive rating of the results because of the following reasons. In two studies case numbers were too low to allow for statistical calculations, but HPT elevations following acupuncture were observed in all subjects [53, 54]. In another study, two thirds of all subjects (6 out of 9, 66.6%) showed an elevated HPT after EA, but not after stimulation at non-acupuncture points [94]. In contrast, the results of two studies which claim to indicate an effect of acupuncture on the HPT were rated as unclear, because mean HPT changes were not calculated [55], or data were depicted in a graph only [35].

All but two studies (92.3%) assessed the effect of acupuncture on the HPT in healthy volunteers. Seventeen of these studies (70.8%) focused on the effect of EA only [30, 34, 36, 42, 53–55, 61, 66, 69, 80, 85, 90, 94, 101, 102, 105], five studies (20.8%) on the effect of MA only [31, 32, 35, 67, 99], and two studies (8.3%) evaluated both treatment methods. Of the 19 studies evaluating EA, 15 (78.9%) indicated an elevation of the HPT [30, 34, 53, 54, 60, 61, 66, 80, 85, 90, 94, 100–102, 105] such as did five of seven (71.4%) MA studies [31, 32, 60, 67, 100].

Price et al. assessed low back pain patients and found a significant effect of EA on the HPT [75]. Lundeberg et al., however, did not find any effect of MA or EA on the HPT neither in patients suffering from sinus pain nor in healthy volunteers [104].

The methods used to determine the HPT varied largely, and some studies evaluated several measures. In 16 out of 26 studies (61.5%), the HPT was defined as the time period that subjects were able to tolerate a defined heat stimulus. Eleven of these studies showed a positive outcome. Defined heat stimuli were produced by a projection lamp held on blackened skin [30, 34–36, 42, 53–55, 66, 69, 85, 90], by a hot plate [32, 67], or by a thermode containing a peltier element [31]. In one study no information was given about the method used for

the application of heat stimuli [94]. In six studies (23.1%) the HPT was defined as the temperature that was first experienced as painful when increasing heat was applied with a thermal sensory analyzer [99–102, 104, 105]. Results of three of these studies (50%) indicated an effect of acupuncture on the HPT [100, 102, 105]. Six of the 26 studies (23.1%) assessed the pain intensity evoked by a heat stimuli applied with a thermode [60, 61, 75, 80, 101, 102]. All six found a reduction of the pain intensity after acupuncture.

Cold Pain Threshold (CPT)

Of the 85 included studies, 17 (20%) addressed the effect of acupuncture on the CPT. In 11 of these 17 studies (64.7%), subjects were less sensitive to cold pain after than before acupuncture. All studies focused on the immediate effect after one single acupuncture treatment. Sixteen studies (94.1%) were conducted with healthy volunteers and one with chronic pain patients. When assessing healthy volunteers, eight studies (50%) showed a significant change of the CPT through EA [25, 57, 58, 71, 84, 92, 97, 105], while four studies (25%) found no effect of EA [56, 59, 101, 102]. Three studies (18.8%) conducted with healthy volunteers investigated the effect of MA. Two of these showed a desensitizing effect [24, 26] and one no effect of acupuncture [99]. Lang et al. assessed both, MA and EA, and did not find any effect on the CPT [100]. Ulett et al. showed that sensitivity to cold pain of chronic pain patients was decreased after EA [89].

Different methods were used for the CPT measurements. In ten studies (58.8%), the CPT was assessed by immersion of the subjects' fingers or forearm into ice-water and documenting either the time until subjects withdrew their limb [24, 26, 71] or the pain intensity that was experienced during a defined period of immersion into ice water [25, 56–59, 84, 89]. Eight of these ten studies (80%) showed a significant reduction of the CPT after acupuncture. In one study, contact to an ice cold cylinder was tolerated longer after than before acupuncture [105]. In six studies (35.3%), cold stimuli were applied with a thermode including a peltier element and either the intensity of the pain evoked by defined cold stimuli [97] or the temperature that was considered painful [92, 99–102] was evaluated. Four of the latter showed no changes of the CPT [99–102].

Thermal Detection Thresholds

Warm Detection Threshold (WDT)

The effect of acupuncture on the WDT was investigated in six studies (7.1%) with five evaluating the immediate effect in healthy subjects. Among these, significant changes of the WDT were shown in one study after MA [29] and in two studies after EA [101, 102]. No effect of acupuncture in healthy volunteers was reported in two studies, one on MA [99] and one assessing both MA and EA [100]. Ahn et al. included patients suffering from painful diabetic neuropathy (PDN) and measured the WDT before and after a series of treatments [106]. The results of

this study are rated as unclear due to the lack of a statistical analysis and the fact that data are depicted in a graph only.

Cold Detection Threshold (CDT)

Five of the six studies that investigated the effect of acupuncture on the WDT also assessed changes in the CDT after acupuncture. Correspondingly, results of one study assessing the CDT in PDN patients were rated as unclear [106]. Of the four remaining studies, three found no change of the CDT after MA [99, 100] or EA [101], and two found a decrease of the CDT after EA [100, 102].

All but one study, for which the measuring tool was not described, used a thermode including a peltier element to assess the WDT and the CDT.

Thermal Sensory Limen (TSL)

Three of the included acupuncture studies (3.5%) used the TSL as an outcome measure. All three investigated immediate treatment effects. In two studies by Lundeberg et al., the TSL was found unchanged after EA as well as after MA in healthy volunteers and in patients suffering from sinus pain [103, 104]. In contrast, Lang et al. found a significant elevation of the TSL of up to 3°C through EA but not through MA in healthy volunteers [100]. In all studies, TSL assessments were performed by using a sensory thermal analyzer. The thermode applied on the subjects' skin increases or decreases its temperature. As soon as the subject indicates the feeling of cool or warm, a switch towards the opposite direction of the temperature change is induced.

Pressure Pain Threshold (PPT)

Almost half of the articles (42 out of 85, 49.4%) included in this review describe changes of the PPT through acupuncture.

PPT assessed in Healthy Subjects

Seven of these 42 studies (16.7%) evaluated the immediate effect of acupuncture on the PPT in healthy subjects. All showed a statistically significant elevation of the PPT through acupuncture. Two studies assessed the effect of EA only [28, 39], three studies solely the effect of MA [65, 95, 98], and two the effect of both, MA and EA. Either a manual algometer [39, 65, 95, 100] according to Fischer et al. [107] or an electronic algometer [28, 76, 98] was used as measuring tool (tip size 1 cm²). Changes ranged from 59 to 392 kPa [28, 39, 76, 98, 100] or from 10% to 27%, respectively [65, 95].

PPT assessed in Pain Disorders

The assessment of the PPT was widely used (35 out of 42, 83.3%) in order to evaluate the effectiveness of acupuncture in reducing hyperalgesia in pain disorders. Most common conditions under investigation were myofascial pain, associated with the occurrence of MTrPs (8 out of 35, 22.9%) [33, 38, 41, 48, 50, 63, 70, 83], fibromyalgia (7 out of 35, 20%) [37, 44, 45, 72, 81, 82, 87], chronic neck and back pain (5 out of 35, 14.3%)

[22, 46, 49, 74, 77], temporomandibular disorder (5 out of 35, 14.3%) [40, 43, 68, 79, 91], and experimentally induced delayed-onset muscle soreness (DOMS; 4 out of 35, 11.4%) [27, 47, 51, 78]. PPT was also used as an outcome measure for the treatment effect of acupuncture in tension type headache [52, 93], osteoarthritis of the knee [86], abdominal scar pain [62], plantar fasciitis [96], and syringomyelia [64].

Ten of the 35 patient studies (28.6%) evaluated the immediate effect of one single acupuncture treatment. Nine of these reported the PPT to be significantly elevated [33, 40, 41, 43, 51, 64, 78, 79, 83], and one study found no significant PPT increase [50]. Among the 26 studies (74.3%) that assessed the effectiveness of an acupuncture series, 18 (69.2%) showed a PPT increase after treatment [37, 38, 46, 50, 52, 62, 63, 68, 70, 72, 74, 77, 81, 82, 86, 87, 91, 93] while six studies (23.1%) observed no effect of acupuncture on the PPT [22, 27, 47–49, 96]. The outcome of two studies was rated as unclear, because the analysis was performed by combining data of all groups [44, 45]. Although no statistics were provided, the outcome of two other studies was rated as positive. Singh et al. found that, after a MA treatment of two months, the majority of fibromyalgia patients felt less pain, although greater pressure than at baseline was applied [81]. Perez-Palomarez et al. included a large number of patients and showed a prominent mean increase of the PPT (74.5 kPa–202.0 kPa) after DN and EA [74].

The effect of EA on the PPT was evaluated in four out of the 35 studies (11.4%) conducted with patients [37, 64, 74, 93]. All four showed an increase of the PPT after EA. MA was investigated in 22 studies (62.9%) of which 17 (77.3%) revealed an increase of the PPT. In two of these studies, MA was applied as an add-on treatment to stretching [70] or tricyclic antidepressants and exercise [87] for myofascial pain or fibromyalgia, respectively. Seven of the 35 studies (20.0%) investigated the effects of DN in the treatment of myofascial pain. DN was shown to be effective in increasing the PPT in six studies [33, 38, 40, 41, 74, 83], but one study found no effect of DN on the PPT at MTrPs of the neck [48]. Four studies used a combination of either EA and MA [22, 46, 68] or of MA and DN [49]. Two of these studies showed a change of the PPT [46, 68] and two no treatment effect [22, 49]. The observed PPT changes varied largely between studies (22.5 kPa to 245.2 kPa) and tended to increase during follow up if assessed.

Three studies also assessed the pressure pain tolerance (PPTo) which was defined as the time subjects tolerated painful pressure [45, 64, 74]. Two studies showed the PPTo to be elevated after EA [64] and DN [74], respectively. Results of one study on MA were rated as unclear [44].

Study outcome was not associated with a certain methodology used for the PPT assessment. One study did not provide details about the measuring tool (spring roller) [64]. In all other studies (34 out of 35, 97.1%) algometers with different tip sizes (between 0.28 and 3.14 cm²) were used to determine the PPT. In 23 studies, algometers were equipped with a rubber tip of 1 cm² or larger [27, 33, 37, 38, 40, 41, 44–48, 50, 62, 68, 70, 74, 79, 81, 82, 86, 87, 91, 96], while in five studies tips of smaller sizes were used [49, 51, 52, 72, 83, 93]. In five articles no

information was given about the characteristics of the algometer [22, 43, 63, 77, 78].

Meta-Analysis

Considering the previously defined quality criteria and clinical homogeneity, only studies investigating the effect of acupuncture on the PPT were eligible for meta-analyses. Four studies could be combined in an analysis regarding the immediate treatment effect [22, 37, 49, 93], and two studies were analyzed regarding long-term effects of acupuncture in pain conditions [49, 52]. The results of these meta-analyses confirm our descriptive findings. The effect of acupuncture directly after a series of treatments (fig. 3A) and at follow up time points (6 weeks [52] and 3 months [49]; fig. 3B) was shown to be significant ($p < 0.05$). However, effects were small (SMD 0.26 [95% CI: 0.1; 0.41] for immediate effects, SMD 0.3 [95% CI: 0.02; 0.52] for long-term effects) and heterogeneity was found to be substantial.

Mechanical Pain Threshold/Sensitivity (MPT/MPS)

Three of the 85 articles (3.5%) included in this review investigated the effect of acupuncture on the MPT and/or the MPS. All three studies were conducted with healthy volunteers assessing the MPT/MPS before and immediately after a single acupuncture treatment. All showed a desensitizing effect of EA as well as of MA [73, 100, 105]. Just Lang et al., who assessed both, the MPT and the MPS, did not find an effect of MA on the MPT [100]. However, different measuring tools were used; scaled forces applied with pin-pricks [100], increasing pressure applied by a blunt needle [73], or gauged forceps [105].

Mechanical Detection Threshold

We identified two acupuncture studies (2.4%), in which the MDT was one of the outcome measures. Both studies were conducted with healthy volunteers, used Von Frey Filaments in order to evaluate the MDT, and focused on immediate effects of acupuncture; one on EA [102] and one on both EA and MA [100]. Results of these two studies indicate that neither EA nor MA has an impact on the MDT in healthy subjects.

Vibration Detection Threshold (VDT)

Four articles (4.7%) describe the effect of acupuncture on the VDT. In two studies the impact of a MA treatment series on the VDT in patients suffering from diabetic peripheral neuropathy was explored. Tong et al. noted an improvement of the ability to detect vibration [88] while Abuaisha et al. found no changes of the VDT after MA [23, 88]. Lang et al. and Lundeborg et al. reported that a single application of neither EA nor MA had an effect on the VDT in healthy volunteers [100, 103]. Vibration stimuli were applied with a Rydel-Seiffer tuning fork [100] or an electromechanical device [23, 88, 103].

Effect of Needle Location relative to the Sites of Measurements

Twenty (23.5%) studies compared ipsi- to contralateral and/or close to distant needling either in healthy subjects (15 out of 20; 75%) or in pain patients (5 out of 20; 25%). Of the ten studies comparing close to distant needling, eight (80%) showed a larger increase of at least one sensory threshold close to the needling location [35, 51, 53, 67, 83, 101, 102, 105]. In six out of eleven studies (54.5%), threshold changes were more pronounced after ipsi- than after contralateral needling [25, 34, 67, 100–102]. In one high quality study bilateral needle placement was superior to unilateral needling [65], and results of three studies suggest that needle stimulation at LI 4 is more effective in changing pain thresholds than needling at other acupuncture points [39, 53, 65]. Four studies (20%) found significant effects of acupuncture on sensory perception independent of the needle location [28, 60, 66, 75, 95], and in two studies no change of any sensory threshold was observed [103, 104].

Responder versus Non-responders

Six of all included studies (7.1%) distinguished between subjects that responded to acupuncture and those who did not [32, 35, 42, 55, 60, 94]. All of these studies assessed the HPT in healthy volunteers and all were conducted with relatively small case numbers (11.8 ± 3.0). Proportions of responders ranged from one third to two thirds. Chae et al. found genetic difference between acupuncture responders and non-responders [32], but their results have not been reproduced by further investigations. Furthermore, the role of hypnotic susceptibility in responsiveness to acupuncture was investigated in three studies, with contradictory results [56, 58, 89]. The influence of expectancy was found to be substantial in two studies [57, 61]. Knox et al. found no effect of EA neither in oriental nor occidental subjects [59].

Discussion

Result Interpretation

Our results revealed that in 76.5% of 85 eligible studies at least one sensory threshold was changed after acupuncture, indicating an activation of neuromodulatory mechanisms. However, results displayed substantial heterogeneity, which is illustrated for the PPT by results of the meta-analyses (fig. 3).

Over half of the sham-controlled studies found larger effects in the verum group than in the sham group. However, a quarter of these studies found significant threshold changes also after sham acupuncture, while only one of the included articles reported changes of sensory thresholds after an inert control procedure. These facts go in line with the previously drawn conclusion that there are effects specific to acupuncture, but that sham acupuncture may cause physiological reactions exceeding pure placebo effects [6, 108, 109]. It can be

assumed that this in part explains the clinical effects of sham acupuncture interventions observed in acupuncture randomized controlled trials.

Most studies conducted with patients used the PPT as an outcome measure. This reflects the frequent use of acupuncture in clinical practice for treating pain conditions e.g. musculoskeletal disorders, in which the PPT correlates well with clinical status [110]. More than 80% of the studies – i.e. 27 out of 35 clinical studies and all seven studies conducted with healthy volunteers – showed that acupuncture reduced pain evoked by blunt pressure which is mainly mediated by deep tissue nociceptors (A δ - and C-fibers) [110]. PPT reductions of up to 245.2 kPa as observed in some studies can be interpreted as clinically relevant. In addition, results of two meta-analyses show significant short- and long-term effects of acupuncture on the PPT in pain conditions (fig. 3). Thus, these findings provide a physiological basis for the growing body of evidence for the effectiveness of acupuncture in locomotor conditions associated with tenderness [111–113].

Pain thresholds were elevated after acupuncture also when painful pressure was exerted on a rather small skin area (≤ 1 cm²) and when pin-prick like stimuli (MPS and MPT) were applied. This finding is derived from few studies, but suggests that acupuncture also affects mechanical pain evoked by punctate objects which is primarily mediated by intra-epidermal nociceptors (mainly A- δ fibers) [114].

Studies investigating whether thermal pain is reduced through acupuncture are abundant but almost exclusively conducted with healthy volunteers. Results of such investigations are more ambiguous than data of included studies on changes of the PPT, MPT and MPS. After acupuncture, sensitivity to painful heat was reduced in 19 out of 27 studies, while sensitivity to painful cold was reduced in 11 out of 17 experiments. The transmission of heat pain is mainly evoked by C-fiber mechano-heat nociceptors (CMH) responding to heat stimuli ranging from 41 °C to 49 °C [115] and is linked to the capsaicin sensitive vanilloid receptor VR1 which is also found in type II A δ -nociceptors [116]. In contrast, the transmission of cold pain is mediated by both cold sensitive C- and A δ -fibers [117, 118] which are insensitive to vanilloid compounds [119]. Our results suggest that both types of nociception are likely to be affected by acupuncture in healthy subjects. Reasons for the heterogeneity of data might partly be explained by methodological issues. Studies in which painful cold was applied through a thermode found changes of the CPT less frequently (two versus four) than other studies (eight versus two), in which e.g. the subject's hand or arm was immersed into ice-water. It is conceivable that differences in study outcome are related to the size of the skin area to which cold stimuli were applied or to the intensity of these stimuli (see 4.2. *Limitations* for further discussion).

In contrast to pain perception, data on the effect of acupuncture on sensory detection are sparse. The ability to detect temperature changes (WDT, CDT, TSL) was reduced after acupuncture in half of all experiments. Mechanical detection (VDT and MDT), in contrast, was not affected by acupuncture in five out of six trials. These findings provide a first hint that acupuncture might not affect mechanical detection which is mainly mediated by A β -fiber signaling [120, 121],

while the influence on thermal detection which is linked to signaling of warm sensitive C-fibers or cold sensitive A δ - and C-fibers [122, 123] remains unclear.

Another important finding is that ipsilateral needling and needling close to the measure sites were found to exhibit stronger effects on sensory thresholds than needling at contralateral or remote body sites, respectively. This underscores the importance of local mechanisms such as the release of neuromodulators at the needling site [14] and spinal mechanisms such as segmental inhibition [11]. It is supposed, that activation of A-fiber afferents results in the activation of spinal inhibitory interneurons, those achieving primary analgesia within the same segment [124, 125]. Nevertheless, more studies adopting a sophisticated selection of measure sites in combination with imaging studies are needed to clearly differentiate between local, spinal and supraspinal mechanisms of acupuncture. In contrast, it seems to be of minor importance whether the needle is stimulated electrically or manually.

Limitations

The overall poor study quality (fig. 2) and the consequently low number of studies included in the meta-analyses are the major limitations to our findings. However, limitations resulting from the quality assessment itself need to be taken into account. Many publications were characterized by poor reporting which renders an estimation of the real number of high quality studies/studies with low risk of bias impossible. In particular details of blinding and randomization procedures as well as treatment regimens were often missing. Second, the STRICTA guidelines provide an essential tool for assuring the quality of reporting in acupuncture trials. Nevertheless, it bears the striking disadvantage that the quality of reporting of treatment related items might be underestimated in studies investigating the effectiveness of individualized acupuncture regimens since precise instructions for such studies are missing.

A further limitation comes with the methodology of QST itself. Although when performed with standardized methods, QST involves subjective ratings of perceptions and, therefore, is susceptible to bias due to psychological factors such as expectation and conditioning [126]. These should be assessed and/or controlled for [127]. Comparability of studies was also limited due to methodological variability with regard to treatment, study population, and outcome assessment. Appropriate selection of the acupuncture points needled, number of needles and stimulation technique are, according to the traditional concept of acupuncture, crucial for achieving an optimal treatment effect. These concepts are in part supported by research findings. For example, current evidence suggests an association between the number of needles applied and the clinical outcome [128], but overall there is still no consensus within the scientific community about how to appraise these parameters correctly. The same holds for the socio-cultural background of the study population. Previous acupuncture experience and social valuation of acupuncture is very likely to have an impact on treatment outcome; likewise to any other treatment. Yet, there is very little emphasis on sociocultural

aspects in acupuncture research. There were striking differences between acupuncture interventions applied in the included studies and, although not reported in detail, very probably also between the different study populations. However, due to the still poor knowledge about the impact of these factors, there are no guidelines on how to appraise them in a review apart from pure reporting as performed in [table 1](#) and [2](#).

Additionally, study outcome might in part be influenced by the measuring tool. Investigations comparing different methods evaluating the same sensory thresholds are missing. It also remains unclear to what extent the test stimuli might interfere with the treatment effect. For example, strong noxious stimulation is known to activate aspects of the endogenous pain-modulating network [[129–131](#)]; a phenomenon known as counter irritation. It is postulated that the main underlying mechanisms namely DNIC is also involved in the analgesic effect of acupuncture [[98, 132–134](#)]. This is however controverted by recent work and needs to be further explored [[98, 134](#)].

Furthermore, the interpretation of the clinical relevance of the effect of acupuncture is limited by the fact that studies - besides those assessing the PPT - were almost exclusively conducted with healthy volunteers. There is also limited amount of data encoding the effect of acupuncture on stimuli above pain thresholds, which have been shown to activate other nerve fibers than stimuli around the pain threshold [[135](#)].

Future Perspectives

Recently, efforts are increasing to comprehensively characterize painful conditions by means of QST [[136–139](#)]. Attempts are made to classify patients on the basis of symptoms, signs, or patterns of somatosensory abnormalities [[139–141](#)]. Those might reflect the underlying pathological mechanisms [[142](#)] and might, therefore, be related to different treatment responses [[137, 143](#)]. Such subgroup analysis should also be adopted in acupuncture research since responsiveness to acupuncture varies largely between individuals. This inter-individual variability in response to acupuncture was also demonstrated by studies included in this review, but proposed hypotheses explaining this phenomenon have not been verified. Comprehensive QST assessments applied in large scale studies might provide hints about whether individual patterns of somatosensory alterations might be of predictive value for the effectiveness of acupuncture. It is striking that only one study applied a comprehensive, standardized QST test battery [[100](#)].

Comprehensive QST assessments might, furthermore, provide information about the extent to which signs of peripheral and central sensitization may be reduced by acupuncture. For example, it has not yet been subject to clinical trials whether thermal and mechanical hypaesthesia or increased wind-up, which can be associated with chronic pain conditions [[144–146](#)], might vanish along with clinical improvement after an acupuncture therapy.

Conclusion

This review provides evidence for the effect of acupuncture on sensory perception, especially pain perception. Most compelling evidence supports the reduction of pain evoked by blunt pressure through acupuncture; particularly as a measure of tenderness in pain conditions. Moderate evidence was found that acupuncture reduces the perception of noxious heat or cold. The outcome of these studies seemed to depend on measuring methods. Little but consistent evidence was found that pin-prick like pain (MPT, MPS) is reduced but that mechanical detection (VDT, MDT) is not influenced by acupuncture. No conclusions can be drawn about whether acupuncture affects the ability to detect temperature changes.

Sham-acupuncture approaches can evoke such effects as well. Thus, it is questionable to classify them as pure placebos, a fact that needs to be taken into account when conducting and interpreting acupuncture studies.

Data support the importance of local and spinal mechanisms involved in the neurophysiological effect of acupuncture. More high quality studies are needed to characterize the effect of acupuncture on the whole sensory profile by means of comprehensive QST assessments. In addition, investigating defined patient populations accordingly might also clarify whether certain characteristics of the somatosensory profile are of predictive value for the analgesic effect of acupuncture and to which extent pathologic alterations of sensory perception vanish along with clinical improvement achieved by acupuncture.

Supporting Information

S1 Table. PRISMA Checklist. Overview of the reporting items in accordance with the PRISMA statement

[doi:10.1371/journal.pone.0113731.s001](https://doi.org/10.1371/journal.pone.0113731.s001) (DOC)

Acknowledgments

We thank Mrs. Franziska Benedikt for her support when performing the data extraction.

Author Contributions

Conceived and designed the experiments: PIB JF ST DI TS MS. Performed the experiments: PIB JF ST. Analyzed the data: PIB JF DI MS ST. Contributed reagents/materials/analysis tools: PIB JF DI MS. Wrote the paper: PIB JF ST DI TS MS.

References

1. **Barnes PM, Bloom B, Nahin RL** (2008) Complementary and alternative medicine use among adults and children: United States, 2007. *Natl Health Stat Report*: 1–23.
2. **Eardley S, Bishop FL, Prescott P, Cardini F, Brinkhaus B, et al.** (2012) A Systematic Literature Review of Complementary and Alternative Medicine Prevalence in EU. *Forsch Komplementmed* 19 Suppl 2: 18–28.
3. **Jones L, Othman M, Dowswell T, Alfirevic Z, Gates S, et al.** (2012) Pain management for women in labour: an overview of systematic reviews. *Cochrane Database Syst Rev* 3: CD009234.
4. **Lee JH, Choi TY, Lee MS, Lee H, Shin BC** (2013) Acupuncture for acute low back pain: a systematic review. *Clin J Pain* 29: 172–185.
5. **Smith CA, Zhu X, He L, Song J** (2011) Acupuncture for primary dysmenorrhoea. *Cochrane Database Syst Rev*: CD007854.
6. **Vickers AJ, Cronin AM, Maschino AC, Lewith G, MacPherson H, et al.** (2012) Acupuncture for chronic pain: individual patient data meta-analysis. *Arch Intern Med* 172: 1444–1453.
7. **Pittler MH, Ernst E** (2008) Complementary therapies for neuropathic and neuralgic pain: systematic review. *Clin J Pain* 24: 731–733.
8. **Langhorst J, Klose P, Musial F, Irnich D, Hauser W** (2010) Efficacy of acupuncture in fibromyalgia syndrome—a systematic review with a meta-analysis of controlled clinical trials. *Rheumatology (Oxford)* 49: 778–788.
9. **Huang W, Pach D, Napadow V, Park K, Long X, et al.** (2012) Characterizing acupuncture stimuli using brain imaging with fMRI—a systematic review and meta-analysis of the literature. *PLoS One* 7: e32960.
10. **Lin JG, Chen WL** (2008) Acupuncture analgesia: a review of its mechanisms of actions. *Am J Chin Med* 36: 635–645.
11. **Zhao ZQ** (2008) Neural mechanism underlying acupuncture analgesia. *Prog Neurobiol* 85: 355–375.
12. **Han JS** (2004) Acupuncture and endorphins. *Neurosci Lett* 361: 258–261.
13. **Zhang Y, Zhang RX, Zhang M, Shen XY, Li A, et al.** (2012) Electroacupuncture inhibition of hyperalgesia in an inflammatory pain rat model: involvement of distinct spinal serotonin and norepinephrine receptor subtypes. *Br J Anaesth* 109: 245–252.
14. **Goldman N, Chen M, Fujita T, Xu Q, Peng W, et al.** (2010) Adenosine A1 receptors mediate local anti-nociceptive effects of acupuncture. *Nat Neurosci* 13: 883–888.
15. **Arendt-Nielsen L, Yarnitsky D** (2009) Experimental and clinical applications of quantitative sensory testing applied to skin, muscles and viscera. *J Pain* 10: 556–572.
16. **Dyck PJ, O'Brien PC, Kosanke JL, Gillen DA, Karnes JL** (1993) A 4, 2, and 1 stepping algorithm for quick and accurate estimation of cutaneous sensation threshold. *Neurology* 43: 1508–1512.
17. **Dyck PJ, Zimmerman I, Gillen DA, Johnson D, Karnes JL, et al.** (1993) Cool, warm, and heat-pain detection thresholds: testing methods and inferences about anatomic distribution of receptors. *Neurology* 43: 1500–1508.
18. **Rolke R, Baron R, Maier C, Tolle TR, Treede RD, et al.** (2006) Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values. *Pain* 123: 231–243.
19. **Moher D, Liberati A, Tetzlaff J, Altman DG** (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS medicine* 6: e1000097.
20. **Higgins JPT, Altman DG, Sterne JAC, Cochrane Statistical Methods Group** (2011) *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0. - Chapter 8: Assessing risk of bias in included studies. Available: www.cochrane-handbook.org Accessed 2013 November 12.
21. **MacPherson H, Altman DG, Hammerschlag R, Youping L, Taixiang W, et al.** (2010) Revised Standards for Reporting Interventions in Clinical Trials of Acupuncture (STRICTA): extending the CONSORT statement. *PLoS medicine* 7: e1000261.
22. **Zhu XM, Polus B** (2002) A controlled trial on acupuncture for chronic neck pain. *American Journal of Chinese Medicine* 30: 13–28.

23. **Abuaisha BB, Costanzi JB, Boulton AJM** (1998) Acupuncture for the treatment of chronic painful peripheral diabetic neuropathy: A long-term study. *Diabetes Research and Clinical Practice* 39: 115–121.
24. **Amand M, Nguyen-Huu F, Balestra C** (2011) Acupuncture effect on thermal tolerance and electrical pain threshold: A randomised controlled trial. *Acupuncture in Medicine* 29: 47–50.
25. **Anderson DG, Jamieson JL, Man SC** (1974) Analgesic effects of acupuncture on the pain of ice water: a double blind study. *CANAD.J.PSYCHOL.* 28: 239–244.
26. **Ashton H, Ebenezer I, Golding JF, Thompson JW** (1984) Effects of acupuncture and transcutaneous electrical nerve stimulation on cold-induced pain in normal subjects. *Journal of Psychosomatic Research* 28: 301–308.
27. **Barlas P, Robinson J, Allen J, Baxter GD** (2000) Lack of effect of acupuncture upon signs and symptoms of delayed onset muscle soreness. *Clin Physiol* 20: 449–456.
28. **Barlas P, Ting SL, Chesterton LS, Jones PW, Sim J** (2006) Effects of intensity of electroacupuncture upon experimental pain in healthy human volunteers: a randomized, double-blind, placebo-controlled study. *Pain* 122: 81–89.
29. **Benoliel R, Zaidan S, Eliav E** (2011) Acupuncture modulates facial warm sensory thresholds. *Journal of orofacial pain* 25: 32–38.
30. **Berlin FS, Bartlett R, Black JD** (1975) Acupuncture and placebo: Effects on delaying the terminating response to a painful stimulus. *Anesthesiology* 42: 527–531.
31. **Brockhaus A, Elger CE** (1990) Hypalgesic efficacy of acupuncture on experimental pain in man. Comparison of laser acupuncture and needle acupuncture. *Pain* 43: 181–185.
32. **Chae Y, Park HJ, Hahm DH, Yi SH, Lee H** (2006) Individual differences of acupuncture analgesia in humans using cDNA microarray. *J Physiol Sci* 56: 425–431.
33. **Chou LW, Hsieh YL, Chen HS, Hong CZ, Kao MJ, et al.** (2011) Remote therapeutic effectiveness of acupuncture in treating myofascial trigger point of the upper trapezius muscle. *Am J Phys Med Rehabil* 90: 1036–1049.
34. **Clark WC, Yang JC** (1974) Acupunctural analgesia? Evaluation by signal detection theory. *Science* 184: 1096–1098.
35. **Croze S, Antonietti C, Duclaux R** (1976) Changes in burning pain threshold induced by acupuncture in man. *Brain Research* 104: 335–340.
36. **Day RL, Kitahata LM, Kao FF** (1975) Evaluation of acupuncture anesthesia: a psychophysical study. *Anesthesiology* 43: 507–517.
37. **Deluze C, Bosia L, Zirbs A, Chantraine A, Vischer TL** (1992) Electroacupuncture in fibromyalgia: results of a controlled trial. *BMJ* 305: 1249–1252.
38. **Edwards J, Knowles N** (2003) Superficial Dry Needling and Active Stretching in the Treatment of Myofascial Pain - A Randomised Controlled Trial. *Acupuncture in Medicine* 21: 80–86.
39. **Farber PL, Tachibana A, Campiglia HM** (1997) Increased pain threshold following electroacupuncture: Analgesia is induced mainly in meridian acupuncture points. *Acupuncture and Electro-Therapeutics Research* 22: 109–117.
40. **Fernandez-Carnero J, La Touche R, Ortega-Santiago R, Galan-del-Rio F, Pesquera J, et al.** (2010) Short-term effects of dry needling of active myofascial trigger points in the masseter muscle in patients with temporomandibular disorders. *J Orofac Pain* 24: 106–112.
41. **Fu ZH, Wang JH, Sun JH, Chen XY, Xu JG** (2007) Fu's subcutaneous needling: Possible clinical evidence of the subcutaneous connective tissue in acupuncture. *Journal of Alternative and Complementary Medicine* 13: 47–51.
42. **Galloon S, Evans RJ** (1977) Acupuncture analgesia. *Deutsches Arzteblatt* 74: 879–880.
43. **Goddard G, Karibe H, McNeill C, Villafuerte E** (2002) Acupuncture and sham acupuncture reduce muscle pain in myofascial pain patients. *Journal of Orofacial Pain* 16: 71–76.
44. **Harris RE, Gracely RH, McLean SA, Williams DA, Giesecke T, et al.** (2006) Comparison of Clinical and Evoked Pain Measures in Fibromyalgia. *Journal of Pain* 7: 521–527.

45. **Harris RE, Sundgren PC, Pang Y, Hsu M, Petrou M, et al.** (2008) Dynamic levels of glutamate within the insula are associated with improvements in multiple pain domains in fibromyalgia. *Arthritis and Rheumatism* 58: 903–907.
46. **He D, Veiersted KB, Hostmark AT, Medbo JI** (2004) Effect of acupuncture treatment on chronic neck and shoulder pain in sedentary female workers: a 6-month and 3-year follow-up study. *Pain* 109: 299–307.
47. **Hubscher M, Vogt L, Bernhorster M, Rosenhagen A, Banzer W** (2008) Effects of acupuncture on symptoms and muscle function in delayed-onset muscle soreness. *J Altern Complement Med* 14: 1011–1016.
48. **Ibuldu E, Cakmak A, Disci R, Aydin R** (2004) Comparison of laser, dry needling, and placebo laser treatments in myofascial pain syndrome. *Photomedicine and Laser Surgery* 22: 306–311.
49. **Irnich D, Behrens N, Molzen H, Konig A, Gleditsch J, et al.** (2001) Randomised trial of acupuncture compared with conventional massage and "sham" laser acupuncture for treatment of chronic neck pain. *BMJ* 322: 1574–1578.
50. **Irnich D, Karg H, Behrens N, Lang PM, Schreiber MA, et al.** (2003) Controlled trial on point specificity of acupuncture in the treatment of lateral epicondylitis (Tennis elbow). *Physikalische Medizin Rehabilitationsmedizin Kurortmedizin* 13: 215–219.
51. **Itoh K, Minakawa Y, Kitakoji H** (2011) Effect of acupuncture depth on muscle pain. *Chinese Medicine* 6.
52. **Karst M, Rollnik JD, Fink M, Reinhard M, Piepenbrock S** (2000) Pressure pain threshold and needle acupuncture in chronic tension-type headache - A double-blind placebo-controlled study. *Pain* 88: 199–203.
53. **Kitade T, Hyodo M** (1979) The effects of stimulation of ear acupuncture points on the body's pain threshold. *American Journal of Chinese Medicine* 7: 241–252.
54. **Kitade T, Minamikawa M, Nawata T** (1981) An experimental study on the enhancing effects of phenylalanine on acupuncture analgesia. *American Journal of Chinese Medicine* 9: 243–248.
55. **Kitade T, Odahara Y, Shinohara S, Ikeuchi T, Sakai T, et al.** (1988) Studies on the enhanced effect of acupuncture analgesia and acupuncture anesthesia by D-phenylalanine (first report)—effect on pain threshold and inhibition by naloxone. *Acupunct Electrother Res* 13: 87–97.
56. **Knox VJ, Gekoski WL, Shum K, McLaughlin DM** (1981) Analgesia for experimentally induced pain: Multiple sessions of acupuncture compared to hypnosis in high- and low-susceptible subjects. *Journal of Abnormal Psychology* 90: 28–34.
57. **Knox VJ, Handfield-Jones CE, Shum K** (1979) Subject expectancy and the reduction of cold pressor pain with acupuncture and placebo acupuncture. *Psychosomatic Medicine* 41: 477–486.
58. **Knox VJ, Shum K** (1977) Reduction of cold-pressor pain with acupuncture analgesia in high- and low-hypnotic subjects. *Journal of Abnormal Psychology* 86: 639–643.
59. **Knox VJ, Shum K, McLaughlin DM** (1977) Response to cold pressor pain and to acupuncture analgesia in oriental and occidental subjects. *Pain* 4: 49–57.
60. **Kong J, Fufa DT, Gerber AJ, Rosman IS, Vangel MG, et al.** (2005) Psychophysical outcomes from a randomized pilot study of manual, electro, and sham acupuncture treatment on experimentally induced thermal pain. *Journal of Pain* 6: 55–64.
61. **Kong J, Kaptchuk TJ, Polich G, Kirsch I, Vangel M, et al.** (2009) Expectancy and treatment interactions: A dissociation between acupuncture analgesia and expectancy evoked placebo analgesia. *NeuroImage* 45: 940–949.
62. **Kotani N, Kushikata T, Suzuki A, Hashimoto H, Muraoka M, et al.** (2001) Insertion of intradermal needles into painful points provides analgesia for intractable abdominal scar pain. *Regional Anesthesia and Pain Medicine* 26: 532–538.
63. **Kummerddee W** (2009) Effectiveness comparison between Thai traditional massage and Chinese acupuncture for myofascial back pain in Thai military personnel: a preliminary report. *Journal of the Medical Association of Thailand = Chotmaihet thangphaet* 92 Suppl 1: S117–123.
64. **Li S, Jiang C, Chen G** (1983) The relationship between needling sensation and acupuncture effects, with special reference to their ascending pathway in the spinal cord. *Acupuncture and Electro-Therapeutics Research* 8: 105–110.

65. **Li W, Cobbin D, Zaslowski C** (2008) A comparison of effects on regional pressure pain threshold produced by deep needling of L14 and L111, individually and in combination. *Complementary Therapies in Medicine* 16: 278–287.
66. **Lim TW, Loh T, Kranz H, Scott D** (1977) Acupuncture—effect on normal subjects. *Med J Aust* 1: 440–442.
67. **Lin MT, Chandra A, Chen-Yen SM** (1981) Effects of needle stimulation of acupuncture loci Nei-Kuan (EH-6), Tsu-San-Li (St-36), San-Yin-Chiao (Sp-6) and Chu-Chih (LI-11) on cutaneous temperature and pain threshold in normal adults. *Am J Chin Med* 9: 305–314.
68. **List T, Helkimo M, Karlsson R** (1993) Pressure pain thresholds in patients with craniomandibular disorders before and after treatment with acupuncture and occlusal splint therapy: a controlled clinical study. *Journal of Orofacial Pain* 7: 275–282.
69. **Lloyd MA, Wagner MK** (1976) Acupuncture analgesia and radiant heat pain: a signal detection analysis. *Anesthesiology* 44: 147–150.
70. **Ma C, Wu S, Li G, Xiao X, Mai M, et al.** (2010) Comparison of miniscalpel-needle release, acupuncture needling, and stretching exercise to trigger point in myofascial pain syndrome. *Clinical Journal of Pain* 26: 251–257.
71. **Moret V, Forster A, Laverriere MC, Lambert H, Gaillard RC, et al.** (1991) Mechanism of analgesia induced by hypnosis and acupuncture: is there a difference? *Pain* 45: 135–140.
72. **Nabeta T, Kawakita K** (2002) Relief of chronic neck and shoulder pain by manual acupuncture to tender points - A sham-controlled randomized trial. *Complementary Therapies in Medicine* 10: 217–222.
73. **Pauser G, Reichmann C, Baum M** (1975) The influence of acupuncture analgesia on the sensation and feeling of pain and the autonomic state of the organism. *Wiener Klinische Wochenschrift* 87: 25–28.
74. **Perez-Palomares S, Oliván-Blázquez B, Magallon-Botaya R, De-La-Torre-Beldarrain MML, Gaspar-Calvo E, et al.** (2010) Percutaneous electrical nerve stimulation versus dry needling: Effectiveness in the treatment of chronic low back pain. *Journal of Musculoskeletal Pain* 18 (1): 23–30.
75. **Price DD, Rafii A, Watkins LR, Buckingham B** (1984) A psychophysical analysis of acupuncture analgesia. *Pain* 19: 27–42.
76. **Schliessbach J, Van Der Klift E, Arendt-Nielsen L, Curatolo M, Streitberger K** (2011) The Effect of Brief Electrical and Manual Acupuncture Stimulation on Mechanical Experimental Pain. *Pain Medicine* 12: 268–275.
77. **Seidel U, Uhlemann C** (2002) A randomised controlled double-blind trial comparing dosed lasertherapy on acupuncture points and acupuncture for chronic cervical syndrome. *Deutsche Zeitschrift für Akupunktur* 45: 258–269.
78. **Shen YF, Goddard G** (2007) The short-term effects of acupuncture on myofascial pain patients after clenching. *Pain Practice* 7: 256–264.
79. **Shen YF, Younger J, Goddard G, Mackey S** (2009) Randomized clinical trial of acupuncture for myofascial pain of the jaw muscles. *Journal of Orofacial Pain* 23: 353–359.
80. **Shukla S, Torossian A, Duann JR, Leung A** (2011) The analgesic effect of electroacupuncture on acute thermal pain perception—a central neural correlate study with fMRI. *Mol Pain* 7: 45.
81. **Singh BB, Wu WS, Hwang SH, Khorsan R, Der-Martirosian C, et al.** (2006) Effectiveness of acupuncture in the treatment of fibromyalgia. *Alternative Therapies in Health and Medicine* 12: 34–41.
82. **Sprott H, Jeschonneck M, Grohmann G, Hein G** (2000) Changes in microcirculation above tender points in patients with fibromyalgia mediated by acupuncture. *Wiener Klinische Wochenschrift* 112: 580–586.
83. **Srbely JZ, Dickey JP, Lee D, Lowerison M** (2010) Dry needle stimulation of myofascial trigger points evokes segmental anti-nociceptive effects. *Journal of Rehabilitation Medicine* 42: 463–468.
84. **Stern JA, Brown M, Ulett GA, Sletten I** (1977) A comparison of hypnosis, acupuncture, morphine, valium, aspirin, and placebo in the management of experimentally induced pain. *Annals of the New York Academy of Sciences* 296: 175–193.
85. **Stewart D, Thomson J, Oswald I** (1977) Acupuncture analgesia: an experimental investigation. *British Medical Journal* 1: 67–70.

86. **Takeda W, Wessel J** (1994) Acupuncture for the treatment of pain of osteoarthritic knees. *Arthritis Care and Research* 7: 118–122.
87. **Targino RA, Imamura M, Kaziyama HH, Souza LP, Hsing WT, et al.** (2008) A randomized controlled trial of acupuncture added to usual treatment for fibromyalgia. *J Rehabil Med* 40: 582–588.
88. **Tong Y, Guo H, Han B** (2010) Fifteen-day Acupuncture Treatment Relieves Diabetic Peripheral Neuropathy. *JAMS Journal of Acupuncture and Meridian Studies* 3: 95–103.
89. **Ulett GA, Parwatikar SD, Stern JA, Brown M** (1978) Acupuncture, hypnosis and experimental pain-II. Study with patients. *Acupuncture and Electro-Therapeutics Research* 3: 191–201.
90. **Umino M, Shimada M, Kubota Y** (1984) Effects of acupuncture anesthesia on the pituitary gland. *The Bulletin of Tokyo Medical and Dental University* 31: 93–98.
91. **Vicente-Barrero M, Si-Lei YL, Bingxin Z, Bocanegra-Perez S, Duran-Moreno D, et al.** (2012) The efficacy of acupuncture and decompression splints in the treatment of temporomandibular joint pain-dysfunction syndrome. *Med Oral Patol Oral Cir Bucal*.
92. **Wang SM, Lin EC, Maranets I, Kain ZN** (2009) The impact of asynchronous electroacupuncture stimulation duration on cold thermal pain threshold. *Anesth Analg* 109: 932–935.
93. **Xue CCL, Dong L, Polus B, English RA, Zheng Z, et al.** (2004) Electroacupuncture for Tension-type Headache on Distal Acupoints only: A Randomized, Controlled, Crossover Trial. *Headache* 44: 333–341.
94. **Yoon SH, Koga Y, Matsumoto I, Ikezono E** (1986) Clinical study of objective pulse diagnosis. *Am J Chin Med* 14: 179–183.
95. **Zaslowski CJ, Cobbin D, Lidums E, Petocz P** (2003) The impact of site specificity and needle manipulation on changes to pain pressure threshold following manual acupuncture: A controlled study. *Complementary Therapies in Medicine* 11: 11–21.
96. **Zhang SP, Yip TP, Li QS** (2009) Acupuncture Treatment for Plantar Fasciitis: A Randomized Controlled Trial with Six Months Follow-up. *Evid Based Complement Alternat Med*.
97. **Zhang WT, Jin Z, Huang J, Zhang L, Zeng YW, et al.** (2003) Modulation of cold pain in human brain by electric acupoint stimulation: evidence from fMRI. *Neuroreport* 14: 1591–1596.
98. **Schliessbach J, Van Der Klift E, Siegenthaler A, Arendt-Nielsen L, Curatolo M, et al.** (2012) Does acupuncture needling induce analgesic effects comparable to diffuse noxious inhibitory controls? *Evidence-based Complementary and Alternative Medicine Article ID:785613*.
99. **Downs NM, Kirk K, MacSween A** (2005) The effect of real and sham acupuncture on thermal sensation and thermal pain thresholds. *Archives of Physical Medicine and Rehabilitation* 86: 1252–1257.
100. **Lang PM, Stoer J, Schober GM, Audette JF, Irnich D** (2010) Bilateral acupuncture analgesia observed by quantitative sensory testing in healthy volunteers. *Anesthesia and Analgesia* 110: 1448–1456.
101. **Leung A, Khadivi B, Duann JR, Cho ZH, Yaksh T** (2005) The effect of Ting point (tendinomuscular meridians) electroacupuncture on thermal pain: A model for studying the neuronal mechanism of acupuncture analgesia. *Journal of Alternative and Complementary Medicine* 11: 653–661.
102. **Leung AY, Kim SJ, Schulteis G, Yaksh T** (2008) The effect of acupuncture duration on analgesia and peripheral sensory thresholds. *BMC Complementary and Alternative Medicine* 8.
103. **Lundeberg T, Eriksson S, Lundeberg S, Thomas M** (1989) Acupuncture and sensory thresholds. *American Journal of Chinese Medicine* 17: 99–110.
104. **Lundeberg T, Laurell G, Thomas M** (1988) Effect of acupuncture on sinus pain and experimentally induced pain. *Ear, Nose and Throat Journal* 67: 565–566+571.
105. **Lynn B, Perl ER** (1977) Failure of acupuncture to produce localized analgesia. *Pain* 3: 339–351.
106. **Ahn AC, Bennani T, Freeman R, Hamdy O, Kaptchuk TJ** (2007) Two styles of acupuncture for treating painful diabetic neuropathy—a pilot randomised control trial. *Acupunct Med* 25: 11–17.
107. **Fischer AA** (1987) Pressure algometry over normal muscles. Standard values, validity and reproducibility of pressure threshold. *Pain* 30: 115–126.
108. **Irnich D, Salih N, Offenbacher M, Fleckenstein J** (2011) Is sham laser a valid control for acupuncture trials? *Evid Based Complement Alternat Med* 2011: 485945.

109. **Linde K, Niemann K, Schneider A, Meissner K** (2010) How large are the nonspecific effects of acupuncture? A meta-analysis of randomized controlled trials. *BMC Med* 8: 75.
110. **Treede RD, Rolke R, Andrews K, Magerl W** (2002) Pain elicited by blunt pressure: neurobiological basis and clinical relevance. *Pain* 98: 235–240.
111. **Linde K, Allais G, Brinkhaus B, Manheimer E, Vickers A, et al.** (2009) Acupuncture for tension-type headache. *Cochrane Database Syst Rev*: CD007587.
112. **Manheimer E, Cheng K, Linde K, Lao L, Yoo J, et al.** (2010) Acupuncture for peripheral joint osteoarthritis. *Cochrane Database Syst Rev*: CD001977.
113. **Vickers AJ, Cronin AM, Maschino AC, Lewith G, Macpherson H, et al.** (2012) Acupuncture for Chronic Pain: Individual Patient Data Meta-analysis. *Arch Intern Med*: 1–10.
114. **Garnsworthy RK, Gully RL, Kenins P, Mayfield RJ, Westerman RA** (1988) Identification of the physical stimulus and the neural basis of fabric-evoked prickle. *J Neurophysiol* 59: 1083–1097.
115. **LaMotte RH, Campbell JN** (1978) Comparison of responses of warm and nociceptive C-fiber afferents in monkey with human judgments of thermal pain. *J Neurophysiol* 41: 509–528.
116. **Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, et al.** (1997) The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 389: 816–824.
117. **Yarnitsky D, Ochoa JL** (1991) Warm and cold specific somatosensory systems. Psychophysical thresholds, reaction times and peripheral conduction velocities. *Brain* 114 (Pt 4): 1819–1826.
118. **Verdugo R, Ochoa JL** (1992) Quantitative somatosensory thermotest. A key method for functional evaluation of small calibre afferent channels. *Brain* 115 (Pt 3): 893–913.
119. **Caterina MJ, Leffler A, Malmberg AB, Martin WJ, Trafton J, et al.** (2000) Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science* 288: 306–313.
120. **Ochoa J, Torebjork E** (1983) Sensations evoked by intraneural microstimulation of single mechanoreceptor units innervating the human hand. *J Physiol* 342: 633–654.
121. **Wiesenfeld-Hallin Z, Hallin RG, Persson A** (1984) Do large diameter cutaneous afferents have a role in the transmission of nociceptive messages? *Brain Res* 311: 375–379.
122. **Schmelz M, Schmidt R** (2010) Microneurographic single-unit recordings to assess receptive properties of afferent human C-fibers. *Neurosci Lett* 470: 158–161.
123. **Darian-Smith I, Johnson KO, Dykes R** (1973) "Cold" fiber population innervating palmar and digital skin of the monkey: responses to cooling pulses. *J Neurophysiol* 36: 325–346.
124. **Le Bars D** (2002) The whole body receptive field of dorsal horn multireceptive neurones. *Brain Res Brain Res Rev* 40: 29–44.
125. **Sandkuhler J, Chen JG, Cheng G, Randic M** (1997) Low-frequency stimulation of afferent Adelta-fibers induces long-term depression at primary afferent synapses with substantia gelatinosa neurons in the rat. *J Neurosci* 17: 6483–6491.
126. **Finniss DG, Kaptchuk TJ, Miller F, Benedetti F** (2010) Biological, clinical, and ethical advances of placebo effects. *Lancet* 375: 686–695.
127. **Petersen GL, Finnerup NB, Norskov KN, Grosen K, Pilegaard HK, et al.** (2012) Placebo manipulations reduce hyperalgesia in neuropathic pain. *Pain* 153: 1292–1300.
128. **MacPherson H, Maschino AC, Lewith G, Foster NE, Witt CM, et al.** (2013) Characteristics of acupuncture treatment associated with outcome: an individual patient meta-analysis of 17,922 patients with chronic pain in randomised controlled trials. *PLoS One* 8: e77438.
129. **Sprenger C, Bingel U, Buchel C** (2011) Treating pain with pain: Spinal mechanisms of endogenous analgesia elicited by heterotopic noxious conditioning stimulation. *Pain* 152: 428–439.
130. **Talbot JD, Duncan GH, Bushnell MC, Boyer M** (1987) Diffuse noxious inhibitory controls (DNICs): psychophysical evidence in man for intersegmental suppression of noxious heat perception by cold pressor pain. *Pain* 30: 221–232.
131. **Wand-Tetley JI** (1956) Historical methods of counter-irritation. *Ann Phys Med* 3: 90–99.
132. **Carlsson C** (2002) Acupuncture mechanisms for clinically relevant long-term effects—reconsideration and a hypothesis. *Acupunct Med* 20: 82–99.

133. **Irnich D, Beyer A** (2002) [Neurobiological mechanisms of acupuncture analgesia]. *Schmerz* 16: 93–102.
134. **Tobbackx Y, Meeus M, Wauters L, De Vilder P, Roose J, et al.** (2013) Does acupuncture activate endogenous analgesia in chronic whiplash-associated disorders? A randomized crossover trial. *Eur J Pain* 17: 279–289.
135. **Garell PC, McGillis SL, Greenspan JD** (1996) Mechanical response properties of nociceptors innervating feline hairy skin. *J Neurophysiol* 75: 1177–1189.
136. **Tampin B, Slater H, Hall T, Lee G, Briffa NK** (2012) Quantitative sensory testing somatosensory profiles in patients with cervical radiculopathy are distinct from those in patients with nonspecific neck-arm pain. *Pain*.
137. **Maier C, Baron R, Tolle TR, Binder A, Birbaumer N, et al.** (2010) Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): somatosensory abnormalities in 1236 patients with different neuropathic pain syndromes. *Pain* 150: 439–450.
138. **Wylde V, Palmer S, Learmonth ID, Dieppe P** (2012) Somatosensory abnormalities in knee OA. *Rheumatology (Oxford)* 51: 535–543.
139. **Pfau DB, Rolke R, Nickel R, Treede RD, Daublaender M** (2009) Somatosensory profiles in subgroups of patients with myogenic temporomandibular disorders and Fibromyalgia Syndrome. *Pain* 147: 72–83.
140. **Jensen TS, Baron R** (2003) Translation of symptoms and signs into mechanisms in neuropathic pain. *Pain* 102: 1–8.
141. **Konopka KH, Harbers M, Houghton A, Kortekaas R, van Vliet A, et al.** (2012) Somatosensory profiles but not numbers of somatosensory abnormalities of neuropathic pain patients correspond with neuropathic pain grading. *PLoS One* 7: e43526.
142. **Cruccu G, Sommer C, Anand P, Attal N, Baron R, et al.** (2010) EFNS guidelines on neuropathic pain assessment: revised 2009. *Eur J Neurol* 17: 1010–1018.
143. **Pfau DB, Geber C, Birklein F, Treede RD** (2012) Quantitative sensory testing of neuropathic pain patients: potential mechanistic and therapeutic implications. *Curr Pain Headache Rep* 16: 199–206.
144. **Agostinho CM, Scherens A, Richter H, Schaub C, Rolke R, et al.** (2009) Habituation and short-term repeatability of thermal testing in healthy human subjects and patients with chronic non-neuropathic pain. *Eur J Pain* 13: 779–785.
145. **Westermann A, Ronnau AK, Krumova E, Regeniter S, Schwenkreis P, et al.** (2011) Pain-associated mild sensory deficits without hyperalgesia in chronic non-neuropathic pain. *Clin J Pain* 27: 782–789.
146. **Staud R, Vierck CJ, Cannon RL, Mauderli AP, Price DD** (2001) Abnormal sensitization and temporal summation of second pain (wind-up) in patients with fibromyalgia syndrome. *Pain* 91: 165–175.
147. **World Health Organisation** (1993) Standard Acupuncture Nomenclature - Second Edition. Available: http://www.centerfortraditionalmedicine.org/uploads/2/3/7/5/23750643/standard_acupuncture_nomenclature.pdf Accessed 2014 May 20.