



CASE REPORT

Open Access

A rapid and accurate new bedside test to assess maximal liver function: a case report

Sascha A Müller^{1,3*}, Ignazio Tarantino¹, Marcello Corazza¹, Frank Pianka¹, Jürgen Fornaro², Ulrich Beutner¹, Cornelia Lüthi¹ and Bruno M Schmied¹

Abstract

Background: In liver surgery, appropriate preoperative evaluation and preparation of the patient is of cardinal importance. The up-to-date, preoperative prediction of residual liver function has thus far been limited. As post-hepatectomy liver failure is a major cause of mortality, a new and simple bedside test (LiMAx) has been developed to predict postoperative liver function in conjunction with preoperative volumetric analysis of the liver.

Case presentation: A 45-year-old patient presented with a cecal carcinoma and a large synchronous liver metastasis for major liver surgery. Liver function was determined by the LiMAx-test for the enzymatic capacity of cytochrome P450 1A2, which is ubiquitously and solely active in the liver. A solution of 2 mg/kg body weight ¹³C-labeled methacetin was injected as a bolus into an intravenous catheter and, thereafter, was metabolized into acetaminophen and ¹³CO₂ and pulmonarily exhaled. The analysis of the ¹³CO₂/¹²CO₂ ratio was performed using online breath sampling over a period of maximally 60 minutes. Based on this test, a value of more than 315 µg/kg/h represents normal liver function. A laparoscopic right hemihepatectomy was planned during virtual resection with a residual liver volume of 48% and a preoperative anticipated residual LiMAx of 301 µg/kg/h. After successful resection, the initial postoperative LiMAx value was 316 µg/kg/h, indicating good liver function and a correct prediction of the outcome.

Conclusion: In the presented patient, residual liver function could be accurately predicted preoperatively using a combination of the new LiMAx test with CT-volumetry. This test might significantly improve preoperative evaluation and postoperative outcomes in liver surgery.

Keywords: Liver function, Liver volumetry, Liver resection, Liver function measurement, LiMAx

Background

Improvements in surgical techniques, anesthetic protocols, and perioperative care have decreased morbidity and mortality and have contributed to an increase in extended liver resections. This more aggressive approach inevitably leads to smaller remnant livers with a substantially elevated risk of post-resectional liver failure (PLF) [1-3]. It was demonstrated that 80% of deaths caused by PLF occur after resecting more than 50% of the liver volume [4-6], and the PLF incidence increases with the number of resected segments [5]. Therefore, accurate preoperative volumetric calculation of total and partial liver volumes based on preoperative imaging and

resection planning is crucial in the assessment of hepatic functional reserve and resectability, especially in cases of major liver resection and patients with underlying parenchymal disease [7]. Liver steatosis and steatohepatitis, for example, are associated with an increased risk of PLF after partial liver resection, especially after neoadjuvant chemotherapy, or in living donor liver transplantation [8]. Traditional methods estimating the hepatic functional reserve, such as liver enzyme tests and Child-Pugh classification, have clinical and practical limitations [9]. The major disadvantage of most quantitative liver function tests, such as the indocyanine green clearance test and the galactose elimination capacity, is that they are influenced by several factors such as hepatic blood flow and hyperbilirubinemia [10]. Moreover, no reliable cutoffs and guidelines could be defined. Consequently, clinical decisions mainly rely on the surgeon's experience [1].

* Correspondence: Sascha.mueller@kssg.ch

¹Department of Surgery, Kantonsspital St.Gallen, St.Gallen, Switzerland

³Rorschacherstrasse 95, 9007 St.Gallen, Switzerland

Full list of author information is available at the end of the article

Until now no biological marker or score has been able to reliably measure total liver function or predict the postoperative outcome. Here, we present the LiMAx-test, which has been developed at the Department of General, Visceral and Transplantation Surgery at the Charité in Berlin [11]. The test has been successfully evaluated in many patients undergoing liver surgery and liver transplantation since 2003 [11,12]. This test is now used in all patients with planned liver resection in our surgical department and in several other European centers.

The new LiMAx-test

This breath test is based on the hepatocyte-specific metabolism of the cytochrome P450 1A2 enzyme, which is active throughout the liver. After 3 hours of fasting, patients are placed in a resting horizontal position. The activity of the enzyme is measured by i.v. bolus injection of non-radioactive ^{13}C -methacetin (2 mg/kg body weight). Ten minutes prior to the injection with ^{13}C -methacetin, the baseline $^{13}\text{CO}_2/^{12}\text{CO}_2$ ratio is recorded. ^{13}C -methacetin (Euriso-top, Saint-Aubin, Cedex, France) is metabolized into acetaminophen and $^{13}\text{CO}_2$, which is pulmonarily exhaled. The exhaled air is collected using a face mask (Humedics GmbH, Berlin Germany), and the $^{13}\text{CO}_2/^{12}\text{CO}_2$ ratio in the air is determined using a modified nondispersive isotope-selective infrared spectroscopy (FANci2-db16, Fischer Analysen Instrumente, Leipzig, Germany, Humedics GmbH, Berlin, Germany) [11,13]. Based on the $^{13}\text{CO}_2/^{12}\text{CO}_2$ ratio and the body weight, the cytochrome P450 1A2 activity is determined using the so-called LiMAx value with the unit $\mu\text{g}/\text{kg}/\text{h}$ (μg methacetin/kg body weight/hour). The analysis of the $^{13}\text{CO}_2/^{12}\text{CO}_2$ ratio is performed over a period of 20 to 60 minutes. For normal liver function, a LiMAx value over 315 $\mu\text{g}/\text{kg}/\text{h}$ is required. In healthy volunteers, the normal range of LiMAx was found to be 425 ± 67 $\mu\text{g}/\text{kg}/\text{h}$ (range 311-575 $\mu\text{g}/\text{kg}/\text{h}$). Furthermore, the correlation between repeated measurements was found to be 0.82 [11].

Case report

A 45-year-old patient in good general health presented with a cecal carcinoma and a large synchronous liver metastasis. After neo-adjuvant chemotherapeutic treatment with 5-FU, Irinotecan and Oxaliplatin, a "liver first" approach was decided by the tumor board. The preoperative LiMAx value was 622 $\mu\text{g}/\text{kg}/\text{h}$, functional liver volume was 1736 mL, and tumor volume was 351 mL. A laparoscopic left hemihepatectomy was planned during virtual resection with a functional resection of 895 mL, resulting in a residual liver volume of 841 mL (48%) and a pre-operatively anticipated residual LiMAx of 301 $\mu\text{g}/\text{kg}/\text{h}$ (Figure 1). The laparoscopic left hemihepatectomy was performed under general anesthesia in the supine position using 5 ports according to the technique described by




Figure 1 Three-dimensional computation from a preoperative CT scan of the patient with a large liver metastasis (superior view). The portal vein is pink and green, the hepatic veins are blue, and the tumor is yellow.

Pearce et al. [14]. Briefly after applying a nylon tape around the portal triad for the Pringle maneuver, the liver was completely mobilized. The resection plane was defined by intraoperative ultrasound. After vascular control, the dissection was performed by harmonic scalpel (Harmonic ACE; Ethicon Endo-Surgery, Cincinnati, OH, USA) and Cavitron Ultrasonic Surgical Aspirator (Valleylab, Boulder, CO, USA) under intermittent Pringle maneuver. After completion of adequate hemostasis, the specimen was removed in an impermeable bag (Endocatch; Ethicon Endo-Surgery) introduced through a 15-mm suprapubic port subsequently extended through a Pfannenstiel incision. The operation time was 5.5 hours, and the intraoperative blood loss was 300 mL.

After successful resection, the initial postoperative LiMAx value on postoperative day one was 316 $\mu\text{g}/\text{kg}/\text{h}$, indicating good liver function and correct prediction of the outcome. After an uneventful postoperative course, the patient was discharged on postoperative day six. Eight days later, an uneventful laparoscopic right hemicolectomy was performed to remove the primary tumor.

Discussion

As impressively demonstrated by the presented case, the LiMAx-test in combination with a CT-based liver volumetry can reliably predict the postoperative liver function. Therefore, we adopted these preoperative tests as a standard procedure for every patient undergoing liver resection at our institution.

In 2010, Stockmann et al. published the LiMAx algorithm, which is an easily applicable clinical decision tree for preoperative resection planning [15]. This allows

effective preoperative surgical risk evaluation to prevent major complications or even liver failure-related death. Normal liver function (LiMAX >315 µg/kg/h) allows the resection of up to four liver segments without further consideration. By contrast, patients with strongly impaired liver function (LiMAX <140 µg/kg/h, indicating significant hepatic injury) are not eligible for surgery as they are prone to developing postoperative liver failure even after minor liver resection. Patients with limited hepatic impairment and a preoperative LiMAX-value between 140–315 µg/kg/h are the most challenging ones. In these patients, precise liver volumetry is mandatory for a planned major liver resection (>four liver segments). Resections with an expected residual LiMAX value less than 80 µg/kg/h should be avoided because they are associated with a high mortality of up to 40%. Planned resections with a predicted residual LiMAX value between 80–100 µg/kg/h are considered critical (10.5% postoperative mortality), and thus alternative therapies or additional preoperative procedures should be considered (i.e., portal vein embolization, neo-adjuvant chemotherapy to downstage tumors). Liver resections with expected residual LiMAX values of >100 µg/kg/h can safely be performed.

The LiMAX test can clearly help to avoid post-resectional liver failure. Thus, one could expect a relevant impact on health care costs. Recently, Lock et al. published a study examining 45 patients after major liver resection [16]. Eight patients developed postoperative liver failure, seven of these died in the postoperative course. Patients developing post-resectional liver failure generated higher costs than patients without post-resectional liver failure (82,199 € [95%CI: 42,812 – 21,586 €] vs. 25,980 € [95%CI: 9,559 – 42,401 €]; $p = 0.013$) [14]. The proportion of intensive-care unit cost relative to total costs increased from 30% to more than 50%, and the cost for blood tests and products from the blood bank increased more than eightfold from 819 € [95%CI: 0 – 1,837 €] to 6,953 € [95%CI: 4,512 – 9,395 €]; $p \leq 0.001$). In covariate analysis, overall costs related to post-resectional liver failure were 56,219 € [95%CI: 12,327 – 100,066 €]. Routinely applying the LiMAX test, we can achieve an optimized and individualized treatment for patients scheduled for liver resections. This ultimately leads to economic savings by significantly decreasing morbidity and mortality and the overall length of hospital stay.

Conclusion

In the presented patient, residual liver function could be accurately predicted preoperatively by the combination of the new LiMAX test with CT-volumetry. Considering the existing literature, the LiMAX test significantly

improves preoperative evaluation and postoperative outcomes in liver surgery.

Consent

Written informed consent was obtained from the patient for the publication of this case report.

Competing interests

The authors declare no conflicts of interest. In particular, the authors do not have any specific conflict of interest related to the manufacturer of the LiMAX test, including intellectual property, patents or patents pending, industrial relation to the manufacturer, or speaker's honoraria.

Authors' contributions

Data were prepared and reviewed by SAM, MC, CL and FP. JF reviewed radiological images. The manuscript was drafted by SAM, IT and UB and critically edited by BMS. All authors read and approved the final manuscript.

Author details

¹Department of Surgery, Kantonsspital St.Gallen, St.Gallen, Switzerland.

²Institute of Radiology, Kantonsspital St.Gallen, St.Gallen, Switzerland.

³Rorschacherstrasse 95, 9007 St.Gallen, Switzerland.

Received: 13 February 2013 Accepted: 17 April 2013

Published: 25 April 2013

References

1. Schneider PD: Preoperative assessment of liver function. *Surg Clin North Am* 2004, **84**(2):355–373.
2. Belghiti J, Hiramatsu K, Benoist S, Massault P, Sauvanet A, Farges O: Seven hundred forty-seven hepatectomies in the 1990s: an update to evaluate the actual risk of liver resection. *J Am Coll Surg* 2000, **191**(1):38–46.
3. Clavien P-A, Petrowsky H, DeOliveira ML, Graf R: Strategies for safer liver surgery and partial liver transplantation. *N Engl J Med* 2007, **356**(15):1545–1559.
4. Poon RT, Fan ST, Lo CM, Liu CL, Lam CM, Yuen WK, Yeung C, Wong J: Improving perioperative outcome expands the role of hepatectomy in management of benign and malignant hepatobiliary diseases: analysis of 1222 consecutive patients from a prospective database. *Ann Surg* 2004, **240**(4):698–708.
5. Schindl MJ, Redhead DN, Fearon KCH, Garden OJ, Wigmore SJ: The value of residual liver volume as a predictor of hepatic dysfunction and infection after major liver resection. *Gut* 2005, **54**(2):289–296.
6. Hammond JS, Guha IN, Beekingham IJ, Lobo DN: Prediction, prevention and management of postresection liver failure. *Br J Surg* 2011, **98**(9):1188–1200.
7. Pianka F, Baumhauer M, Stein D, Radeleff B, Schmied BM, Meinzer H-P, Müller SA: Liver tissue sparing resection using a novel planning tool. *Langenbecks Arch Surg* 2011, **396**(2):201–208.
8. Veteläinen R, van Vliet A, Gouma DJ, van Gulik TM: Steatosis as a risk factor in liver surgery. *Ann Surg* 2007, **245**(1):20–30.
9. Mullin EJ, Metcalfe MS, Maddern GJ: How much liver resection is too much? *Am J Surg* 2005, **190**(1):87–97.
10. Stehr A, Ploner F, Traeger K, Theisen M, Zuelke C, Radermacher P, Matejovic M: Plasma disappearance of indocyanine green: a marker for excretory liver function? *Intensive Care Med* 2005, **31**(12):1719–1722.
11. Stockmann M, Lock JF, Riecke B, Heyne K, Martus P, Fricke M, Lehmann S, Niehues SM, Schwabe M, Lemke AJ, Neuhaus P: Prediction of postoperative outcome after hepatectomy with a new bedside test for maximal liver function capacity. *Ann Surg* 2009, **250**(1):119–125.
12. Lock JF, Schwabauer E, Martus P, Videv N, Pratschke J, Malinowski M, Neuhaus P, Stockmann M: Early diagnosis of primary nonfunction and indication for reoperation after liver transplantation. *Liver Transpl* 2010, **16**(2):172–180.
13. Riecke B, Neuhaus P, Stockmann M: Major influence of oxygen supply on ¹³CO₂:¹²CO ratio measurement by nondispersive isotope-selective infrared spectroscopy. *Helicobacter* 2005, **10**(6):620–622.
14. Pearce NW, Di Fabio F, Teng MJ, Syed S, Primrose JN, Abu Hilal M: Laparoscopic right hepatectomy: a challenging, but feasible, safe and efficient procedure. *Am J Surg* 2011, **202**(5):e52–e58.

15. Stockmann M, Lock JF, Malinowski M, Niehues SM, Seehofer D, Neuhaus P: **The LiMAx test: a new liver function test for predicting postoperative outcome in liver surgery.** *HPB (Oxford)* 2010, **12**(2):139–146.
16. Lock JF, Reinhold T, Malinowski M, Pratschke J, Neuhaus P, Stockmann M: **The costs of postoperative liver failure and the economic impact of liver function capacity after extended liver resection—a single-center experience.** *Langenbecks Arch Surg* 2009, **394**(6):1047–1056.

doi:10.1186/1754-9493-7-11

Cite this article as: Müller *et al.*: A rapid and accurate new bedside test to assess maximal liver function: a case report. *Patient Safety in Surgery* 2013 7:11.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

