

RESEARCH

Open Access



# Interleukin-1 $\beta$ mediates alterations in mitochondrial fusion/fission proteins and memory impairment induced by amyloid- $\beta$ oligomers

Andre F. Batista<sup>1</sup>, Tayná Rody<sup>1</sup>, Leticia Forny-Germano<sup>1</sup>, Suzana Cerdeiro<sup>1</sup>, Maria Bellio<sup>2</sup>, Sergio T. Ferreira<sup>1,3</sup>, Douglas P. Munoz<sup>4</sup> and Fernanda G. De Felice<sup>1,4,5\*</sup>

## Abstract

**Background:** The lack of effective treatments for Alzheimer's disease (AD) reflects an incomplete understanding of disease mechanisms. Alterations in proteins involved in mitochondrial dynamics, an essential process for mitochondrial integrity and function, have been reported in AD brains. Impaired mitochondrial dynamics causes mitochondrial dysfunction and has been associated with cognitive impairment in AD. Here, we investigated a possible link between pro-inflammatory interleukin-1 (IL-1), mitochondrial dysfunction, and cognitive impairment in AD models.

**Methods:** We exposed primary hippocampal cell cultures to amyloid- $\beta$  oligomers (A $\beta$ Os) and carried out A $\beta$ O infusions into the lateral cerebral ventricle of cynomolgus macaques to assess the impact of A $\beta$ Os on proteins that regulate mitochondrial dynamics. Where indicated, primary cultures were pre-treated with mitochondrial division inhibitor 1 (mdivi-1), or with anakinra, a recombinant interleukin-1 receptor (IL-1R) antagonist used in the treatment of rheumatoid arthritis. Cognitive impairment was investigated in C57BL/6 mice that received an intracerebroventricular (i.c.v.) infusion of A $\beta$ Os in the presence or absence of mdivi-1. To assess the role of interleukin-1 beta (IL-1 $\beta$ ) in A $\beta$ O-induced alterations in mitochondrial proteins and memory impairment, interleukin receptor-1 knockout (*Il1r1*<sup>-/-</sup>) mice received an i.c.v. infusion of A $\beta$ Os.

**Results:** We report that anakinra prevented A $\beta$ O-induced alteration in mitochondrial dynamics proteins in primary hippocampal cultures. Altered levels of proteins involved in mitochondrial fusion and fission were observed in the brains of cynomolgus macaques that received i.c.v. infusions of A $\beta$ Os. The mitochondrial fission inhibitor, mdivi-1, alleviated synapse loss and cognitive impairment induced by A $\beta$ Os in mice. In addition, A $\beta$ Os failed to cause alterations in expression of mitochondrial dynamics proteins or memory impairment in *Il1r1*<sup>-/-</sup> mice.

(Continued on next page)

\* Correspondence: [felice@bioqmed.ufrj.br](mailto:felice@bioqmed.ufrj.br)

<sup>1</sup>Institute of Medical Biochemistry Leopoldo de Meis, Federal University of Rio de Janeiro, CCS, room H2-019, Rio de Janeiro, RJ 21941-590, Brazil

<sup>4</sup>Centre for Neuroscience Studies, Queen's University, Kingston, Ontario K7L3N6, Canada

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

**Conclusion:** These findings indicate that IL-1 $\beta$  mediates the impact of A $\beta$ Os on proteins involved in mitochondrial dynamics and that strategies aimed to prevent pathological alterations in those proteins may counteract synapse loss and cognitive impairment in AD.

**Keywords:** Alzheimer's disease, Mitochondrial dynamics, Mitochondrial dysfunction, IL-1 $\beta$ , Neuroinflammation, A $\beta$  oligomers

## Introduction

Alzheimer's disease (AD) is characterized by progressive memory loss and brain dysfunction that results from deterioration of synapses and brain cells [1, 2]. Accumulation of extracellular amyloid plaques, intraneuronal neurofibrillary tangles, and oxidative stress are among the most prominent alterations observed in AD brains [2]. Oxidative stress occurs when mitochondrial function is impaired [3–5]. Mitochondrial function requires proper mitochondrial dynamics, which comprises continuous and balanced processes of mitochondrial fission, fusion, and transport [6]. In mammalian cells, mitochondrial fusion is controlled by two outer-membrane-localized proteins, mitofusins 1 and 2 (MFN-1 and MFN-2), and one inner-membrane-localized protein, optic atrophy 1 (OPA-1) [6]. Mitochondrial fission, on the other hand, is mainly controlled by dynamin-related protein 1 (Drp1), which is predominantly localized in the cytosol and is recruited to the mitochondrial outer membrane to drive this process [6].

Several studies have indicated that mitochondria are important for synaptic development and plasticity [7–10], with mitochondria fission required for the establishment and maintenance of long-term potentiation [8]. Impaired mitochondrial dynamics has been implicated in neurodegenerative disorders including Alzheimer's, Parkinson's, and Huntington's diseases [10–17]. Reduced expression of OPA-1, MFN-1, and MFN-2 and increased expression of fission protein 1 (Fis-1) and Drp1 have been reported in post-mortem AD brains [12, 18]. Mitochondrial dysfunction has further been observed in mouse models of AD [10, 12, 13, 19]. An elegant recent study showed that Drp1 inhibition protects against mitochondrial dysfunction and cognitive impairment in a mouse model of AD [13]. Amyloid- $\beta$  oligomers (A $\beta$ Os), toxins that accumulate in the brains of AD patients and impair synaptic function and structure [2, 20], have been shown to induce mitochondrial fragmentation and dysfunction in neurons [12, 13, 21].

Brain inflammation is an important feature of AD [22–24]. Specifically, the interleukin-1 beta (IL-1 $\beta$ ) pathway has been implicated in the pathogenesis of AD [25–29]. Studies have shown elevated IL-1 $\beta$  in the brain, cerebrospinal fluid, and plasma of AD patients [30–32]. Interestingly, IL-1 $\beta$  and its receptor, interleukin-1

receptor (IL-1R), are predominantly expressed in the hippocampus, consistent with a major role of IL-1 $\beta$  signaling in modulating hippocampal memory function [33].

The mechanisms underlying altered neuronal mitochondrial dynamics in AD remain poorly understood. Because A $\beta$ Os have been shown to trigger an increase in brain IL-1 $\beta$  levels [25, 34], we hypothesized that abnormal activation of this pro-inflammatory pathway could lead to synapse loss and memory impairment via alterations in mitochondrial dynamics. We show that anakinra, an IL-1R antagonist used in the treatment of rheumatoid arthritis [35, 36], protected against alterations in proteins involved in mitochondrial fission and fusion induced by A $\beta$ Os in hippocampal cultures. Alterations in mitochondrial proteins were observed in the frontal cortex of cynomolgus macaques that received intracerebroventricular (i.c.v.) infusions of A $\beta$ Os. Additionally, we found that mdivi-1, an inhibitor of mitochondrial fission that further affects cellular physiology in ways that are independent of mitochondrial fusion [37–41], prevented the increases in expression of Drp1 and IL-1 $\beta$  and attenuated cognitive deficits caused by A $\beta$ Os in mice. Interestingly, A $\beta$ Os failed to alter hippocampal expression of mitochondrial proteins and to cause cognitive impairment in IL-1 receptor knockout mice (*Il1r1*<sup>-/-</sup>) mice, implicating an inflammatory mechanism in the impaired balance of mitochondrial fission/fusion proteins that is linked to synapse loss and cognitive impairment. Together, results provide a basis for targeting IL-1 $\beta$  signaling and the expression of mitochondrial fission and fusion proteins as a potential therapeutic approach for AD.

## Material and methods

### Reagents

A $\beta$ <sub>1–42</sub> was from American Peptide (Sunnyvale, CA, USA). Ham's F12 medium (HF12-02) was from Caisson Labs (Smithfield, UT, USA). 1,1,1,3,3,3-Hexafluoro-2-propanol (HFIP), dimethyl sulfoxide (DMSO), bovine serum albumin (BSA), and mdivi-1 were from Sigma (St Louis, MO, USA). Prolong Gold Antifade, normal goat serum, and 4',6-diamidino-2-phenylindole (DAPI) were from Invitrogen (Carlsbad, CA, USA). Commercial IL-1ra (anakinra, Kineret) was from Swedish Orphan

Biovitrum (Stockholm, Sweden). MitoSOX™ Red Mitochondrial Superoxide Indicator was from Thermo Fisher (Waltham, MA, USA).

### A $\beta$ oligomers (A $\beta$ Os)

A $\beta$ Os were prepared from A $\beta$ <sub>1–42</sub> in aliquots as a dried hexafluoro-2-propanol film and stored at – 80 °C. The peptide film was dissolved in undiluted, sterile Me2SO to make a 5 mM solution. The solution was diluted to 100  $\mu$ M with PBS medium without glutamine (BioSource International, Rockville, MD, USA) and aged overnight at 4 °C. The preparation was centrifuged at 14,000 $\times$ g for 10 min at 4 °C to remove insoluble aggregates (protofibrils and fibrils), and the supernatants containing soluble A $\beta$ Os were transferred to clean tubes and stored at 4 °C. A $\beta$ Os were routinely prepared and characterized by size-exclusion chromatography (HPLC) and immunocytochemistry to detect oligomer binding to neurons with NU4 and absence of larger A $\beta$  aggregates (Suppl. Fig. 1) [42]. Biophysical/biochemical characterization of A $\beta$ Os has also been described previously by our group [43–46]. For experiments with cynomolgus macaques, A $\beta$ Os were prepared in cold Ham's-F12 and routinely characterized by HPLC. A $\beta$ Os were always kept in 4 °C and used within 24 h.

The dosages of A $\beta$ Os used in vitro and in vivo experiments were based on previous studies from our group. We used a subtoxic concentration of A $\beta$ Os in cultures [47]. The in vivo dosage was chosen to be the lowest dose capable of causing persistent (3-week) memory impairment in mice [44].

### Treatments of cultures or mice

Hippocampal cultures were exposed to 500 nM A $\beta$ Os or an equivalent volume of vehicle (2% DMSO in phosphate-buffered saline (PBS) for 24 h). For IL-1 receptor blockade in hippocampal cultures, anakinra was added at a concentration of 100 ng/ $\mu$ L an hour before exposure to A $\beta$ Os. Mitochondrial Division inhibitor 1 (Mdivi-1) (25  $\mu$ M) was added to cultures 1 h before addition of A $\beta$ Os. Dosage of mdivi-1 was based on previous in vitro studies using this inhibitor [13, 48]. For in vitro experiments with MitoSOX Red, a mitochondrial superoxide indicator, 2.5  $\mu$ M was added to cultures for 10 min following the end of treatment with either A $\beta$ Os, mdivi-1, or anakinra. For inhibition of mitochondrial division in vivo, mice received mdivi-1 (40 mg/kg) or an equal volume of saline (i.p.) for 7 days. The choice of dose used in vivo was based on a previous literature report [13] and on the distribution and pharmacokinetics of mdivi-1 in mice [49].

### Mature hippocampal cultures and immunocytochemistry

Mixed hippocampal cultures were used after 18–21 days in vitro. Primary rat hippocampal cultures were prepared

and developed according to established procedure [50] and adapted from the original protocol described by Brewer [51]. Briefly, hippocampi from 18-day-old embryos were dissected and the meninges were removed. After that, tissue was dissociated in 0.05% PBS trypsin/EDTA solution at 37 °C for 5–10 min under occasional gentle shaking, followed by mechanical dissociation with glass Pasteur pipettes in DMEM medium supplemented with 5% horse-serum and 100 U/ml penicillin/streptomycin. Cells in suspension were counted in a Neubauer chamber and were plated on glass coverslips previously coated with 1 mg/ml poly-L-lysine, respectively. The density was 50,000 cells/well (260 cells/mm<sup>2</sup>) on coverslips for immunocytochemistry experiments. Cell cultures were maintained at 37 °C in a humidified atmosphere containing 5% CO<sub>2</sub> for 60 min, and DMEM was replaced by Neurobasal plus medium supplemented with 2% B27 supplement, 0.5 mM Glutamax. Cultures were maintained at 37 °C in a humidified atmosphere containing 5% CO<sub>2</sub> for 18–21 days prior to use. As shown in Suppl. Fig. 2a, we used mixed hippocampal neurons (containing ~ 40% neuronal cells and ~60% non-neuronal cells) in our study, to allow the cross talk between neuronal and non-neuronal cells.

All procedures were approved by the Institutional Animal Care and Use Committee of the Federal University of Rio de Janeiro (protocol #IBQM 022). After the pharmacological treatments described above, cultures were fixed with 4% paraformaldehyde for 15 min, permeabilized with 0.1% Triton X-100 for 5 min at room temperature, and blocked with 10% normal goat serum in PBS for 1 h before immunoreactions with anti-synaptophysin (mouse, 1:1,000; Abcam 14692), anti-postsynaptic density protein 5 (PSD-95) (rabbit, 1:500; Synaptic Systems 124 003), anti-MFN-1 (mouse, 1:200; Abcam Ab57602), or anti-MFN-2 (rabbit, 1:200; Sigma M6319). After overnight incubation with primary antibodies, cells were washed with PBS and incubated with Alexa-conjugated secondary antibodies for 2 h at room temperature. Nuclei were counterstained with ProLong containing DAPI antifade mounting solution. Coverslips were imaged on a Nikon C2 confocal microscope. MFN-1 and MFN-2 immunofluorescence intensities were analyzed in at least 3–4 experiments each (see figure legends) using independent neuronal cultures and A $\beta$ O preparations. In each experiment, 20–30 images (from 2 to 3 coverslips) were acquired from each experimental condition. Histogram analysis of fluorescence at each pixel across the images was performed using FIJI (National Institutes of Health), as described previously [52]. For quantification of synaptic punctae, a total of 15–20 images per experimental condition were acquired. PSD-95 (red) and synaptophysin (green) immunofluorescent puncta were analyzed and quantified using the Puncta

Analyzer plugin in NIH FIJI as previously described [43, 53, 54].

### Animals

Male C57BL/6 and *Il1r1*<sup>-/-</sup> (on a C57BL/6 background) mice, originally obtained from The Jackson Laboratories, were obtained from the animal facility at the Federal University of Rio de Janeiro and were 3 months old at the beginning of experiments. All procedures were approved by the Federal University of Rio de Janeiro Animal Care Committee (protocol number 022/18) and were in full compliance with the NIH Guide for Care and Use of Laboratory. All mice in this study were housed five per cage with ad libitum access to food and water in temperature- and humidity-controlled rooms on a 12-h light/dark cycle. The studies performed with macaques were approved by the Queen's University Animal Care Committee and were in full compliance with the Canada Council on Animal Care (Animal Care Protocol Original Munoz, 2011-039-Or). All macaques were maintained at the Centre for Neuroscience Studies at Queen's University (Kingston, Canada) and closely monitored by a lab animal technician and the Institute's veterinarian.

### Brain infusion of A $\beta$ Os in mice and treatment with mdivi-1

For i.c.v. infusion of A $\beta$ Os, mice were anesthetized for 7 min with 2.5% isoflurane (Cristália, São Paulo, Brazil) using a vaporizer system and gently restrained during the injection procedure. A final volume of 3  $\mu$ l (100 pmol A $\beta$ Os or vehicle) was injected into the lateral ventricle of the mice as described previously [44, 53, 55–57]. Briefly, a 2.5-mm needle was inserted unilaterally 1 mm to the right of the midline point equidistant from each eye and 1 mm posterior to a line drawn through the anterior base of the eye. Mice that showed signs of misplaced injections or any sign of hemorrhage (5% of cases, on average) were excluded from statistical analysis. For inhibition of mitochondrial fission, mice received daily i.p. injections of mdivi-1 (40 mg/Kg) or vehicle (PBS) for 7 days. Dosage of mdivi-1 was based on previous in vivo studies using this inhibitor [10, 13, 58, 59].

### RNA extraction and qPCR

Hippocampi were homogenized in 1 ml Trizol (Thermo Fisher Scientific), and RNA extraction was performed using the SV Total RNA Isolation System (Promega) according to the manufacturer's instructions. Purity and amount of RNA were determined by the 260/280 nm absorbance ratio. Only preparations with ratios between 2.2 and 1.8 and no signs of RNA degradation were used. One microgram RNA was used for cDNA synthesis using the High-Capacity cDNA Reverse Transcription

Kit (Thermo Fisher Scientific). Expression of genes of interest was analyzed by qPCR on an Applied Biosystems 7500 RT-PCR system using the Power SYBR kit (Applied Biosystems, Waltham, MA, USA). Actin was used as an endogenous control. Cycle threshold (Ct) values were used to calculate fold-changes in gene expression using the  $2^{-\Delta\Delta C_t}$  method [60]. In all cases, reactions were performed in 15  $\mu$ l. Primer sequences for genes studied were: MFN-1, Forward: 5'-TCTCCAAGCCCAACATCTTCA-3' and Reverse: 5'-ACTCCGGCTCCGAAGCA-3'; MFN-2, Forward: 5'-GGGGCCTACATCCAAGAGAG-3' and Reverse: 5'-GGAGAACTTTGTCCCAGAGC-3'; OPA-1, Forward: 5'-TGGGCTGCAGAGGATGG-3' and Reverse: 5'-CCTGATGTCACGGTGTGAT-3'; Drp1, Forward: 5'-AGACGCTTAATCTGACGTTTGA C-3' and Reverse: 5'-AGGTGGCCTTAACACTAT TGACA-3'; Actb, Forward: 5'-TGTGACGTTGACAT CCGTAAA-3' and Reverse: 5'-GTACTTGCGC TCAGGAGGAG-3'.

### Brain infusion of A $\beta$ Os in macaques

Seven female cynomolgus macaques (*Maccaca fascicularis*, body weights: 4.7–7.0 kg) were used. Three sham-operated macaques, 9 years of age ( $n = 1$ ) and 16 years of age ( $n = 2$ ), were used as controls and underwent the full surgical procedure but did not receive intracerebroventricular injections. After a recovery period of 2–4 weeks, four macaques, 9 years old ( $n = 2$ ) and 16 years old ( $n = 2$ ), had a guide cannula inserted into the lateral ventricle. Control animals had the cannula implanted into the lateral ventricle in the same manner as the experimental animals but did not receive oligomer injections. A total of 4 animals received 10–100  $\mu$ g i.c.v. of A $\beta$ Os (1 injection per day every 3 days for up to 24 days) while sedated using ketamine (Vetoquinol, Québec, ON, Canada; 3mg/kg), medetomidine (Sandoz, Québec, ON, Canada; 0.15 mg/kg), and glycopyrrolate (Sandoz; 0.013 mg/kg). Although the experimental design involved injection of a fixed amount of 100  $\mu$ g of A $\beta$ Os per injection in all macaques, this amount was somewhat variable among animals, due to procedural limitations at the moment of injection, due to procedural limitations at the moment of injection, including partial clogging of the cannulas and liquid reflux through the cannula following injection. The total amount of A $\beta$ Os effectively introduced into the lateral ventricle in each animal was 300  $\mu$ g (macaque 1), 560  $\mu$ g (macaque 2), 544  $\mu$ g (macaque 3), and 474  $\mu$ g (macaque 4), as described in detail in our previous study [61]. Approximately 1 week after completion of the experimental protocol, macaques were sedated with intramuscular ketamine (10 mg/kg) followed by intravenous sodium pentobarbital (25 mg/kg) and heparin. They were then perfused intracardially with PBS followed by 4% paraformaldehyde in PBS. Brains



were removed, and coronal 40- $\mu$ m-thick sections were obtained.

#### Immunohistochemistry in macaque brain sections

For immunofluorescence analysis of macaque brain sections, tissue autofluorescence was quenched by incubation with 0.06% potassium permanganate (Merck) for 10 min at room temperature. Sections were blocked in 5% BSA and 5% normal goat serum in 1% Triton X-100 for 3 h at room temperature. Primary antibodies anti-MFN-1 (mouse, 1:200; Abcam Ab57602), MFN-2 (rabbit, 1:200; Sigma, M6319), OPA-1 (rabbit, 1:500; Abcam Ab42364), and Drp1 (rabbit, 1:100; Novus Biological NB11055237) were diluted in blocking solution, and sections were incubated at 4 °C for 16 h, followed by incubation with Alexa594-conjugated secondary antibodies (donkey anti-goat, 1:2,000; Thermo Fisher Scientific R37119) for 2 h at room temperature. Nuclei were stained with DAPI (Invitrogen, Carlsbad, CA, USA) for 5 min. Immunostaining was carried out in parallel in brain sections from all animals, and constant parameters for image acquisition and analysis were used. Slides were mounted with Prolong Gold Antifade with DAPI (Invitrogen) and imaged on a Zeiss Axio Observer Z1 microscope equipped with an Apotome module to minimize out-of-focus light. Cells positive for MFN-1, MFN-2, and Drp1 were manually counted in 20 z-stack images acquired using the same parameters from each region of interest for each animal. For quantification of OPA-1, a total of 15–20 z-stack images from each region of interest for each animal were acquired with the same parameters. The number of puncta was analyzed and quantified using the Puncta Analyzer plugin in NIH FIJI.

#### Behavioral testing

##### *Novel object recognition (NOR)*

The novel object recognition (NOR) memory task was carried out in an open field arena measuring 0.3 × 0.3 × 0.45 m as described [44]. Test objects were made of glass or plastic and had different shapes, colors, sizes, and textures. During both training and test sessions, objects were taped to the floor of the arena so that the animals could not move them. None of the objects used in the current study evoked innate preference. Initially, animals were trained in a 5-min habituation session, in which they were allowed to freely explore the empty arena. Total distance and velocity during the 5-min open field session were recorded as measures of locomotor activity, and no differences were found among experimental groups. After that, animals were placed in the arena for the training session, in which animals were placed at the center of the arena in the presence of two identical objects. The amount of time spent exploring each object was recorded. After 1 h, animals were again placed in

the arena for the test session, when one of the familiar objects was replaced by a different (novel) object. Results are expressed as the percentage of time exploring each object during the training and test sessions. Animals with normal memory remember the familiar object as such and spend more time exploring the novel object in the test session.

##### *Contextual fear conditioning (CFC)*

Contextual fear conditioning (CFC) was evaluated as described [53, 55, 56]. The apparatus (25 × 25 × 25 cm) consisted of a box with aluminum walls with a methacrylate door and a grid floor composed of stainless-steel bars spaced 1 cm apart and connected to an electric shock generator (Panlab®, Harvard Apparatus, Cornellà, Spain). Each mouse was placed in this chamber and allowed to freely explore the conditioning box for 3 min before receiving 2 footshocks (0.35 mA for 2 s, with 30 s interval between shocks). Each mouse remained in the chamber for 30 s after the last shock before being returned to its home cage. After 24 h, animals were again placed in the conditioning chamber for 5 min and freezing behavior was quantified using the Freezing® software version 1.3.04 (Panlab).

##### *Statistical analysis*

Results are presented as means ± SEM. All statistical analyses were performed using GraphPad Prism (version 8.0). Student's *t* test or two-way ANOVA followed by Holm-Šidák post hoc test was used for group comparisons, as indicated in the figure legends. Data from the NOR task were analyzed using a one-sample *t* test, comparing time spent exploring the novel object to the fixed value of 50% (chance level) as previously described [44]. *p* values < 0.05 were considered statistically significant.

## Results

### **Anakinra, an IL-1 receptor antagonist, attenuates the impact of A $\beta$ O<sub>s</sub> on mitochondrial membrane potential and fusion/fission proteins**

IL-1 $\beta$  has been found to be elevated in human AD brains [30–32] and in the brains of A $\beta$ O-infused mice [34]. IL-1 $\beta$  has also been implicated in mitochondrial dysfunction and damage [25, 62–64]. Thus, we initially investigated whether A $\beta$ O could induce alterations in mitochondrial fusion/fission proteins and whether activation of pro-inflammatory IL-1 $\beta$  signaling could be involved in mitochondrial pathology in primary hippocampal mixed cultures. Characterization of A $\beta$ O<sub>s</sub> by HPLC shows the presence of low- and high-molecular oligomers (Suppl. Fig. 1). Immunocytochemistry experiments further reveal A $\beta$ O binding to neurons and the absence of larger aggregated species (Suppl. Fig. 2b).

We first used anakinra, a recombinant form of the human endogenous IL-1 receptor antagonist (IL-1ra), which has been used in the treatment of rheumatoid arthritis [36, 65]. Treatment with anakinra prevented the decrease in mitochondrial membrane potential induced by A $\beta$ Os in hippocampal neurons (Fig. 1a, d). Moreover, anakinra attenuated the reductions in MFN-1 (Fig. 1b, e) and MFN-2 (Fig. 1c, f) immunoreactivities in A $\beta$ O-treated hippocampal neurons. These results suggest that IL-1 receptor activation induces mitochondrial dysfunction in neurons exposed to A $\beta$ Os.

#### **A $\beta$ oligomers alter levels of mitochondrial fusion/fission proteins in the brains of cynomolgus macaques**

We tested whether A $\beta$ Os, implicated as proximal synaptotoxins in AD [2, 20], might alter mitochondrial fission/fusion proteins in the primate brain. To this end, we carried out i.c.v. infusions of A $\beta$ Os in cynomolgus macaques (*Macaca fascicularis*), a model that recapitulates several aspects of human AD pathology, including tau hyperphosphorylation, tangle formation, synapse loss, and brain inflammation [53, 56, 61, 66]. We investigated the levels of mitochondrial proteins in the frontal cortex of macaques that received i.c.v. A $\beta$ O infusions because our previous study indicated that this region is targeted by A $\beta$ Os [61]. In accordance, we observed A $\beta$ O labeling in the frontal cortex of macaques that received A $\beta$ O infusions, while no labeling was observed in control macaques (Suppl. Fig. 2c). A $\beta$ Os caused decreases in densities of MFN-1- and MFN-2-positive cells and in the number of OPA-1 immunoreactive puncta and an increase in the number of Drp1-positive cells in the frontal cortex of macaques (Fig. 2a, b). These results are consistent with alterations in these proteins reported in post-mortem AD brains [12, 18] and suggest that A $\beta$ Os may underlie the alterations of levels of mitochondrial fusion/fission proteins in the AD brain.

#### **Mdivi-1 prevents A $\beta$ O-induced synapse loss and rescues memory impairment in mice**

Mitochondrial fission is upregulated in AD, and inhibitors of fission have been proposed as a strategy to counteract deregulated mitochondrial dynamics in AD [67–69]. This prompted us to evaluate whether mdivi-1, an inhibitor of Drp1, would be effective against the deleterious effects of A $\beta$ Os. In hippocampal cultures, treatment with mdivi-1 attenuated the decreases in pre- and post-synaptic markers (synaptophysin, shown in green, and PSD-95, shown in red, respectively) and synapse loss induced by exposure to A $\beta$ Os (Fig. 3a–d). Results suggest that impaired mitochondrial dynamics contribute to synapse loss induced by A $\beta$ Os.

Next, we determined whether mdivi-1 could prevent memory deficits induced by A $\beta$ Os in mice. To this end,


mice received an i.c.v. infusion of A $\beta$ Os and were assessed in the novel object recognition (NOR) test. Mice were then separated into two groups that were treated with saline or mdivi-1 (40 mg/Kg, i.p.) for 7 days (Fig. 3e) before re-testing in the NOR test. Consistent with our previous reports [43, 44, 53, 55, 56], i.c.v. infusion of A $\beta$ Os caused an impairment in object recognition memory 24-h post-infusion (Fig. 3f). While memory impairment persisted in saline-treated animals, A $\beta$ O-infused mice that were treated with mdivi-1 showed normal performance in the NOR task (Fig. 3g). The same mice were further assessed in the contextual fear conditioning (CFC) task, a hippocampus- and amygdala-dependent test [70]. Twenty-four hours after the training session, A $\beta$ O-infused mice presented significantly less freezing behavior compared to vehicle-injected mice, indicating impaired contextual memory (Fig. 3h). In contrast, A $\beta$ O-infused mice that were treated with mdivi-1 performed similar to control animals (Fig. 3h).

We further examined the effects of mdivi-1 on Drp1 expression in the hippocampi of mice that received an i.c.v. infusion of A $\beta$ Os. While A $\beta$ Os induced an increase in hippocampal expression of Drp1, treatment with mdivi-1 corrected Drp1 expression in A $\beta$ O-infused mice (Fig. 3i). Results demonstrate that mdivi-1 rescued abnormal hippocampal expression of Drp1 and memory deficits induced by A $\beta$ Os in mice.

#### **IL-1R mediates A $\beta$ O-induced altered expression of mitochondrial fusion/fission proteins and memory impairment in mice**


While i.c.v. infusion of A $\beta$ Os induced NOR memory impairment in wild-type (WT) mice, A $\beta$ Os failed to impact memory in *Il1r1*<sup>-/-</sup> mice either 24 h or 7 days post-infusion (Fig. 4a–c). Similar results were observed when memory was assessed in the CFC task 9 days after i.c.v. infusion of A $\beta$ Os (Fig. 4d). *Il1r1*<sup>-/-</sup> mice were further protected against reductions in hippocampal expression of MFN-1, MFN-2, and OPA-1 (Fig. 4e–g) and increased expression of Drp1 (Fig. 4h) induced by A $\beta$ Os. Control measurements showed no significant changes in overall mitochondrial content, as indicated by expression of COX-IV, between vehicle- and A $\beta$ O-infused mice (Suppl. Fig. 3).

Finally, because previous studies have shown that excessive mitochondrial fission exacerbates the production of pro-inflammatory mediators by microglial cells [64, 71–74], we examined whether inhibition of Drp1 would alter IL-1 $\beta$  expression in mice. Whereas A $\beta$ Os induced increased hippocampal expression of IL-1 $\beta$  in control (saline-treated) mice, A $\beta$ Os failed to increase IL-1 $\beta$  expression in the hippocampi of mdivi-1-treated mice (Fig. 4i). Results thus indicate that treatment with mdivi-1 decreases hippocampal IL-1 $\beta$  expression in mice and link



**Fig. 1** IL-1 antagonist alleviates AβO-induced loss of mitochondrial membrane potential and fusion proteins in mature hippocampal cultures. **a** Representative images of mature hippocampal cultures exposed to AβOs (500 nM) or vehicle for 24 h. Where indicated, primary cultures were pre-treated for 40 min with anakinra (100 μM). The cultures were stained with a selective mitochondrial probe, Mitotracker (100 nM) (red). **b, c** Representative images of mature hippocampal cultures immunolabeled for MFN-1 (red) and MFN-2 (green). Scale bar: 60 μm. Optical zoom images of selected regions are also presented. Scale bar: 5 μm. **d** Integrated immunofluorescence of mitotracker (100 nM). Data expressed as means ± SEM from 4 experiments with independent cultures (30 images analyzed per experimental condition per experiment). \**p* < 0.05, \*\**p* < 0.01 compared with vehicle or AβOs; Two-way ANOVA followed by Holm-Šidak post hoc test. **e, f** Integrated immunoreactivities for MFN-1 and MFN-2. Data are expressed as means ± SEM from 4 to 5 experiments with independent cultures (30 images analyzed per experimental condition per experiment), \**p* < 0.05 compared with vehicle or AβOs. Two-way ANOVA followed by Holm-Šidak post hoc test





**Fig. 2** AβOs trigger abnormal mitochondrial dynamic proteins changes in the brains of cynomolgus macaques. **a** Representative images of MFN-1, MFN-2, OPA-1, and Drp1 immunolabeling in the frontal cortex of sham-operated and AβO-injected macaques. Nuclear staining (DAPI) is shown in blue. Scale bar: 20 μM. **b** Quantification of MFN-1-, MFN-2-, OPA-1-, and Drp1-positive cells. Data are expressed as means ± SEM ( $n = 3$  sham;  $n = 4$  AβOs). \*\* $p < 0.01$ , \* $p < 0.05$ , Student's unpaired  $t$  test

pro-inflammatory IL-1 $\beta$  signaling to impairments in mitochondrial dynamics.

## Discussion


In the current work, we show that AβOs, toxins that accumulate in AD brains [75, 76] and have been implicated in synapse damage and memory impairment in AD [2, 20], trigger alterations in proteins involved in mitochondrial dynamics in experimental models of AD, including cynomolgus macaques that received i.c.v. injections of AβOs. Changes in mitochondrial proteins and cognitive impairment induced by AβOs were found to be mediated by IL-1R activation.

Previous studies have reported changes in proteins involved in mitochondrial fission/fusion in vitro and in AD mouse models [6, 9, 10, 13, 19, 77]. In post-mortem AD brains, decreased expression of fusion proteins MFN-1, MFN-2, and OPA-1 and increased expression of


fission proteins Fis-1 and Drp1 have been reported [12, 19]. These findings suggest that an imbalance in levels of proteins involved in mitochondrial dynamics play a role in the pathogenesis of AD.

Aβ has been found to induce decreased expression of fusion proteins (MFN-1, MFN-2, and OPA-1) and increased expression of fission proteins (Drp1 and Fis-1) in AD models [3, 12, 18, 67, 78]. Recent studies further showed that excessive mitochondrial fission caused by increased Drp1 expression led to mitochondrial dysfunction in neurons exposed to Aβ [13, 67]. While our findings are in line with those studies, it is important to note that Aβ has been proposed to induce pathology by inhibiting Drp1 or fission more generally [77, 79, 80]. In addition, decreases in mitochondrial fission have been shown to cause fragmentation and mitochondrial abnormalities. Such studies suggest that any imbalance of





**Fig. 3** Mdivi-1 rescues AβO-induced cognitive impairment, defective mitochondrial fission protein expression, and inflammation in mice and prevents synapse loss in hippocampal neurons. **a** Representative images of cultured hippocampal neurons exposed to 500 nM AβOs or vehicle for 24 h and immunolabeled for synaptophysin (green) or PSD-95 (red). Synapses evidenced by colocalized puncta, appear in yellow. Where indicated, neurons were pre-incubated for 1 h with mdivi-1 (25 μM). Scale bar: 20 μm. **b, d** Integrated immunoreactivities, for synaptophysin (**b**), PSD-95 (**c**), and number of colocalized synaptophysin/PSD95 puncta (**d**). Data are expressed as means ± SEM from 4 experiments from independent cultures (30 images analyzed per experimental condition per independent culture). \**p* < 0.05, \*\**p* < 0.01 compared with vehicle or AβOs; two-way ANOVA followed by Holm-Šidak post hoc test. **e** Experimental design of the treatment of mdivi-1. Male C57BL/6 mice received an i.c.v. injection of AβOs (100 pmol) or vehicle. Twenty-four hours after i.c.v. injection of AβOs (100 pmol) or vehicle, animals underwent the NOR. The percentage of time exploring familiar or novel objects in the test session is shown in **f**. Mdivi-1 (40 mg/kg, i.p.) treatment began on day 3 and lasted for 7 days. On day 10, animals underwent NOR again. **g** after the mdivi-1 (40 mg/kg, i.p.) treatment. **h** 11 days after i.c.v. injection of AβOs (100 pmol), the same animals were trained in the contextual fear conditioning task. The percentage of freezing behavior in this task is shown. In **f** (*n* = 17–18 animals per group) and **g** (*n* = 9–10 animals per group), data are expressed as means ± SEM of the percentage of time spent on each object presented at the test session and \**p* < 0.05, compared with a hypothetical value of 50% of exploration time; one-sample Student's *t* test. In **h**, data are expressed as means ± SEM (*n* = 9–10 animals per group). \*\**p* < 0.01, \**p* < 0.05, compared with vehicle or AβOs; two-way ANOVA followed by Holm-Šidak post hoc test. **i** Drp1 levels in the hippocampi of mice 12 days after i.c.v. injection of AβOs (100 pmol). Following AβO injection, animals were treated for 7 days with saline or mdivi-1 (40 mg/kg; i.p.; *n* = 8–9 animals/group). \**p* < 0.05 compared with vehicle or AβOs, two-way ANOVA followed by Holm-Šidak post hoc test.



**Fig. 4** IL-1 signaling mediates cognitive impairment and abnormal expression of mitochondrial fission and fusion proteins caused by AβOs in mice. **a** Experimental design for behavior experiments with *Il1r1*<sup>-/-</sup> mice i.c.v. injected with vehicle or 100 pmol AβOs. **b, c** Exploration times of WT or *Il1r1*<sup>-/-</sup> mice 24 h or 7 days after a single i.c.v. injection with vehicle or 100 pmol AβOs in the NOR task. Data are expressed as means ± SEM ( $n = 11$ –12 animals per experimental group). Asterisks denote a statistically significant ( $p < 0.05$ ) difference from 50% (reference value). One-sample Student's *t* test. **d** 9 days after i.c.v. injection of AβOs (100 pmol), the same animals were trained in the contextual fear conditioning task. The percentage of freezing behavior in this task is shown. Data are expressed as means ± SEM ( $n = 6$ –7 animals per experimental group). \*\* $p < 0.01$ , \*\*\* $p < 0.001$  compared to vehicle or AβOs, two-way ANOVA followed by Holm-Šidak post hoc test. **e** MFN-1, **f** MFN-2, **g** OPA-1, and **h** Drp1 mRNA levels in the hippocampi of mice 12 days after i.c.v. injection of AβOs (100 pmol). \* $p < 0.05$  compared with vehicle or AβOs ( $n = 6$ –7 animals per experimental group). Two-way ANOVA followed by Holm-Šidak post hoc test. **i** IL-1β mRNA levels in the hippocampi of mice 12 days after i.c.v. injection of AβOs (100 pmol). Following AβO injection, animals were treated for 7 days with saline or mdivi-1 (40 mg/kg; i.p.;  $n = 9$ –10 animals/group). \* $p < 0.05$  compared with vehicle or AβOs, two-way ANOVA followed by Holm-Šidak post hoc test

fusion/fission could be detrimental and future studies aiming at investigating the role of Drp1 in health and disease are warranted. While Drp1 inhibition may be detrimental in physiological conditions, it is possible that pathological conditions that cause increases in

Drp1 levels interfere with mechanisms involved in memory formation.

Mitochondria play a vital role in maintaining cellular function through their capacity to produce energy [81]. Fusion and fission mechanisms participate in the

regulation of mitochondrial function, and disruption of these processes alters cellular localization and function of these organelles. Mitochondrial dynamics is physiologically relevant in regulation of mitochondrial function and metabolism, apoptosis, calcium homeostasis, autophagy, and mitophagy [5, 82]. Altered expression of proteins involved in mitochondrial dynamics has been associated with aging and age-related alterations [6, 10]. We found that alterations in mitochondrial fusion/fission proteins induced by A $\beta$ O are linked to synapse damage/loss and memory impairment. Mitochondrial dysfunction has been associated with synaptic failure [7, 9, 13, 67]. In neurons, mitochondrial dynamics has been implicated in the formation of synapses and dendritic spines [6, 7, 81]. GTPases MFN-1, MFN-2, and OPA-1 appear to be key for mitochondrial fusion, whereas Drp1 is needed for fission [6]. Mdivi-1 is expected to inhibit Drp1 and induce several outcomes that impact cell physiology. It is noteworthy the findings suggesting that mdivi-1 is not a specific Drp1 inhibitor, as mdivi-1 was found to modulate ROS generation by brain mitochondria, reversibly inhibit complex I in WT and Drp1 knockout (KO) fibroblasts [37]. In addition, in cancer cells, mdivi-1 was shown to decrease oxidative metabolism [41]. In the yeast pathogen *Candida albicans*, mdivi-1 was shown to impact endogenous nitric oxide levels [39]. Collectively, these studies indicate that such actions of mdivi-1 may contribute to its effects reported in our study and in other models of disease [38, 40]. Future studies will be important to better understand the role of mdivi-1 in physiological and pathological conditions.

Brain inflammation plays a key role in neuronal dysfunction in AD [23, 83]. Levels of IL-1 $\beta$  have been shown to be elevated in AD brains [30, 84]. Moreover, studies have suggested that neuroinflammation triggered by IL-1 $\beta$  impairs mitochondrial function [25, 62, 63]. Anakinra, a recombinant IL-1R antagonist [36, 65], prevented A $\beta$ O-induced mitochondrial dysfunction and alterations in MFN-1 and MFN-2 levels in mixed hippocampal cultures. A $\beta$ O-induced alterations in expression of mitochondrial fusion/fission proteins and cognitive impairment in mice were found to depend on IL-1R activation. A $\beta$  is thus likely impacting glial cells that are present in hippocampal cultures and such cells are expected to play a key role in the detrimental effects of A $\beta$  on mitochondrial proteins. Our findings implicate altered mitochondrial dynamics in IL-1 $\beta$ -mediated synaptic and memory defects and are consistent with studies showing that blockade of IL-1R restores cognition and alleviates pathogenesis in AD mouse models [26, 27]. While the role of neuroinflammation on neuronal mitochondrial dynamics in AD has not been completely explored, there is evidence showing that production of inflammatory mediators can impair mitochondrial function [63, 85].


A very recent study [63] showed that overexpression of MFN-2 prevented peripheral lipopolysaccharide (LPS)-induced neuroinflammation in mice, with significant reductions in brain IL-1 $\beta$  levels and mitochondrial fragmentation in neurons. Collectively, these data support our hypothesis that neuroinflammation mediated by IL-1 $\beta$  affects mitochondrial dynamics and memory impairment in mice.

Interestingly, decreases in MFN-1 and MFN-2 in the brain were reported in a mouse model of metabolic syndrome that presented microglial activation and increased IL-1 $\beta$  expression [86]. Fang and colleagues (2019) further reported that A $\beta$  inhibited mitophagy and that improvements in neuronal mitophagy enhanced the phagocytic efficiency of microglia and decreased pro-inflammatory cytokines, including IL-1 $\beta$  [87]. These effects were associated with reversion of memory impairment in mice.

Restoration of mitochondrial fission/fusion balance has been proposed as a therapeutic strategy in neurodegenerative disorders, including Alzheimer's, Huntington's, and Parkinson's diseases and in metabolic disorders (i.e., diabetes and obesity) [13, 67–69, 78]. Baek and colleagues (2017) demonstrated that inhibition of Drp1 by mdivi-1 alleviated mitochondrial fragmentation, loss of mitochondrial membrane potential, and ROS production and prevented memory impairment in the APP/PS1 AD mouse model [13]. Consistent with these findings, we observed that inhibition of mitochondrial fission alleviated mitochondrial dysfunction and, notably, reversed memory deficits in A $\beta$ O-infused mice.

Mitochondria are major players in the innate immune response [88, 89]. Studies have demonstrated the role of mitochondria in inflammation through activation of Nod-like receptor family, pyrin domain containing 3 (NLRP3) inflammasome [90, 91]. NLRP3 detects damaged mitochondria and initiates an inflammatory response leading to the release of IL-1 $\beta$  [91, 92]. Our findings extend those of previous studies in models of blood-brain barrier (BBB) disruption [72] and LPS-stimulated microglial cells [63, 64, 71], where treatment with mdivi-1 decreased levels of IL-1 $\beta$  and other pro-inflammatory mediators. Even though the acute effects of mdivi-1 on IL-1 $\beta$  have not been investigated in our study, our experimental design allowed the determination of IL-1 $\beta$  expression in mice that received an i.c.v. infusion of A $\beta$ O and were treated for 7 consecutive days with mdivi-1. Our findings that mdivi-1 blocks the increase in IL-1 $\beta$  expression induced by A $\beta$ O under these conditions suggests that blocking mitochondrial pathology may be relevant to block the sustained induction of IL-1 $\beta$  in mice. Fan and colleagues [72] reported that treatment with mdivi-1 decreased brain expression of IL-1 $\beta$ , TNF- $\alpha$ , and IL-6 24 h after subarachnoid





hemorrhage injury in rats. This suggests that, in addition to the longer-term effects revealed in our study, mdivi-1 may also have an acute impact on IL-1 $\beta$  in rodents. Our results further suggest that regulation of proper mitochondrial dynamics may comprise an approach to counteract excessive brain inflammation promoted by IL-1 $\beta$ . Interestingly, we found that A $\beta$ Os failed to cause cognitive impairment in *Il1r1*<sup>-/-</sup> mice and that anakinra attenuated alterations in the levels of mitochondrial proteins induced by A $\beta$ Os in vitro. Our results thus implicate IL-1 $\beta$ -dependent signaling in the mechanism of mitochondrial dysfunction in AD experimental models.

## Conclusion

Taken together, our findings indicate that activation of IL-1 receptor mediates the alteration in levels of mitochondrial fission/fusion proteins, leading to memory impairment induced by A $\beta$ Os (Fig. 5). Moreover, IL-1 $\beta$  expression in the hippocampus was found to be reduced following mdivi-1 treatment, suggesting a negative feedback loop between mitochondrial dynamics and IL-1 $\beta$ -mediated inflammation (Fig. 5). In conclusion, our results point to a bidirectional link between brain mitochondrial dynamics and IL-1 $\beta$  and suggest that strategies aimed to neutralize excessive IL-1 $\beta$  signaling

may be interesting therapeutic approaches to protect mitochondrial dynamics, synapses and memory in AD.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12974-021-02099-x>.

**Additional file 1.** Supplemental Figure 1. Characterization of A $\beta$ Os by HPLC. HPLC size-exclusion chromatography revealed that oligomer preparations comprised two major peaks. High-molecular-weight A $\beta$ Os are the major components in peak 1, while low-molecular-weight oligomers constitute the majority of peak 2

**Additional file 2.** Supplemental Figure 2. Characterization of A $\beta$ Os in hippocampal cultures and in the frontal cortex of macaques. (A) Percentage of MAP2-positive (neuronal) and MAP2-negative (non-neuronal) cells in five independent hippocampal cultures from E17-E18 rat embryos. Total cell count was determined by DAPI staining. (B) Representative images from hippocampal neurons in culture exposed to 500 nM A $\beta$ Os or vehicle for 24 h and immunolabeled for A $\beta$ Os (NU4 antibody). Scale bar: 20  $\mu$ m. (C) A $\beta$ O binding in the frontal cortex of a macaque that received A $\beta$ O injections (right panel). No A $\beta$ O immunostaining was detected in the frontal cortex of the sham-operated macaques (left panel). Scale bar: 25  $\mu$ m

**Additional file 3** Supplemental Figure 3. A $\beta$ Os did not cause a reduction of COX-IV expression in hippocampus of *Il1r1*<sup>-/-</sup> mice. COX-IV messenger ribonucleic acid (mRNA) levels in the hippocampi of *Il1r1*<sup>-/-</sup> mice 12 days after i.c.v. injection of A $\beta$ Os (100 pmol). Data are expressed as means  $\pm$  SEM ( $n = 6$ –7 animals per experimental group). Two-way ANOVA followed by Holm-Sidak post hoc test ( $p = 0.83$ )

## Abbreviations

Actb:  $\beta$ -Actin; AD: Alzheimer's disease; A $\beta$ O: Amyloid- $\beta$  oligomer; BBB: Blood-brain barrier; CFC: Contextual fear conditioning; Ct: Cycle threshold; DAPI: 4',6-Diamidino-2-phenylindole; Drp1: Dynamin-related protein 1; Fis-1: Fission protein 1; IL-1: Interleukin-1; IL-1 $\beta$ : Interleukin-1 beta; IL-1R: Interleukin-1 receptor; *Il1r1*<sup>-/-</sup>: IL-1 receptor knockout mice; i.c.v.: Intracerebroventricular; KO: Knockout; LPS: Lipopolysaccharide; Mdivi-1: Mitochondrial Division inhibitor 1; MFN-1: Mitofusin 1; MFN-2: Mitofusin 2; mRNA: Messenger ribonucleic acid; NLRP3: Nod-like receptor family, pyrin domain containing 3; NOR: Novel object recognition task; OPA-1: Optic atrophy 1; PSD-95: Postsynaptic density protein 5; PBS: Phosphate-buffered saline; ROS: Reactive oxygen species; RNA: Ribonucleic acid; RT-qPCR: Reverse transcription quantitative real-time polymerase chain reaction; WT: Wild-type

## Acknowledgements

We thank Maira Oliveira and Mariangela Viana for administrative and technical support. We thank Dr. William Klein (Northwestern University) for the kind gift of oligomer-specific NU4 antibody and Dr. Luis Eduardo Santos for performing A $\beta$ O preparation and characterization. We further acknowledge Dr. Helen Maciqueira de Melo, Dr. Mychael Lourenço, and Guilherme Braga for the characterization of mature hippocampal cultures.

## Authors' contributions

F.G.D.F., D.P.M., S.T.F., and A.F.B. did the conceptualization and design of the work. F.G.D.F., D.P.M., S.T.F., and A.F.B. did the methodology. A.F.B., T.R., L.F.G., and S.C. did the acquisition, analysis, and interpretation of data. M.B. contributed the resources. F.G.D.F., S.T.F., M.B., A.F.B., and T.R. did the writing. The authors read and approved the final manuscript.

## Funding

This work was supported by grants from the Human Frontiers Science Program (HFSP) and the John Simon Guggenheim Foundation (to F.G.D.F.), the National Institute for Translational Neuroscience (INNT/Brazil), and the Brazilian funding agencies Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Fundação de Amparo Pesquisa do Estado do Rio de Janeiro (FAPERJ) (to F.G.D.F. and S.T.F.), Canadian Institutes for Health Research (Grant MOP-38854 to D.P.M. and F.G.D.F.), and Canada Research Chair Program (D.P.M.). A.F.B. received a CNPq postdoctoral fellowship. T.R. is

supported by a CNPq predoctoral fellowship. L.F.G. is supported by a CNPq senior postdoctoral fellowship.

## Availability of data and materials

The datasets collected and analyzed during the current study are available from the corresponding author on reasonable request.

## Ethics approval and consent to participate

All protocols were approved by the Institutional Animal Care and Use Committee of the Federal University of Rio de Janeiro (protocol #IBQM 022). The studies performed with macaques were approved by the Queen's University Animal Care Committee and were in full compliance with the Canada Council on Animal Care (Animal Care Protocol Original Munoz, 2011-039-Or).

## Consent for publication

Not applicable.

## Competing interests

The authors declare no competing interests.

## Author details

<sup>1</sup>Institute of Medical Biochemistry Leopoldo de Meis, Federal University of Rio de Janeiro, CCS, room H2-019, Rio de Janeiro, RJ 21941-590, Brazil. <sup>2</sup>Department of Immunology, Institute of Microbiology Paulo de Góes, Federal University of Rio de Janeiro, Rio de Janeiro, RJ 21941-902, Brazil. <sup>3</sup>Institute of Biophysics Carlos Chagas Filho, Federal University of Rio de Janeiro, Rio de Janeiro, RJ 21941-902, Brazil. <sup>4</sup>Centre for Neuroscience Studies, Queen's University, Kingston, Ontario K7L3N6, Canada. <sup>5</sup>Department of Psychiatry, Queen's University, Kingston, Ontario K7L3N6, Canada.

Received: 2 July 2020 Accepted: 2 February 2021

Published online: 21 February 2021

## References

- Masters CL, Bateman R, Blennow K, Rowe CC, Sperling RA, Cummings JL. Alzheimer's disease. *Nat Rev Dis Primers*. 2015;1:15056.
- Selkoe DJ, Hardy J. The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol Med*. 2016;8(6):595–608.
- Barsoum MJ, Yuan H, Gerencser AA, Liot G, Kushnareva Y, Graber S, et al. Nitric oxide-induced mitochondrial fission is regulated by dynamin-related GTPases in neurons. *EMBO J*. 2006;25(16):3900–11.
- Martin-Maestro P, Gargini R, Garcia E, Perry G, Avila J, Garcia-Escudero V. Slower dynamics and aged mitochondria in sporadic Alzheimer's disease. *Oxid Med Cell Longev*. 2017;2017:9302761.
- Wai T, Langer T. Mitochondrial dynamics and metabolic regulation. *Trends Endocrinol Metab*. 2016;27(2):105–17.
- DuBoff B, Feany M, Gotz J. Why size matters - balancing mitochondrial dynamics in Alzheimer's disease. *Trends Neurosci*. 2013;36(6):325–35.
- Guo L, Tian J, Du H. Mitochondrial dysfunction and synaptic transmission failure in Alzheimer's disease. *J Alzheimers Dis*. 2017;57(4):1071–86.
- Rangaraju V, Lauterbach M, Schuman EM. Spatially stable mitochondrial compartments fuel local translation during plasticity. *Cell*. 2019;176(1–2):73–84 e15.
- Du H, Guo L, Yan S, Sosunov AA, McKhann GM, Yan SS. Early deficits in synaptic mitochondria in an Alzheimer's disease mouse model. *Proc Natl Acad Sci U S A*. 2010;107(43):18670–5.
- Manczak M, Reddy PH. Abnormal interaction between the mitochondrial fission protein Drp1 and hyperphosphorylated tau in Alzheimer's disease neurons: implications for mitochondrial dysfunction and neuronal damage. *Hum Mol Genet*. 2012;21(11):2538–47.
- Zhang L, Trushin S, Christensen TA, Bachmeier BV, Gateno B, Schroeder A, et al. Altered brain energetics induces mitochondrial fission arrest in Alzheimer's Disease. *Sci Rep*. 2016;6:18725.
- Wang X, Su B, Lee HG, Li X, Perry G, Smith MA, et al. Impaired balance of mitochondrial fission and fusion in Alzheimer's disease. *J Neurosci*. 2009;29(28):9090–103.
- Baek SH, Park SJ, Jeong JI, Kim SH, Han J, Kyung JW, et al. Inhibition of Drp1 ameliorates synaptic depression, A $\beta$  deposition, and cognitive impairment in an Alzheimer's disease model. *J Neurosci*. 2017;37(20):5099–110.

14. Exner N, Treske B, Paquet D, Holmstrom K, Schiesling C, Gispert S, et al. Loss-of-function of human PINK1 results in mitochondrial pathology and can be rescued by parkin. *J Neurosci*. 2007;27(45):12413–8.
15. Mortiboys H, Thomas KJ, Koopman WJ, Klaffke S, Abou-Sleiman P, Olpin S, et al. Mitochondrial function and morphology are impaired in parkin-mutant fibroblasts. *Ann Neurol*. 2008;64(5):555–65.
16. Wang H, Lim PJ, Karbowski M, Monteiro MJ. Effects of overexpression of huntingtin proteins on mitochondrial integrity. *Hum Mol Genet*. 2009;18(4):737–52.
17. Shirendeb U, Reddy AP, Manczak M, Calkins MJ, Mao P, Tagle DA, et al. Abnormal mitochondrial dynamics, mitochondrial loss and mutant huntingtin oligomers in Huntington's disease: implications for selective neuronal damage. *Hum Mol Genet*. 2011;20(7):1438–55.
18. Manczak M, Calkins MJ, Reddy PH. Impaired mitochondrial dynamics and abnormal interaction of amyloid beta with mitochondrial protein Drp1 in neurons from patients with Alzheimer's disease: implications for neuronal damage. *Hum Mol Genet*. 2011;20(13):2495–509.
19. Calkins MJ, Manczak M, Mao P, Shirendeb U, Reddy PH. Impaired mitochondrial biogenesis, defective axonal transport of mitochondria, abnormal mitochondrial dynamics and synaptic degeneration in a mouse model of Alzheimer's disease. *Hum Mol Genet*. 2011;20(23):4515–29.
20. Ferreira ST, Klein WL. The Abeta oligomer hypothesis for synapse failure and memory loss in Alzheimer's disease. *Neurobiol Learn Mem*. 2011;96(4):529–43.
21. Paula-Lima AC, Adasme T, SanMartin C, Sebollela A, Hetz C, Carrasco MA, et al. Amyloid beta-peptide oligomers stimulate RyR-mediated Ca<sup>2+</sup> release inducing mitochondrial fragmentation in hippocampal neurons and prevent RyR-mediated dendritic spine remodeling produced by BDNF. *Antioxid Redox Signal*. 2011;14(7):1209–23.
22. De Felice FG, Ferreira ST. Inflammation, defective insulin signaling, and mitochondrial dysfunction as common molecular denominators connecting type 2 diabetes to Alzheimer disease. *Diabetes*. 2014;63(7):2262–72.
23. Heneka MT, Carson MJ, El Khoury J, Landreth GE, Brosseron F, Feinstein DL, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol*. 2015;14(4):388–405.
24. Heneka MT, Kummer MP, Latz E. Innate immune activation in neurodegenerative disease. *Nat Rev Immunol*. 2014;14(7):463–77.
25. Parajuli B, Sonobe Y, Horiuchi H, Takeuchi H, Mizuno T, Suzumura A. Oligomeric amyloid beta induces IL-1beta processing via production of ROS: implication in Alzheimer's disease. *Cell Death Dis*. 2013;4:e975.
26. Kitazawa M, Cheng D, Tsukamoto MR, Koike MA, Wes PD, Vasilevko V, et al. Blocking IL-1 signaling rescues cognition, attenuates tau pathology, and restores neuronal beta-catenin pathway function in an Alzheimer's disease model. *J Immunol*. 2011;187(12):6539–49.
27. Martini AC, Gomez-Arboledas A, Forner S, Rodriguez-Ortiz CJ, McQuade A, Danhash E, et al. Amyloid-beta impairs TOM1-mediated IL-1R1 signaling. *Proc Natl Acad Sci U S A*. 2019;116(42):21198–206.
28. Ghosh S, Wu MD, Shaftel SS, Kyrkanides S, LaFerla FM, Olschowka JA, et al. Sustained interleukin-1beta overexpression exacerbates tau pathology despite reduced amyloid burden in an Alzheimer's mouse model. *J Neurosci*. 2013;33(11):5053–64.
29. Rivera-Escalera F, Pinney JJ, Owlett L, Ahmed H, Thakar J, Olschowka JA, et al. IL-1beta-driven amyloid plaque clearance is associated with an expansion of transcriptionally reprogrammed microglia. *J Neuroinflammation*. 2019;16(1):261.
30. Brosseron F, Trasschütz A, Widmann CN, Kummer MP, Tacik P, Santarelli F, et al. Characterization and clinical use of inflammatory cerebrospinal fluid protein markers in Alzheimer's disease. *Alzheimers Res Ther*. 2018;10(1):25.
31. Griffin WS, Stanley LC, Ling C, White L, MacLeod V, Perrot LJ, et al. Brain interleukin 1 and S-100 immunoreactivity are elevated in Down syndrome and Alzheimer disease. *Proc Natl Acad Sci U S A*. 1989;86(19):7611–5.
32. Cacabelos R, Barquero M, Garcia P, Alvarez XA, Varela de Seijas E. Cerebrospinal fluid interleukin-1 beta (IL-1 beta) in Alzheimer's disease and neurological disorders. *Methods Find Exp Clin Pharmacol*. 1991;13(7):455–8.
33. Schneider H, Pitossi F, Balschun D, Wagner A, del Rey A, Besedovsky HO. A neuromodulatory role of interleukin-1beta in the hippocampus. *Proc Natl Acad Sci U S A*. 1998;95(13):7778–83.
34. Ledo JH, Azevedo EP, Clarke JR, Ribeiro FC, Figueiredo CP, Foguel D, et al. Amyloid-beta oligomers link depressive-like behavior and cognitive deficits in mice. *Mol Psychiatry*. 2013;18(10):1053–4.
35. Mertens M, Singh JA. Anakinra for rheumatoid arthritis: a systematic review. *J Rheumatol*. 2009;36(6):1118–25.
36. Cvetkovic RS, Keating G. Anakinra. *BioDrugs*. 2002;16(4):303–11 discussion 13–4.
37. Bordt EA, Clerc P, Roelofs BA, Saladino AJ, Tretter L, Adam-Vizi V, et al. The putative Drp1 inhibitor mdivi-1 is a reversible mitochondrial complex I inhibitor that modulates reactive oxygen species. *Dev Cell*. 2017;40(6):583–94 e6.
38. Koch B, Traven A. Mdivi-1 and mitochondrial fission: recent insights from fungal pathogens. *Curr Genet*. 2019;65(4):837–45.
39. Koch B, Barugahare AA, Lo TL, Huang C, Schittenhelm RB, Powell DR, et al. A metabolic checkpoint for the yeast-to-hyphae developmental switch regulated by endogenous nitric oxide signaling. *Cell Rep*. 2018;25(8):2244–58 e7.
40. So EC, Hsing CH, Liang CH, Wu SN. The actions of mdivi-1, an inhibitor of mitochondrial fission, on rapidly activating delayed-rectifier K(+) current and membrane potential in HL-1 murine atrial cardiomyocytes. *Eur J Pharmacol*. 2012;683(1–3):1–9.
41. Dai W, Wang G, Chwa J, Oh ME, Abeywardana T, Yang Y, et al. Mitochondrial division inhibitor (mdivi-1) decreases oxidative metabolism in cancer. *Br J Cancer*. 2020;122(9):1288–97.
42. Lambert MP, Velasco PT, Chang L, Viola KL, Fernandez S, Lacor PN, et al. Monoclonal antibodies that target pathological assemblies of Abeta. *J Neurochem*. 2007;100(1):23–35.
43. Brito-Moreira J, Lourenco MV, Oliveira MM, Ribeiro FC, Ledo JH, Diniz LP, et al. Interaction of amyloid-beta (Abeta) oligomers with neuroligin 2alpha and neuroligin 1 mediates synapse damage and memory loss in mice. *J Biol Chem*. 2017;292(18):7327–37.
44. Figueiredo CP, Clarke JR, Ledo JH, Ribeiro FC, Costa CV, Melo HM, et al. Memantine rescues transient cognitive impairment caused by high-molecular-weight abeta oligomers but not the persistent impairment induced by low-molecular-weight oligomers. *J Neurosci*. 2013;33(23):9626–34.
45. Seixas da Silva GS, Melo HM, Lourenco MV, Lyra ESNM, de Carvalho MB, Alves-Leon SV, et al. Amyloid-beta oligomers transiently inhibit AMP-activated kinase and cause metabolic defects in hippocampal neurons. *J Biol Chem*. 2017;292(18):7395–406.
46. Sebollela A, Freitas-Correa L, Oliveira FF, Paula-Lima AC, Saraiva LM, Martins SM, et al. Amyloid-beta oligomers induce differential gene expression in adult human brain slices. *J Biol Chem*. 2012;287(10):7436–45.
47. De Felice FG, Wu D, Lambert MP, Fernandez SJ, Velasco PT, Lacor PN, et al. Alzheimer's disease-type neuronal tau hyperphosphorylation induced by A beta oligomers. *Neurobiol Aging*. 2008;29(9):1334–47.
48. Manczak M, Reddy PH. Mitochondrial division inhibitor 1 protects against mutant huntingtin-induced abnormal mitochondrial dynamics and neuronal damage in Huntington's disease. *Hum Mol Genet*. 2015;24(25):7308–25.
49. Rappold PM, Cui M, Grima JC, Fan RZ, de Mesy-Bentley KL, Chen L, et al. Drp1 inhibition attenuates neurotoxicity and dopamine release deficits in vivo. *Nat Commun*. 2014;5:5244.
50. De Felice FG, Vieira MN, Bomfim TR, Decker H, Velasco PT, Lambert MP, et al. Protection of synapses against Alzheimer's-linked toxins: insulin signaling prevents the pathogenic binding of Abeta oligomers. *Proc Natl Acad Sci U S A*. 2009;106(6):1971–6.
51. Brewer GJ. Isolation and culture of adult rat hippocampal neurons. *J Neurosci Methods*. 1997;71(2):143–55.
52. Abramoff MD, Magalhaes PJ, Ram SJ. Image processing with ImageJ. *Biophotonics International*. 2004;11(7):36–42.
53. Batista AF, Forny-Germano L, Clarke JR, Lyra ESNM, Brito-Moreira J, Boehnke SE, et al. The diabetes drug liraglutide reverses cognitive impairment in mice and attenuates insulin receptor and synaptic pathology in a non-human primate model of Alzheimer's disease. *J Pathol*. 2018;245(1):85–100.
54. Diniz LP, Tortelli V, Matias I, Morgado J, Bergamo Araujo AP, Melo HM, et al. Astrocyte transforming growth factor beta 1 protects synapses against Abeta oligomers in Alzheimer's disease model. *J Neurosci*. 2017;37(28):6797–809.
55. Lourenco MV, Frozza RL, de Freitas GB, Zhang H, Kincheski GC, Ribeiro FC, et al. Exercise-linked FNDG5/irisin rescues synaptic plasticity and memory deficits in Alzheimer's models. *Nat Med*. 2019;25(1):165–75.
56. Lourenco MV, Clarke JR, Frozza RL, Bomfim TR, Forny-Germano L, Batista AF, et al. TNF-alpha mediates PKR-dependent memory impairment and brain IRS-1 inhibition induced by Alzheimer's beta-amyloid oligomers in mice and monkeys. *Cell Metab*. 2013;18(6):831–43.
57. Clarke JR, Lyra ESNM, Figueiredo CP, Frozza RL, Ledo JH, Beckman D, et al. Alzheimer-associated Abeta oligomers impact the central nervous system to



- induce peripheral metabolic deregulation. *EMBO Mol Med.* 2015;7(2):190–210.
58. Bido S, Soria FN, Fan RZ, Bezard E, Tieu K. Mitochondrial division inhibitor-1 is neuroprotective in the A53T-alpha-synuclein rat model of Parkinson's disease. *Sci Rep.* 2017;7(1):7495.
  59. Li YH, Xu F, Thome R, Guo MF, Sun ML, Song GB, et al. Mdivi-1, a mitochondrial fission inhibitor, modulates T helper cells and suppresses the development of experimental autoimmune encephalomyelitis. *J Neuroinflammation.* 2019;16(1):149.
  60. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods.* 2001;25(4):402–8.
  61. Forny-Germano L, Lyra E, Silva NM, Batista AF, Brito-Moreira J, Gralle M, Boehnke SE, et al. Alzheimer's disease-like pathology induced by amyloid-beta oligomers in nonhuman primates. *J Neurosci.* 2014;34(41):13629–43.
  62. Wilkins HM, Swerdlow RH. Relationships between mitochondria and neuroinflammation: implications for Alzheimer's disease. *Curr Top Med Chem.* 2016;16(8):849–57.
  63. Harland M, Torres S, Liu J, Wang X. Neuronal mitochondria modulation of LPS-induced neuroinflammation. *J Neurosci.* 2020;40(8):1756–65.
  64. Nair S, Sobotka KS, Joshi P, Gressens P, Fleiss B, Thornton C, et al. Lipopolysaccharide-induced alteration of mitochondrial morphology induces a metabolic shift in microglia modulating the inflammatory response in vitro and in vivo. *Glia.* 2019;67(6):1047–61.
  65. Fiocco U, Vezzu M, Cozzi L, Todesco S. IL-1Ra (recombinant human IL-1 receptor antagonist) in the treatment of rheumatoid arthritis: the efficacy. *Reumatismo.* 2004;56(1 Suppl 1):62–73.
  66. Bomfim TR, Forny-Germano L, Sathler LB, Brito-Moreira J, Houzel JC, Decker H, et al. An anti-diabetes agent protects the mouse brain from defective insulin signaling caused by Alzheimer's disease-associated Aβ oligomers. *J Clin Invest.* 2012;122(4):1339–53.
  67. Reddy PH, Manczak M, Yin X. Mitochondria-division inhibitor 1 protects against Amyloid-beta induced mitochondrial fragmentation and synaptic damage in Alzheimer's disease. *J Alzheimers Dis.* 2017;58(1):147–62.
  68. Reddy AP, Reddy PH. Mitochondria-targeted molecules as potential drugs to treat patients with Alzheimer's disease. *Prog Mol Biol Transl Sci.* 2017;146:173–201.
  69. Reddy PH. Inhibitors of mitochondrial fission as a therapeutic strategy for diseases with oxidative stress and mitochondrial dysfunction. *J Alzheimers Dis.* 2014;40(2):245–56.
  70. Phillips RG, LeDoux JE. Differential contribution of amygdala and hippocampus to cued and contextual fear conditioning. *Behav Neurosci.* 1992;106(2):274–85.
  71. Park J, Choi H, Min JS, Park SJ, Kim JH, Park HJ, et al. Mitochondrial dynamics modulate the expression of pro-inflammatory mediators in microglial cells. *J Neurochem.* 2013;127(2):221–32.
  72. Fan LF, He PY, Peng YC, Du QH, Ma YJ, Jin JX, et al. Mdivi-1 ameliorates early brain injury after subarachnoid hemorrhage via the suppression of inflammation-related blood-brain barrier disruption and endoplasmic reticulum stress-based apoptosis. *Free Radic Biol Med.* 2017;112:336–49.
  73. Katoh M, Wu B, Nguyen HB, Thai TQ, Yamasaki R, Lu H, et al. Polymorphic regulation of mitochondrial fission and fusion modifies phenotypes of microglia in neuroinflammation. *Sci Rep.* 2017;7(1):4942.
  74. Ho DH, Je AR, Lee H, Son I, Kwon HS, Kim HG, et al. LRRK2 kinase activity induces mitochondrial fission in microglia via Drp1 and modulates neuroinflammation. *Exp Neurobiol.* 2018;27(3):171–80.
  75. Gong Y, Chang L, Viola KL, Lacor PN, Lambert MP, Finch CE, et al. Alzheimer's disease-affected brain: presence of oligomeric Aβ ligands (ADDLs) suggests a molecular basis for reversible memory loss. *Proc Natl Acad Sci U S A.* 2003;100(18):10417–22.
  76. Lesne SE, Sherman MA, Grant M, Kuskowski M, Schneider JA, Bennett DA, et al. Brain amyloid-beta oligomers in ageing and Alzheimer's disease. *Brain.* 2013;136(Pt 5):1383–98.
  77. Wang X, Su B, Zheng L, Perry G, Smith MA, Zhu X. The role of abnormal mitochondrial dynamics in the pathogenesis of Alzheimer's disease. *J Neurochem.* 2009;109(Suppl 1):153–9.
  78. Wang W, Yin J, Ma X, Zhao F, Siedlak SL, Wang Z, et al. Inhibition of mitochondrial fragmentation protects against Alzheimer's disease in rodent model. *Hum Mol Genet.* 2017;26(21):4118–31.
  79. Wang X, Su B, Fujioka H, Zhu X. Dynamin-like protein 1 reduction underlies mitochondrial morphology and distribution abnormalities in fibroblasts from sporadic Alzheimer's disease patients. *Am J Pathol.* 2008;173(2):470–82.
  80. Wang X, Su B, Siedlak SL, Moreira PI, Fujioka H, Wang Y, et al. Amyloid-beta overproduction causes abnormal mitochondrial dynamics via differential modulation of mitochondrial fission/fusion proteins. *Proc Natl Acad Sci U S A.* 2008;105(49):19318–23.
  81. Cai Q, Tammineni P. Mitochondrial aspects of synaptic dysfunction in Alzheimer's disease. *J Alzheimers Dis.* 2017;57(4):1087–103.
  82. Sebastian D, Palacin M, Zorzano A. Mitochondrial dynamics: coupling mitochondrial fitness with healthy aging. *Trends Mol Med.* 2017;23(3):201–15.
  83. Stephenson J, Nutma E, van der Valk P, Amor S. Inflammation in CNS neurodegenerative diseases. *Immunology.* 2018;154(2):204–19.
  84. Brosseron F, Krauthausen M, Kummer M, Heneka MT. Body fluid cytokine levels in mild cognitive impairment and Alzheimer's disease: a comparative overview. *Mol Neurobiol.* 2014;50(2):534–44.
  85. Zhou R, Yazdi AS, Menu P, Tschopp J. A role for mitochondria in NLRP3 inflammasome activation. *Nature.* 2011;469(7329):221–5.
  86. Tyagi A, Nguyen CU, Chong T, Michel CR, Fritz KS, Reisdorph N, et al. SIRT3 deficiency-induced mitochondrial dysfunction and inflammasome formation in the brain. *Sci Rep.* 2018;8(1):17547.
  87. Fang EF, Hou Y, Palikaras K, Adriaanse BA, Kerr JS, Yang B, et al. Mitophagy inhibits amyloid-beta and tau pathology and reverses cognitive deficits in models of Alzheimer's disease. *Nat Neurosci.* 2019.
  88. Sandhir R, Halder A, Sunkaria A. Mitochondria as a centrally positioned hub in the innate immune response. *Biochim Biophys Acta Mol Basis Dis.* 2017;1863(5):1090–7.
  89. West AP, Shadel GS, Ghosh S. Mitochondria in innate immune responses. *Nat Rev Immunol.* 2011;11(6):389–402.
  90. Iyer SS, He Q, Janczy JR, Elliott EJ, Zhong Z, Olivier AK, et al. Mitochondrial cardiolipin is required for Nlrp3 inflammasome activation. *Immunity.* 2013;39(2):311–23.
  91. Shimada K, Crother TR, Karlin J, Dagvadorj J, Chiba N, Chen S, et al. Oxidized mitochondrial DNA activates the NLRP3 inflammasome during apoptosis. *Immunity.* 2012;36(3):401–14.
  92. Nakahira K, Haspel JA, Rathinam VA, Lee SJ, Dolinay T, Lam HC, et al. Autophagy proteins regulate innate immune responses by inhibiting the release of mitochondrial DNA mediated by the NALP3 inflammasome. *Nat Immunol.* 2011;12(3):222–30.

# Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

