




Corrigendum

Corrigendum to “Factors Associated with the Prevalence of Thyroid Nodules and Goiter in Middle-Aged Euthyroid Subjects”

Dalia Dauksiene ¹, **Janina Petkeviciene** ², **Jurate Klumbiene**,² **Rasa Verkauskiene**,¹
Jelena Vainikonyte-Kristapone,³ **Audrone Seibokaite**,¹ **Jonas Ceponis**,¹
Vygantas Sidlauskas,¹ **Laura Daugintyte-Petrusiene**,³ **Antanas Norkus**,¹
and **Birute Zilaitiene** ¹

¹*Institute of Endocrinology, Medical Academy, Lithuanian University of Health Sciences, Kaunas, Lithuania*

²*Faculty of Public Health, Medical Academy, Lithuanian University of Health Sciences, Kaunas, Lithuania*

³*Department of Endocrinology, Medical Academy, Lithuanian University of Health Sciences, Kaunas, Lithuania*

Correspondence should be addressed to Dalia Dauksiene; dalia.dauksiene@gmail.com

Received 1 January 2019; Accepted 8 January 2019; Published 29 January 2019

Copyright © 2019 Dalia Dauksiene et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “Factors Associated with the Prevalence of Thyroid Nodules and Goiter in Middle-Aged Euthyroid Subjects” [1], an acknowledgment should be added as follows: “This research was funded by a grant no. SEN-14/2015 from the Research Council of Lithuania.”

References

- [1] D. Dauksiene, J. Petkeviciene, J. Klumbiene et al., “Factors associated with the prevalence of thyroid nodules and goiter in middle-aged euthyroid subjects,” *International Journal of Endocrinology*, vol. 2017, Article ID 8401518, 8 pages, 2017.