



Triglycerides and aortic pulse wave velocity in patients with chronic inflammation

Luca Zanoli*, Pasquale Fatuzzo, Pietro Castellino

Department of Clinical and Experimental Medicine, University of Catania, Catania, Italy

ARTICLE INFO

Article history:

Received 10 April 2018

Accepted 16 April 2018

Available online 26 April 2018

Keywords:

Arterial stiffness

Atherosclerosis

Inflammatory bowel disease

In a work published in a recent issue of the Egyptian Heart Journal,¹ Youssef et al. found that carotid stiffness was increased in subjects with rheumatoid arthritis, a disease characterized by chronic inflammation, and associated with disease duration. This finding is of clinical interest because support current concepts that chronic inflammation may lead to arterial stiffening.² Moreover, an association between arterial stiffness and triglycerides was also confirmed in subjects with rheumatoid arthritis, as previously reported.³ Similar results were also reported in a sample of the general population and in patients with metabolic syndrome.^{4,5} The relationship between triglycerides and arterial stiffness may involve local and systemic inflammation, oxidative stress and endothelial dysfunction. In this regard, triglycerides are associated with C-reactive protein levels in patients with chronic inflammation.⁶

We have recently reported that arterial stiffness and wave reflection are increased in patients with inflammatory bowel disease (IBD),^{7,8} a model of chronic inflammation characterized by low traditional cardiovascular risk factors and increased cardiovascular risk (the IBD paradox⁹), and associated with inflammation.¹⁰ Here, we reanalyze data of our recent individual participant data meta-analysis¹⁰ to test whether aortic pulse wave velocity (aPWV) was associated with triglycerides in patients with IBD. All analyses were performed in 111 participants with ulcerative colitis, 125




Fig. 1. Association between aortic pulse wave velocity (aPWV) and triglycerides in control subjects and in patients with ulcerative colitis (UC) or Crohn's disease (CD).

with Crohn's disease, and 147 control patients without missing data. The aPWV values were standardized for the patients with IBD and their respective control patients enrolled in each study included in these analyses by using the standard score (z-score).¹⁰ Because aPWV and triglycerides values were positively skewed, we used \log_e -transformed in the analyses. aPWV was associated with triglycerides at univariate analysis (Fig. 1). This association was confirmed ($b = 0.25$ z score; 95% confidence interval, 0.03–0.47 z

Peer review under responsibility of Egyptian Society of Cardiology.

* Corresponding author at: Nephrology, Department of Clinical and Experimental Medicine, Policlinico Universitario, University of Catania, Via Santa Sofia 78, 95123 Catania, Italy.

E-mail address: dott.zanoli@gmail.com (L. Zanoli).

<https://doi.org/10.1016/j.ehj.2018.04.005>

1110-2608/© 2018 Egyptian Society of Cardiology. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

score [$P = 0.02$]) in an outlier-robust multivariate linear regression model adjusted for age, sex, mean blood pressure, IBD, and study of origin. Longitudinal studies are needed to better clarify in patients with IBD the role of triglycerides on aortic stiffening.

Conflicts of interest

None.

References

1. Youssef G, Allam NT, Gaber W, Affi A, Hesham D. Increased arterial stiffness in rheumatoid arthritis and its relation to disease activity: a cross sectional study. *Egypt Heart J*. 2018;70:35–40. <https://doi.org/10.1016/j.ehj.2017.11.002>.
2. Zanolì L, Rastelli S, Inserra G, Castellino P. Arterial structure and function in inflammatory bowel disease. *World J Gastroenterol*. 2015;21:11304–11311. <https://doi.org/10.3748/wjg.v21.i40.11304>.
3. Sliem H, Nasr G. Change of the aortic elasticity in rheumatoid arthritis: relationship to associated cardiovascular risk factors. *J Cardiovasc Dis Res*. 2010;1:110–115. <https://doi.org/10.4103/0975-3583.70901>.
4. Zachariah JP, Rong J, Larson MG, et al. Metabolic predictors of change in vascular function: prospective associations from a community-based cohort. *Hypertension*. 2018;71:237–242. <https://doi.org/10.1161/HYPERTENSIONAHA.117.10054>.
5. Lopes-Vicente WRP, Rodrigues S, Cepeda FX, et al. Arterial stiffness and its association with clustering of metabolic syndrome risk factors. *Diabetol Metab Syndr*. 2017;9:87. <https://doi.org/10.1186/s13098-017-0286-1>.
6. Fan F, Galvin A, Fang L, et al. Comparison of inflammation, arterial stiffness and traditional cardiovascular risk factors between rheumatoid arthritis and inflammatory bowel disease. *J Inflamm (Lond)*. 2014;11:29. <https://doi.org/10.1186/s12950-014-0029-0>.
7. Zanolì L, Rastelli S, Granata A, et al. Arterial stiffness in inflammatory bowel disease: a systematic review and meta-analysis. *J Hypertens*. 2016;34:822–829. <https://doi.org/10.1097/HJH.0000000000000867>.
8. Zanolì L, Granata A, Lentini P, Gaudio A, Castellino P. Augmentation index is increased in patients with inflammatory bowel disease, a meta-analysis. *Eur J Intern Med*. 2017;39:e31–e32. <https://doi.org/10.1016/j.ejim.2016.12.012>.
9. Zanolì L, Inserra G, Castellino P. Increased cardiovascular risk in subjects with a low prevalence of classic cardiovascular risk factors: the inflammatory bowel disease paradox. *Trends Cardiovasc Med*. 2015;25:705–706. <https://doi.org/10.1016/j.tcm.2015.04.001>.
10. Zanolì L, Boutouyrie P, Fatuzzo P, et al. Inflammation and aortic stiffness: an individual participant data meta-analysis in patients with inflammatory bowel disease. *J Am Heart Assoc*. 2017;6. <https://doi.org/10.1161/JAHA.117.007003>.