

Research Article



Restoration of miR-193a-5p and miR-146 a-5p Expression Induces G1 Arrest in Colorectal Cancer through Targeting of MDM2/p53

Saeed Noorolyai¹, Elham Baghbani^{1,2}, Leili Aghebati Maleki¹, Amir Baghbanzadeh Kojabad¹, Dariush Shانهbansdi¹, Vahid Khaze Shahgoli¹, Ahad Mokhtarzadeh^{1*}, Behzad Baradaran^{1*}

¹Immunology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

²Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran.

Article info

Article History:

Received: 22 June 2019

Revised: 4 Aug. 2019

Accepted: 13 Aug. 2019

published: 11 Dec. 2019

Keywords:

- Colorectal cancer
- miRNA-193a-5p
- miRNA-146a-5p
- MDM2/p53
- Cell cycle
- Restoration

Abstract

Purpose: Colorectal cancer (CRC) remains a universal and lethal cancer owing to metastatic and relapsing disease. Currently, the role of microRNAs has been checked in tumorigenesis. Numerous studies have revealed that between the tumor suppressor miRNAs, the reduced expression of miR-146a-5p and -193a-5p in several cancers including CRC tissues are related with tumor progression and poor prognosis of patients. The purpose of this study is to examine the role of miR-146 a-5p and -193 a-5p in CRC cell cycle progression.

Methods: The miR-193a-5p and -146 a-5p mimics were transfected into HT-29 CRC cells via jetPEI transfection reagent and their impact was assessed on p53, cyclin B, and NF-κB gene expression. The inhibitory effect of these miRNAs on cell cycle was assessed by flow cytometry. The consequence of miR-193a-5p and miR-146 a-5p on the protein expression level of Murine double minute 2 (MDM2) was assessed by western blotting.

Results: miR193a-5p and -146a-5p regulated the expression of MDM2 protein and p53, cyclin B, and NF-κB gene expression in CRC cells. Treatment of HT-29 cells with miRNA-146a-5p and -193a-5p induced G1 cell cycle arrest.

Conclusion: The findings of our study suggest that miR146a-5p and -193a-5p may act as a potential tumor suppressor by their influence on cell cycle progression in CRC cells. Thus, miRNA-146a-5p and -193a-5p restoration may be recommended as a potential therapeutic goal in the treatment of CRC patients.

Introduction

Colorectal cancer (CRC) is the third most prevalent cancer and second largest reason of cancer death in Europe and North America¹. Its general frequency is 5% and the 5-year survival rate varies from 40 to 60%.² Nevertheless, the accurate genetic and epigenetic aberrations in colon cancer development have not been comprehended entirely and further studies need to be carried out in this regard. MicroRNAs (miRNAs, miR) are endogenously stated small noncoding RNAs that prevent gene expression by base-pairing with complementary sequences of the 3'-untranslated region (3'-UTR) of messenger RNAs.³ Owing to the extensive regulator of gene expression, miRNAs take part in critical roles in several biological progressions, as well as metabolism, cell growth, transformation, and apoptosis.⁴ Dysregulations of miRNAs might be consequently found with cancer progression and carcinogenesis. Investigations have discovered that multiple miRNAs are upregulated or

downregulated in CRC and might contribute to the pathogenesis of CRC via directing the expression of crucial signaling molecules.⁵ Growing proofs have established that miRNAs function as tumor suppressors or oncogenes, proposing their significant probability as novel therapeutic goals.⁶ Multiple studies have revealed that between the tumor suppressor miRNAs, miR-146 a-5p and -193 a-5p expression in multiple cancers such as CRC are linked with tumor progression, chemotherapy resistance, metastasis, and poor prognosis of patients.^{7,8} Considering essential molecular apparatuses in CRC development and progression gives new prophecies in emerging unique factors for CRC cures.⁹ However, insufficient clues exist establishing the dependency among miRNAs and target genes in CRC and to disband their roles in tumorigenesis.¹⁰

MDM2-p53 pathway is regularly affected in CRC and is strongly related with poor prognosis.¹¹ Murine double minute 2 (MDM2) was duplicated from 3T3-DM, the transformed mouse cell line. MDM2 is the negative

*Corresponding Authors: Behzad Baradaran and Ahad Mokhtarzadeh, Tel: +98 41 33371440, Fax: +98 41 33371311, Email: baradaranb@tbzmed.ac.ir, Email: mokhtarzadehah@tbzmed.ac.ir

© 2020 The Author (s). This is an Open Access article distributed under the terms of the Creative Commons Attribution (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.

modulator of p53 gene and performs through impeding p53 transcriptional activity and transposition from the nucleus to the cytoplasm. The tumor suppressor protein p53 performs through directing downstream apparatus in cell cycle that clues to cell growth/postponement via apoptosis, DNA repair, and several other apparatuses.¹² Moreover, p53 is a crucial controller of cell cycle checkpoints and apoptosis, directing the transcription of genes associated with over-proliferation, together with p21 and BAX.¹³ Adjustment of MDM2 gene expression is related with some transcription factors as well as NF- κ B; furthermore, the MDM2 gene has been stated to have NF- κ B binding sites.¹⁴

The aim of this study was to restore and increase the expression levels of miRNA-193a-5p and -146a-5p in the CRC cell line using microRNA replacement therapy. It also attempted to explore the effects of this miRNA replacement in multiple molecular and cellular aspects of CRC cells such as cell cycle controlling.

Materials and Methods

Cell culture

HT-29 cell line was purchased from Pasture Institute (Tehran, Iran). This cell line was cultured in RPMI-1640 medium with 100 μ g/mL streptomycin and 100 IU/mL penicillin (Gibco, Maryland) and completed with 10% fetal bovine serum (FBS; Gibco Laboratories, Grand Island, NY). Cells were retained in an incubator with 5% CO₂ and 95% moistened atmosphere at 37°C.

Transfection of miRNAs

For this aim 2×10^5 HT-29 cells were cultured in a 6-well-plate with RPMI-1640 medium. After 24 h, the medium was changed with FBS and antibiotics-free Opti-MEM (Gibco Life Technologies, Gaithersburg, MD). The microRNA-146a-5p mimic and microRNA-193a-5p mimic with 100 pmol concentration were transfected to HT-29 cells separately and simultaneously using jetPEI in vitro transfection reagent (Polyplus, Illkirch, France) and conforming to the manufacturer's guidelines.

The N/P ratio is a measure of the ionic balance of the complexes. It refers to the number of nitrogen residues of jetPEITM per miRNA phosphate. Approximately one in three nitrogen atom of PEI is a cation, so electroneutrality of jetPEITM/miRNA complexes is reached for N/P = 2-3.

In practice, the best transfection results are obtained for N/P = 5 - 10. jetPEITM is supplied as a 7.5 mM solution (expressed in nitrogen residues) and 1 μ g of miRNA contains 3 nmoles of anionic phosphate.

RNA isolation and quantitative real-time PCR (qRT-PCR)

Whole RNA was extracted from the cells using a RiboEX reagent (GeneAll; GeneAll Biotechnology, Seoul, Korea). The quality and the amount of extracted RNAs were determined with NanoDrop spectrophotometer (Thermo

Fisher Scientific Life Sciences, USA). The mRNAs were reverse transcribed to cDNA using cDNA Synthesis Kit from Biofact Company (Daejeon, Korea) and following the manufacturer's protocols.

The qRT-PCR test was conducted using the LightCycler 96 (Roche Diagnostics, Mannheim, Germany) and SYBR green-master mix (Biofact, Daejeon, Korea).

The primer sequences have been presented in Table 1. The assessment of p53, cyclin B, and NF- κ B was carried out by 10 minutes of initial denaturation at 94°C followed by 45 cycles of 94°C for 10 seconds and 60°C for 60 seconds. Finally, using 2^{- $\Delta\Delta$ Ct} cycle threshold method, the relative expression levels of genes were standardized with internal control U6 snRNA and GAPDH for miRNA and mRNA, respectively.

Table 1 illustrates the primer sequences.

Cell cycle analysis using DAPI and flow cytometry

Cells were harvested 48 hours after transfection and washed with cold PBS. Cell pellets were fixed with 70% ethanol at -20°C overnight. Subsequently, cells were washed with cold PBS and resuspended in 500 μ L PBS. Then, 5 μ L RNase A was added (10 mg/mL, Pishgam Biotech Co, Iran) and incubated for 30 minutes at 37°C. Next, it was resuspended in 500 μ L PBS, and 1 μ L DAPI (5 mg/mL, Sigma) and 1 μ L Triton-x100 (ACROS Organics, USA) were added and protected from light. Samples were analyzed through flow cytometry (MacsQuant Analyser 10, Miltenyi Biotech, Germany).

Western blotting

The total protein of whole cells was extracted using RIPA lysis buffer (Santa Cruz Biotechnology) and following the manufacturer's procedure. Fifty micrograms of extracted protein was added on 4% stacking and 12.5% running gel on SDS-PAGE electrophoresis; then, protein bands were blotted into the polyvinylidene difluoride (PVDF) (Roche Diagnostics GmbH, Germany) membrane through semidry immunoblotting. The membrane was blocked with 0.5% Tween and then incubated with MDM-2 and β -actin (as a housekeeping protein) monoclonal antibodies with 1: 1000 concentrations overnight at 4°C (Santa Cruz Biotechnology, USA). Next, the membrane was incubated with rabbit anti-goat secondary antibody

Table 1. The primers sequences

Name	Forward and reverse	Sequences
NF- κ B	F	5'-GCTACACAGGACCAGGGACAGT-3'
	R	5'-AGCTCAGCCTCATAGAAGCCATC-3'
Cyclin B	F	5'-TTGGTGTCACTGCCATGTTT-3'
	R	5'-GATGCTCTCCGAAGGAAGTG-3'
p53	F	5'-AAAGTCTAGAGCCACCGTCC-3'
	R	5'-AATCCAGGGAAGCGTGTCA-3'
GAPDH	F	5'-CAAGATCATCAGCAATGCCT-3'
	R	5'-GCCATCACGCCACAGTTTCC-3'

for MDM2 and rabbit anti-mouse antibody for β -actin, conjugated with horseradish peroxidase (1:2000; diluted in PBS) for 1 hour at room temperature. The protein bands were visualized using the electrochemiluminescence (ECL) kit (Roche Diagnostics, Germany) and western blot imaging system (Sabz Bimedicals Co., Iran). Finally, the amount of proteins was examined using the ImageJ software (National Institutes of Health, Bethesda, MD) and standardized with the β -actin.

Results and Discussion

The miR-193a-5p and -146a-5p replacement altered the expression of p53, cyclin B, and NF- κ B in HT-29 cell line

According to the results of qRT-PCR, while co-transfection and single-transfection with miRNA-146a-5p and -193a-5p significantly decreased the expression levels of cyclin B (Figure 1A) and NF- κ B (Figure 1B), they increased the expression level of p53 (Figure 1C).

The association between NF- κ B and p53 and its importance in the expansion and progression of cancer are fine recognized. NF- κ B is crucial for p53-mediated cell death.¹⁵ In our study, the expression level of NF- κ B was significantly reduced, and the expression of p53 was increased after miRNA-146a-5p and -193a-5p replacement. Along with these findings, we showed that transfection of miR-193a-5p and miR-146a-5p individually and simultaneously in HT-29 cell line increased the expression level of p53.

Co-treatment with miR-146a-5p and -193a-5p significantly suppressed MDM2 protein in HT-29 cells

To define the consequences of miRNA-146a-5p and

-193a-5p restoration on the activity of MDM2, we considered mimic replacement in colorectal cell line. The replacement of these miRNAs alone and in combination with each other affected the activity of MDM2. The miRNA-146a-5p and miRNA-193a-5p individually led to the reduction of MDM2 protein compared to control group. This evidence is particularly more significant in a group of cells transfected simultaneously with both of these miRNAs (The data represent mean \pm SD. n = 3, **** P < 0.0001, Figure 2).

In CRC, the overexpression of MDM2 is an early occurrence in the tumor progression. The MDM2 protein controls the action of p53 through developing a protein compound with p53, which could result in the beginning of p53 tumor suppressor role.¹⁶ Several studies and evidences have showed that MDM2 might be controlled by multiple cellular pathways, for example those related with miRNAs.^{17,18} Recently, abnormal miRNA expression profiles in multiple cancers including CRC have been reported by numerous studies.¹⁹ According to several studies, the expression levels of miR-193a-5p and miR-146a-5p are substantially reduced in CRC^{8,20}; these findings were confirmed by the results of our study. In the present study, we observed that replacement of miR-193a-5p and miR-146a-5p individually and simultaneously in HT-29 CRC cell line reduced the ratio of MDM2 protein.

The miR-193a-5p and -146a-5p replacement induced G1 cell cycle arrest

We examined cell cycle progression in different groups of transfected HT-29 cells. As shown in Figure 3, in the groups transfected with miRNA-193a-5p and -146a-5p mimic, as




Figure 1. The miR-193a-5p and -146a-5p restoration altered the expression of p53, cyclin B, and NF- κ B in HT-29 cell line. The outcomes of transfection of miRNA-146a-5p and -193a-5p alone and in combination with each other on the expression of (A)cyclin, (B) NF- κ B, and (C) p53 in comparison with the control group (** P < 0.01, *** P < 0.001, **** P < 0.0001).




Figure 2. The miR193a-5p and -146a-5p regulated the expression of MDM2 in CRC cell line. The immunoblotting of MDM2 protein was evaluated in untransfected (control) and miR193a-5p and miR146a-5p separately and simultaneously transfected groups (The data represent mean \pm SD, **** $P < 0.0001$).




Figure 3. The effect of miR-193a-5p and -146a-5p restoration on cell cycle progression of CRC cells. The miRNA-193a-5p and -146a-5p alone and in combination with each other caused G1 arrest in HT-29 cells.

well as transfected cells with both miRNAs simultaneously, G1 arrest occurred in cell cycle progression.

According to these findings, miR-193a-5p and miR-146a-5p inhibit cell growth and induce p53 dependent cell death. Also, we demonstrated that replacement of these miRNAs induced G1 phase arrest in HT-29 cells. Similar to our research, Liu et al showed that transfection of miRNA-193 into DU-145 cells of prostate cancer caused cell cycle arrest in the G1 phase.²¹ Also Li et al established that replacement of miRNA-146 in lung cancer cell lines caused cell cycle arrest in the phases G1 and G0.²² The p53 is a tumor suppressor gene that is controlled by miRNAs. Several oncogenic miRNAs straightly target p53, while specified tumor-suppressive miRNAs target negative regulators of p53, including the MDM2 and E3 ligase.^{23,24} Cross talk among p53 and miRNAs is multifaceted and numerous miRNAs control the expression and function of p53 such as miR146a-5p and miR193a-5p.^{25,26} Li et al reported that miR-193a directly targets MDM2, and also miR-193a persuades G1 arrest and apoptosis and returns leukemic cell differentiation; though, it is currently

mysterious as this happens through stimulation of p53.^{26,27}

Conclusion

In summary, our results displayed that miR146a-5p and miR193a-5p may affect MDM2 and p53 regulatory complex that controls CRC cells growth and expansion. Also, the results of current study suggest that miR146a-5p and miR193a-5p might function as a probable tumor suppressor displayed by their inhibition of cell cycle progress and elevation of apoptosis in CRC. Beside the assumption that miR146a-5p and miR193a-5p may have several target genes, these different target genes might apply numerous functions of these miRNAs in these biological procedures, enhancing vital queries for upcoming investigations.

Ethical Issues

Not applicable.

Conflict of Interest

The authors declare that there are no conflicts of interest.

Acknowledgments

This study was supported by Immunology Research Center of Tabriz University of Medical Sciences, Tabriz, Iran.

References

- Roseweir AK, Kong CY, Park JH, Bennett L, Powell A, Quinn J, et al. A novel tumor-based epithelial-to-mesenchymal transition score that associates with prognosis and metastasis in patients with Stage II/III colorectal cancer. *Int J Cancer* 2019;144(1):150-9. doi: 10.1002/ijc.31739
- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin* 2016;66(1):7-30. doi: 10.3322/caac.21332
- Hosseinali N, Aghapour M, Duijf PHG, Baradaran B. Treating cancer with microRNA replacement therapy: a literature review. *J Cell Physiol* 2018;233(8):5574-88. doi: 10.1002/jcp.26514
- Chen M, Xu R, Rai A, Suwakulsiri W, Izumikawa K, Ishikawa H, et al. Distinct shed microvesicle and exosome microRNA signatures reveal diagnostic markers for colorectal cancer. *PLoS One* 2019;14(1):e0210003. doi: 10.1371/journal.pone.0210003
- Bushati N, Cohen SM. microRNA functions. *Annu Rev Cell Dev Biol* 2007;23:175-205. doi: 10.1146/annurev.cellbio.23.090506.123406
- Slaby O, Svoboda M, Michalek J, Vyzula R. MicroRNAs in colorectal cancer: translation of molecular biology into clinical application. *Mol Cancer* 2009;8:102. doi: 10.1186/1476-4598-8-102
- Sathyanarayanan A, Chandrasekaran KS, Karunakaran D. microRNA-146a inhibits proliferation, migration and invasion of human cervical and colorectal cancer cells. *Biochem Biophys Res Commun* 2016;480(4):528-33. doi: 10.1016/j.bbrc.2016.10.054
- Shirafkan N, Shomali N, Kazemi T, Shanebandi D, Ghasabi M, Baghbani E, et al. microRNA-193a-5p inhibits migration of human HT-29 colon cancer cells via suppression of metastasis pathway. *J Cell Biochem* 2018. doi: 10.1002/jcb.28164
- Fritzmam J, Morkel M, Besser D, Budczies J, Kosel F, Brembeck FH, et al. A colorectal cancer expression profile that includes transforming growth factor beta inhibitor BAMBI predicts metastatic potential. *Gastroenterology* 2009;137(1):165-75. doi: 10.1053/j.gastro.2009.03.041
- Garofalo M, Croce CM. microRNAs: Master regulators as potential therapeutics in cancer. *Annu Rev Pharmacol Toxicol* 2011;51:25-43. doi: 10.1146/annurev-pharmtox-010510-100517
- Zhao Y, Yao YH, Li L, An WF, Chen HZ, Sun LP, et al. Pokemon enhances proliferation, cell cycle progression and anti-apoptosis activity of colorectal cancer independently of p14ARF-MDM2-p53 pathway. *Med Oncol* 2014;31(12):288. doi: 10.1007/s12032-014-0288-x
- Gupta A, Shah K, Oza MJ, Behl T. Reactivation of p53 gene by MDM2 inhibitors: a novel therapy for cancer treatment. *Biomed Pharmacother* 2019;109:484-92. doi: 10.1016/j.biopha.2018.10.155
- Subash-Babu P, Alshammari GM, Ignacimuthu S, Alshatwi AA. Epoxy clerodane diterpene inhibits MCF-7 human breast cancer cell growth by regulating the expression of the functional apoptotic genes Cdkn2A, Rb1, mdm2 and p53. *Biomed Pharmacother* 2017;87:388-96. doi: 10.1016/j.biopha.2016.12.091
- Busuttill V, Droin N, McCormick L, Bernassola F, Candi E, Melino G, et al. NF-kappaB inhibits T-cell activation-induced, p73-dependent cell death by induction of MDM2. *Proc Natl Acad Sci U S A* 2010;107(42):18061-6. doi: 10.1073/pnas.1006163107
- Patel M, Horgan PG, McMillan DC, Edwards J. NF-kappaB pathways in the development and progression of colorectal cancer. *Transl Res* 2018;197:43-56. doi: 10.1016/j.trsl.2018.02.002
- Mori D, Nakafusa Y, Miyazaki K, Tokunaga O. Differential expression of Janus kinase 3 (JAK3), matrix metalloproteinase 13 (MMP13), heat shock protein 60 (HSP60), and mouse double minute 2 (MDM2) in human colorectal cancer progression using human cancer cDNA microarrays. *Pathol Res Pract* 2005;201(12):777-89. doi: 10.1016/j.prp.2005.06.005
- Asslaber D, Piñón JD, Seyfried I, Desch P, Stöcher M, Tinhofer I, et al. microRNA-34a expression correlates with MDM2 SNP309 polymorphism and treatment-free survival in chronic lymphocytic leukemia. *Blood* 2010;115(21):4191-7. doi: 10.1182/blood-2009-07-234823
- Zhang J, Sun Q, Zhang Z, Ge S, Han ZG, Chen WT. Loss of microRNA-143/145 disturbs cellular growth and apoptosis of human epithelial cancers by impairing the MDM2-p53 feedback loop. *Oncogene* 2013;32(1):61-9. doi: 10.1038/onc.2012.28
- Jansson MD, Lund AH. MicroRNA and cancer. *Mol Oncol* 2012;6(6):590-610. doi: 10.1016/j.molonc.2012.09.006
- Iguchi T, Nambara S, Masuda T, Komatsu H, Ueda M, Kidogami S, et al. miR-146a polymorphism (rs2910164) predicts colorectal cancer patients' susceptibility to liver metastasis. *PLoS One* 2016;11(11):e0165912. doi: 10.1371/journal.pone.0165912
- Liu Y, Xu X, Xu X, Li S, Liang Z, Hu Z, et al. MicroRNA-193a-3p inhibits cell proliferation in prostate cancer by targeting cyclin D1. *Oncol Lett* 2017;14(5):5121-8. doi: 10.3892/ol.2017.6865
- Li YL, Wang J, Zhang CY, Shen YQ, Wang HM, Ding L, et al. MiR-146a-5p inhibits cell proliferation and cell cycle progression in NSCLC cell lines by targeting CCND1 and CCND2. *Oncotarget* 2016;7(37):59287-98. doi: 10.18632/oncotarget.11040
- Jansson MD, Damas ND, Lees M, Jacobsen A, Lund AH. miR-339-5p regulates the p53 tumor-suppressor pathway by targeting MDM2. *Oncogene* 2015;34(15):1908-18. doi: 10.1038/onc.2014.130
- Kim YJ, Lee JH, Jin S, Kim JH, Kim SH. Primate-specific miR-944 activates p53-dependent tumor suppression in human colorectal cancers. *Cancer Lett* 2019;440-441:168-79. doi: 10.1016/j.canlet.2018.10.029
- Gougelet A, Pissaloux D, Besse A, Perez J, Duc A, Dutour A, et al. Micro-RNA profiles in osteosarcoma as a predictive tool for ifosfamide response. *Int J Cancer* 2011;129(3):680-90. doi: 10.1002/ijc.25715
- Li Y, Gao L, Luo X, Wang L, Gao X, Wang W, et al. Epigenetic silencing of microRNA-193a contributes to leukemogenesis in t (8;21) acute myeloid leukemia by activating the PTEN/PI3K signal pathway. *Blood* 2013;121(3):499-509. doi: 10.1182/blood-2012-07-444729
- Tarasov VA, Matishov DG, Shin EF, Boiko NV, Timoshkina NN, Makhotkin MA, et al. Inheritable changes in miRNAs expression in HeLa cells after X-ray and mitomycin C treatment. *Russ J Genet* 2014;50(8):798-806. doi: 10.1134/s1022795414080092