





Blood Transfusion and the Risk of Cancer in the US Population: Is There an Association?

This article was published in the following Dove Press journal:
Clinical Epidemiology

Amr Ehab El-Qushayri ^{1,*}
Sherief Ghozy ^{2,*}
Sara Morsy ³
Faria Ali⁴
Sheikh Mohammed Shariful Islam ⁵

¹Faculty of Medicine, Minia University, Minia, Egypt; ²Faculty of Medicine, Mansoura University, Mansoura, Egypt; ³Medical Biochemistry and Molecular Biology Department, Faculty of Medicine, Tanta University, Tanta, Egypt; ⁴Department of Internal Medicine, Henry Ford Allegiance Health, Jackson, MI 49201, USA; ⁵Institute for Physical Activity and Nutrition (IPAN), School of Exercise and Nutrition Sciences, Deakin University, Melbourne, Australia

*These authors contributed equally to this work

Purpose: We aimed to test if blood transfusion is a risk factor for the prevalence of cancer. **Patients and Methods:** We conducted secondary analyses using the NHANES database from 1999 to 2016. We included all individuals who received a blood transfusion with known cancer comorbidity (diseased or not). We used univariate logistic regression to identify any possible association between history of blood transfusion and the prevalence of cancer with adjustment for different co-founders was done. Regression results were expressed as odds ratios (ORs) and 95% confidence interval (95% CI) for both adjusted and unadjusted models. **Results:** A total of 48,796 individuals were included in the final analysis: 6333 of them received a blood transfusion, while the other 42,463 individuals did not. In individuals who received a blood transfusion, the most prevalent cancer was breast cancer (3.4%), followed by prostate (3.0%), non-melanoma skin (2.4%) cancers, while non-melanoma skin (1.2%), prostate (1.1%) and breast (1.1%) cancers were the most prevalent in the no transfusion individuals. There was a significant association between the reported history of blood transfusion and the overall prevalence of cancer in both the unadjusted (OR= 3.47; 95% CI= 3.23–0.72; P-value< 0.001) and adjusted model (OR= 1.86; 95% CI= 1.72–0.2.01; P-value< 0.001). On the level of individual cancers, a significant reduction in cancer prevalence was found in patients with breast, cervix, larynx, Hodgkin's lymphoma, melanoma, prostate, skin (non-melanoma), skin (unspecified), soft tissue, testicular, thyroid, and uterine cancers.

Conclusion: Results did not imply any concrete association between cancer risk and history of blood transfusion. These findings would help in debunking the myth of increased cancer risk following blood transfusion.

Keywords: blood transfusion, cancer, association, prevalence

Introduction

Blood transfusion is considered to be an indispensable choice for many people with acute and chronic conditions which results in the improvement of tissue perfusion and oxygen delivery. It is often used as a lifesaving treatment for the correction of physiologic abnormalities in acute anemia or hemorrhage. It is also used in different surgical procedures as an emergency treatment of surgical induced hemorrhage. However, the decision for blood transfusion should be taken carefully and after weighing both risks and benefits to the patient, as beneficial outcomes are coupled with multiple adverse reactions, including, but not limited to, infections, transfusion-associated immune modulations and incompatible host responses.¹

Blood transfusion can be either the transfusion of whole blood or blood components. Component transfusion includes red blood cells (RBC), plasma, platelets,

Correspondence: Sherief Ghozy
Faculty of Medicine, Mansoura University,
Mansoura 35516, Egypt
Email sherief_ghozy@yahoo.com

cryoprecipitates, and fractionated plasma products including intravenous immunoglobulins (IVIGs), albumin, or plasma-derived coagulation factors for replacement.^{1,2} The main indication of RBC transfusion depends on the transfusion trigger, which is the hemoglobin (Hb) level below which the transfusion is needed. It is postulated by the strategy of restrictive transfusion rule of Hb of 7.0 g/dl in critically ill patients, where Hb concentrations are maintained at 7.0 through 9.0 g/dl as per The Transfusion Requirements in Critical Care (TRICC) trial recommendations, with the exception in unstable angina and acute myocardial infarction.³ At this level, the benefit of blood transfusion outweighs its risks. The patient's comorbidities also need to be thoroughly evaluated before the decision is made for RBC transfusion.^{1,2} Plasma transfusion is usually indicated for ascitic patients to provide plasma proteins, mainly albumin.⁴ Furthermore, plasma is administered for different coagulopathies, including idiopathic thrombocytopenic purpura.⁵ The main clinical indication for plasma transfusion is the international normalized ratio (INR) above 2, prothrombin time (PT) above 1.5, and prolonged thromboplastin time (PTT) more than 2 times.^{1,2} Platelet transfusion is indicated in defective platelet number or function. Meanwhile, cryoprecipitate is used mainly for dysfibrinogenemia and hypofibrinogenemia.^{1,2,4} IVIG is frequently used to treat immune paraneoplastic diseases and may also prevent tumor cell migration.⁶

Well documented side effects of blood transfusion include acute hemolytic transfusion reaction caused mainly by ABO mismatch.⁷ Moreover, allergic reactions, ranging from mild urticaria to severe anaphylaxis, are well-documented in the history of medicine. Other causes of high morbidity and mortality following transfusion, are transfusion-related acute lung injury (TRALI), circulatory overload, and transfusion dyspnea.^{7,8} These side effects typically occur within a short time after transfusion.

There is, however, contradicting evidence regarding the long-term effects of blood transfusion in populations with or without co-morbid conditions.⁸ Long-term mortality was evaluated in cardiac patients and was significantly associated with high mortality for up to one year.⁹ Another study found that blood transfusion is a strong predictor of mortality in blunt trauma patients¹⁰ and increased mortality by three folds. A serious potential long-term side effect is the transmission of chronic infectious viral diseases, including Epstein-Barr Virus [EBV], Hepatitis C [HCV], or Human Immunodeficiency Virus [HIV].⁸ In every

2 million units transfer, the risk of HCV or HIV is 1. The potential for West Nile virus transmission is more, as it is transferred in every 350,000 units. However, the risk of infections had a regional difference as evidenced in developed countries where there is less risk of infections.⁸

An immensely interesting side effect currently under investigation is the risk of cancer after transfusion. A study reported that after 5–29 years from a blood transfusion, there is a 26% risk of developing non-Hodgkin's lymphoma [NHL].¹¹ Another study in UK women confirmed these findings and found that there is also an associated increase in liver cancer.¹² In a study based on the Surveillance, Epidemiology, and End Results (SEER) database, there was a high risk for cancer within 1 year after transfusion. After this interval, there was no significant risk for cancer. Of note, specific types of cancer showed a significant increase, including stomach, liver, kidney, lip, lymphoma, leukemia, respiratory system, and gastrointestinal tract [GIT].¹³ However, it still depends on the interval between transfusion and cancer risk. GIT, renal cancer, and lymphoma were seen in the first year. In the second year, leukemia, liver, oral and pharyngeal carcinoma were diagnosed.¹³

On the contrary, a different study claimed that there is no risk of cancer after blood transfusion.¹⁴ Besides, lifestyle habits, including cigarette smoking and alcohol are determining factors for the risk of developing cancer and can be confounding factors in the studies connecting transfusion to cancer diagnosis.¹⁵ In this study, we aimed to examine any possible association between cancer prevalence and history of blood transfusion.

Patients and Methods

Data Source and Variable Selection

A secondary analysis was performed using the NHANES database conducted between the years 1999–2016 using 9 surveys (1999–2000, 2001–2002, 2003–2004, 2005–2006, 2007–2008, 2009–2010, 2011–2012, 2013–2014 and 2015–2016) by the National Center for Health Statistics of the Centers for Disease Control and Prevention (CDC). We retrieved several variables from the database, including age (in years), gender (male or female), marital status (married, widowed, divorced, separated, never married and living with a partner), race (Mexican American, other Hispanic, non-Hispanic white, non-Hispanic African American and other, including multi-racial), educational level (12th grade without a diploma or less, high school

graduate GED or equivalent, some college course or AA degree, college graduate or higher, and other [do not know or refused]), blood transfusion status (received or not), year of blood transfusion (before 1972, 1972–1991 and 1992 to present), cancer status (diseased or not) and cancer type (bladder, blood, bone, brain, breast, cervix [cervical], colon, esophagus [esophageal], gallbladder, kidney, larynx/windpipe, leukemia, liver, lung, lymphoma/Hodgkin's disease, melanoma, mouth/tongue/lip, nervous system, ovary [ovarian], pancreas [pancreatic], prostate, rectum [rectal], skin [non-melanoma], skin [do not know what kind], soft tissue [muscle or fat], stomach, testis [testicular], thyroid, uterus [uterine] and other).

Inclusion and Exclusion Criteria

We included all individuals who received a blood transfusion with known cancer comorbidity (diseased or not). We excluded all individuals who had missing, declined, or did not know whether they had previously received a blood transfusion, in addition to individuals who had missing data or did not know their cancer comorbidity (diseased or not).

Statistical Analysis

All data were analyzed using R software version 3.6.2¹⁶ using the packages (Rcmdr)¹⁷ and (glm2).¹⁸ All variables were represented as frequencies, and percentages with Chi2 test (or Fisher's exact test, as appropriate) was used for testing the difference, according to the blood transfusion status. We used univariate logistic regression to identify any possible association between history of blood transfusion and the prevalence of cancer. Moreover, adjustment for different co-founders was done; including age, gender, race, marital status, and education of the included patients.¹⁹ Regression results were expressed as odds ratios (ORs) and 95% confidence interval (95% CI) for both adjusted and unadjusted models.

Results

Characteristics of the Included Individuals

A total of 48,796 individuals were included in the final analysis; 6333 of them received a blood transfusion, while the other 42,463 individuals did not. About half of the individuals (43.8%), with transfusion history, had received blood following 1992. Moreover, cancer history was reported in 21.4% of the transfusion group, compared to only 7.3% for no transfusion group. In individuals who

received a blood transfusion, the most prevalent cancer was breast cancer (3.4%), followed by prostate (3.0%), non-melanoma skin (2.4%) cancers, while non-melanoma skin (1.2%), prostate (1.1%) and breast (1.1%) cancers were the most prevalent in the no transfusion individuals.

In the blood transfusion group, the most common age group was 70–100 years (41.3%) followed by 60–69 years (22.2%), 50–59 years (15.1%), and 40–49 years (10.7%), respectively. Moreover, there was a significant difference in age groups when comparing the included individuals according to their blood transfusion history. A similar difference was found in gender distribution, with a male percentage of 39.9% and 49.2% in individuals with a history of blood transfusion and no history, respectively. As presented in [Table 1](#), all other characteristics show significant differences among included individuals when compared on basis of blood transfusion history; including marital status, race, education, year of blood transfusion, overall cancer prevalence, and specific cancer type prevalence.

Association with the Risk of Cancer

There was a significant association between the reported history of blood transfusion and overall prevalence of cancer; this was evident in both unadjusted (OR [95% CI] = 3.47 [3.23–3.72]; P-value < 0.001) and adjusted models (OR [95% CI] = 1.86 [1.72–2.01]; P-value < 0.001). This association appears to be more related to blood transfusions done following the year 1992 (OR [95% CI] = 1.25 [1.07–1.45]; P-value = 0.005). On adjusting for different co-founding factors, an increase in the association level was detected in transfusion done following 1992 (OR [95% CI] = 1.63 [1.38–1.92]; P-value < 0.001), as well as the association for transfusion years 1972–1991 turned into a significant level (OR [95% CI] = 1.33 [1.11–1.60]; P-value = 0.002) ([Table 2](#)).

On the level of individual cancers, there was an increase in the prevalence rate among adjusted (OR [95% CI] = 1.82 [0.75–4.44]) and unadjusted (OR [95% CI] = 1.34 [0.56–3.2]) models of bone cancer and adjusted model of liver cancer (OR [95% CI] = 1.04 [0.37–2.93]). Noteworthy, a significant reduction in cancer prevalence was found in patients with breast, cervix, larynx, Hodgkin's lymphoma, melanoma, prostate, skin (non-melanoma), skin (unspecified), soft tissue, testicular, thyroid, and uterine cancers ([Supplementary Table 1](#)). [Figure 1](#) shows the graphical presentation of both adjusted and unadjusted models for all cancer types.

Table 1 Characteristics of the Included Individuals

Variables; n (%)	Received Blood Transfusion		P value
	Yes (6333)	No (42,463)	
Age (years)			<0.001*
20:29	258 (4.1)	8431 (19.9)	
30:39	421 (6.6)	7920 (18.7)	
40:49	679 (10.7)	7493 (17.6)	
50:59	954 (15.1)	6116 (14.4)	
60:69	1407 (22.2)	6225 (14.7)	
70:100	2614 (41.3)	6278 (14.8)	
Gender			<0.001*
Male	2525 (39.9)	20,873 (49.2)	
Female	3808 (60.1)	21,590 (50.8)	
Marital status			<0.001*
Married	3179 (50.2)	22,212 (52.3)	
Widowed	1267 (20.0)	3171 (7.5)	
Divorced	807 (12.7)	4052 (9.5)	
Separated	199 (3.1)	1388 (3.3)	
Never married	559 (8.8)	7921 (18.7)	
Living with partner	241 (3.8)	3222 (7.6)	
Race			<0.001*
Mexican American	849 (13.4)	7920 (18.7)	
Other Hispanic	436 (6.9)	3526 (8.3)	
Non-Hispanic White	3395 (53.6)	18,543 (43.7)	
Non-Hispanic Black	1338 (21.1)	8773 (20.7)	
Other Race - Including Multi-Racial	315 (5.0)	3701 (8.7)	
Education			<0.001*
12th grade without diploma or less	2042 (32.2)	11,805 (27.8)	
High school graduate GED or equivalent	1470 (23.2)	9776 (23.0)	
Some college courses or AA degree	1724 (27.2)	11,659 (27.5)	
College graduate or higher	1084 (17.1)	9147 (21.5)	
Other	12 (0.2)	72 (0.2)	
Year received blood transfusion			
Before 1972	1476 (23.3)	NA	
1972–1991	1872 (29.6)		
1992 to present	2777 (43.8)		
Cancer			<0.001*
Yes	1357 (21.4)	3096 (7.3)	
No	4976 (78.6)	39,367 (92.7)	

(Continued)

Table 1 (Continued).

Variables; n (%)	Received Blood Transfusion		P value
	Yes (6333)	No (42,463)	
Type of cancer			<0.001*
Bladder	48 (0.8)	61 (0.1)	
Blood	3 (0.0006)	5 (0.0001)	
Bone	17 (0.3)	12 (0.0003)	
Brain	6 (0.1)	10 (0.0003)	
Breast	214 (3.4)	476 (1.1)	
Cervix (cervical)	60 (0.9)	235 (0.6)	
Colon	138 (2.2)	152 (0.4)	
Esophagus (esophageal)	7 (0.1)	13 (0.0003)	
Gallbladder	0 (0.0)	1 (0.0)	
Kidney	38 (0.6)	36 (0.1)	
Larynx/windpipe	5 (0.1)	19 (0.0004)	
Leukemia	16 (0.3)	27 (0.1)	
Liver	10 (0.2)	10 (0.0003)	
Lung	46 (0.7)	67 (0.2)	
Lymphoma/Hodgkin's disease	30 (0.5)	61 (0.1)	
Melanoma	57 (0.9)	191 (0.4)	
Mouth/tongue/lip	7 (0.1)	18 (0.0005)	
Nervous system	0 (0.0)	1 (0.0)	
Ovary (ovarian)	37 (0.6)	62 (0.1)	
Pancreas (pancreatic)	3 (0.0)	8 (0.0002)	
Prostate	188 (3.0)	486 (1.1)	
Rectum (rectal)	8 (0.1)	11 (0.0003)	
Skin (non-melanoma)	154 (2.4)	508 (1.2)	
Skin (do not know what kind)	107 (1.7)	243 (0.6)	
Soft tissue (muscle or fat)	2 (0.0004)	8 (0.0002)	
Stomach	16 (0.3)	20 (0.0004)	
Testis (testicular)	6 (0.1)	22 (0.1)	
Thyroid	16 (0.3)	63 (0.1)	
Uterus (uterine)	56 (0.9)	130 (0.3)	
Others	57 (0.9)	121 (0.3)	

Note: *Statistically significant.
Abbreviation: NA, not applicable.

Discussion

In this study, we investigated the association between the prevalence of cancer and the history of blood transfusion using a large cohort database. Our results did not imply any concrete association between cancer risk and history of blood transfusion. Blood transfusions are associated

Table 2 Logistic Regression of the Association Between Blood Transfusion and Cancer Prevalence

Predictor	Estimate \ddagger	SE	Z	P-value	Odds Ratio	95% Confidence Interval	
						Lower	Upper
Blood transfusion (unadjusted)							
No	Reference						
Yes	1.24	0.04	34.67	<0.001*	3.47	3.23	3.72
Blood transfusion (adjusted\ddagger)							
No	Reference						
Yes	0.62	0.04	15.91	<0.001*	1.86	1.72	2.01
Year of blood transfusion (unadjusted)							
Before 1972	Reference						
1972–1991	–0.09	0.09	–0.98	0.328	0.92	0.77	1.09
1992 to present	0.22	0.08	2.78	0.005*	1.25	1.07	1.45
Year of blood transfusion (adjusted\ddagger)							
Before 1972	Reference						
1972–1991	0.29	0.09	3.03	0.002*	1.33	1.11	1.60
1992 to present	0.49	0.08	5.73	<0.001*	1.63	1.38	1.92

Notes: \ddagger Estimates represent the log odds of “Cancer = Yes” vs. “Cancer = No”; \ddagger Adjusted for age, gender, race, marital status, and education of the included patients; *Statistically significant.

with both risks and benefits. In transfusion medicine, the current hypothesis on the mechanism of carcinogenesis points toward the synergism of immunosuppression and the existence of viruses as the main risk factors.¹⁸ There may be an involvement of unknown genetic mechanisms, that may make susceptible genotypes behave differently after a blood transfusion.¹⁸

Our analysis revealed an overall increased rate of cancer among patients with a history of blood transfusion, and these rates were higher following 1972. However, there was no significant increased rate of individual cancer types in patients who received a blood transfusion. In addition, in the unadjusted analysis, there was a significantly decreased prevalence of breast, cervix, larynx, lymphoma, melanoma, prostate, skin, testis, thyroid, and uterus. Furthermore, after adjustment for age, gender, race, marital status, and education of the included patients, there was still a decreased prevalence of breast, cervix, prostate, melanoma, and thyroid cancers.

There is still a controversy over the risk of cancer in patients receiving blood transfusion.^{7,8} In our study, we did not find any significant risk of cancer in those patients. Our results partially coincide with the results of another study that found that there was no increased risk of cancer after four years of transfusion.¹³ In the same study, they

found that there was an increased risk of liver cancer within 13–30 years, but this increased risk was eliminated after adjusting for the patients’ characteristics.¹³

Notwithstanding, our study results contradict the results of similar studies. In a large cohort of UK women, blood transfusion was associated with increased risk of liver and non-Hodgkin lymphoma.¹² They found that the risk increased within five years for most cancers suggesting subclinical cancer foci. However, after five years, an increase was noted only in liver cancer and non-Hodgkin lymphoma.¹² The same results were obtained in the Danish study, which found that there was an increased risk of liver and non-Hodgkin lymphoma.¹⁵ Another study, based on the SEER registry, implied that there was an increased overall risk of cancer within the first year after blood transfusion. There was also a site-specific increased risk of stomach, colon, liver, kidney, renal pelvis and ureter, myeloma, leukemia, and leukemia.¹³

All studies that investigated the risk of cancer after blood transfusion found that there is a temporal dependent risk association.^{11–15,20} The risk decreases over time and after a specific period, there was no risk of cancer. This was explained as a reverse causation relationship and that the increased risk of cancer after transfusion does not necessarily mean that the transfusion is the cause of




Figure 1 Showing the unadjusted and adjusted odds ratio (OR) and the corresponding 95% confidence interval (CI) of the association between blood transfusion and cancer site.

cancer.¹³ Precancerous lesions or cancer itself may lead to anemia that is treated through blood transfusion, which explains the increased risk for specific cancers.^{13,21} Other causes should be considered including lifestyle habits of alcohol, smoking, and the presence of occult cancer foci.^{8,12,22} In another study, the increased cancer risk was mainly attributed to the transfer of carcinogenic infectious agents like EBV, HIV, and HCV. This explains why there was an increased risk for specific cancers, such as lymphoma and liver cancer.²²

At this juncture, it is very important to discuss the speculated immunomodulatory effect of blood transfusion, referred to as transfusion-induced immune modulation (TRIM). A study of the change in specific immunological functions in human subjects and mice following blood transfusion, has shown decreases in interleukin 2 (IL-2) secretion, natural killer (NK) cell activity and thus, reduced macrophage activity, abnormal CD4/CD8 ratios and immune surveillance against malignancy, and could be either antigen-specific/non-specific in response. This ends up indirectly enabling malignant proliferation and cell growth, tethering, and dissemination.⁶ The role of viruses themselves has also been implicated as a cause of immunosuppression after blood transfusion.²³ Ergo, immune suppression, and

viruses are most likely intertwined as cofactors in the controversial theory of carcinogenesis.^{23,24}

The current study has some limitations; we performed a secondary analysis using cross-sectional data where the temporal relationship could not be investigated, to validate the findings. Secondly, the data for blood transfusion frequency or quantity may have been skewed by the recall bias of patients. Lastly, we were not able to exclude the criteria of other possible risk factors of cancer e.g. smoking, alcohol, and chronic immunosuppressive medications, which may influence the results as confounding clinical and pathological factors.

Conclusion

Our study results did not imply any concrete association between cancer risk and history of blood transfusion. Our results may have important implications in cancer biology, clinical medicine, and public health of debunking the myth of increased cancer risk following blood transfusion. Efforts to surgically minimize blood loss, implement alternative strategies for blood replacement and restrictive transfusion criteria must also continue meanwhile.

Ethics Approval and Informed Consent

The survey protocol was approved by the Institutional Review Board of the Centers for Disease Control and Prevention (Protocol #98-12, Protocol #2005-06, Continuation of Protocol #2005-06, Protocol #2011-17). Written informed consent was obtained from all subjects.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Yaddanapudi S, Yaddanapudi L. Indications for blood and blood product transfusion. *Indian J Anaesth.* 2014;58(5):538–542. doi:10.4103/0019-5049.144648
2. Nuttall GA, Brost BC, Connis RT, et al. Practice guidelines for perioperative blood transfusion and adjuvant therapies: an updated report by the American Society of anesthesiologists task force on perioperative blood transfusion and adjuvant therapies. *Anesthesiology.* 2006;105(1):198–208. doi:10.1097/00000542-200607000-00030
3. Hébert PC, Wells G, Blajchman MA, et al. A multicenter, randomized, controlled clinical trial of transfusion requirements in critical care. *N Engl J Med.* 1999;340(6):409–417. doi:10.1056/NEJM199902113400601
4. O'Shaughnessy DF, Atterbury C, Bolton Maggs P, et al. Guidelines for the use of fresh-frozen plasma, cryoprecipitate and cryosupernatant. *Br J Haematol.* 2004;126(1):11–28. doi:10.1111/j.1365-2141.2004.04972.x
5. Sharma S, Sharma P, Tyler LN. Transfusion of blood and blood products: indications and complications. *Am Fam Physician.* 2011;83(6):719–724.
6. Goubran HA, Elemary M, Radosevich M, Seghatchian J, El-Ekiaby M, Burnouf T. Impact of transfusion on cancer growth and outcome. *Cancer Growth Metastasis.* 2016;9:CGM.S32797. doi:10.4137/CGM.S32797
7. Sahu S, Hemlata H, Verma A. Adverse events related to blood transfusion. *Indian J Anaesth.* 2014;58(5):543–551. doi:10.4103/0019-5049.144650
8. Rawn J. The silent risks of blood transfusion. *Curr Opin Anaesthesiol.* 2008;21(5):664–668. doi:10.1097/ACO.0b013e32830f1fd1
9. Shishehbor MH, Madhwal S, Rajagopal V, et al. Impact of blood transfusion on short- and long-term mortality in patients with ST-segment elevation myocardial infarction. *JACC Cardiovasc Interv.* 2009;2(1):46–53. doi:10.1016/j.jcin.2008.09.011
10. Charles A, Shaikh AA, Walters M, Huehl S, Pomerantz R. Blood transfusion is an independent predictor of mortality after blunt trauma. *Am Surg.* 2007;73(1):1–5. doi:10.1177/000313480707300101
11. Cerhan JR, Engels EA, Cozen W, et al. Blood transfusion, anesthesia, surgery and risk of non-hodgkin lymphoma in a population-based case-control study. *Int J Cancer.* 2008;123(4):888–894. doi:10.1002/ijc.23561
12. Yang TO, Cairns BJ, Reeves GK, Green J, Beral V, collaborators MWS. Cancer risk among 21st century blood transfusion recipients. *Ann Oncol.* 2017;28(2):393–399. doi:10.1093/annonc/mdw555
13. Riedl R, Engels EA, Warren JL, Berghold A, Ricker W, Pfeiffer RM. Blood transfusions and the subsequent risk of cancers in the United States elderly. *Transfusion.* 2013;53(10):2198–2206. doi:10.1111/trf.12071
14. Edgren G, Hjalgrim H, Reilly M, et al. Risk of cancer after blood transfusion from donors with subclinical cancer: a retrospective cohort study. *Lancet.* 2007;369(9574):1724–1730. doi:10.1016/S0140-6736(07)60779-X
15. Hjalgrim H, Edgren G, Rostgaard K, et al. Cancer incidence in blood transfusion recipients. *J Natl Cancer Inst.* 2007;99(24):1864–1874. doi:10.1093/jnci/djm248
16. R Core Team (2014). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available from: <https://www.r-project.org/>. Accessed September 30, 2020.
17. Rcmdr: R commander. Version 2.6-2. 2020. Available from: <https://CRAN.R-project.org/package=Rcmdr>. Accessed March 22, 2020.
18. glm2: fitting generalized linear models. Version 1.2.1. 2018. Available from: <https://CRAN.R-project.org/package=glm2>. Accessed March 22, 2020.
19. Jiang H, Kulkarni PM, Mallinckrodt CH, Shurzinske L, Molenberghs G, Lipkovich I. Covariate adjustment for logistic regression analysis of binary clinical trial data. *Stat Biopharm Res.* 2017;9(1):126–134. doi:10.1080/19466315.2016.1234973
20. Cerhan JR, Wallace RB, Folsom AR, Potter JD, Munger RG, Prineas RJ. Transfusion history and cancer risk in older women. *Ann Intern Med.* 1993;119(1):8–15. doi:10.7326/0003-4819-119-1-199307010-00002
21. Edgren G, Bagnardi V, Bellocco R, et al. Pattern of declining hemoglobin concentration before cancer diagnosis. *Int J Cancer.* 2010;127(6):1429–1436. doi:10.1002/ijc.25122
22. Martin D, Gutkind JS. Human tumor-associated viruses and new insights into the molecular mechanisms of cancer. *Oncogene.* 2008;27(Suppl 2):S31–S42. doi:10.1038/onc.2009.351
23. Kirkley SA. Proposed mechanisms of transfusion-induced immunomodulation. *Clin Diagn Lab Immunol.* 1999;6(5):652–657. doi:10.1128/CDLI.6.5.652-657.1999
24. Moore PS, Chang Y, Condit RC. Common commensal cancer viruses. *PLoS Pathog.* 2017;13(1):e1006078. doi:10.1371/journal.ppat.1006078

Clinical Epidemiology

Publish your work in this journal

Clinical Epidemiology is an international, peer-reviewed, open access, online journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification,

Submit your manuscript here: <https://www.dovepress.com/clinical-epidemiology-journal>

Dovepress

systematic reviews, risk & safety of medical interventions, epidemiology & biostatistical methods, and evaluation of guidelines, translational medicine, health policies & economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.