



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Reducing mortality from 2019-nCoV: host-directed therapies should be an option

The number of confirmed cases of the 2019 novel coronavirus (2019-nCoV) reported to WHO continues to rise worldwide.¹ As with two other WHO Blueprint priority coronaviruses, SARS-CoV² and MERS-CoV,³ 2019-nCoV is lethal. As of Feb 3, 2020, 2019-nCoV has caused 362 deaths out of 17 391 confirmed cases reported to WHO.¹ No specific anti-viral treatment exists. The mainstay of clinical management is largely symptomatic treatment, with organ support in intensive care for seriously ill patients. The unprecedented flurry of activity by WHO and other global public health bodies has mainly focused on preventing transmission, infection control measures, and screening of travellers. The development of vaccines has received immediate funding; however, as with SARS-CoV and MERS-CoV, support for developing treatments for 2019-nCoV that reduce mortality has not been forthcoming. There is an urgent need for focusing funding and scientific investments into advancing novel therapeutic interventions for coronavirus infections.

All three coronaviruses induce excessive and aberrant non-effective host immune responses that are associated with severe lung pathology, leading to death.²⁻⁴ Similar to patients with SARS-CoV and MERS-CoV, some patients with 2019-nCoV develop acute respiratory distress syndrome (ARDS) with characteristic pulmonary ground glass changes on imaging. In most moribund patients, 2019-nCoV infection is also associated with a cytokine storm, which is characterised by increased plasma concentrations of interleukins 2, 7, and 10, granulocyte-colony stimulating factor, interferon- γ -inducible protein 10, monocyte chemoattractant protein 1, macrophage inflammatory protein 1

alpha, and tumour necrosis factor α .²⁻⁶ In those who survive intensive care, these aberrant and excessive immune responses lead to long-term lung damage and fibrosis, causing functional disability and reduced quality of life.^{7,8}

Specific drugs to treat 2019-nCoV will take several years to develop and evaluate. In the meantime, a range of existing host-directed therapies that have proven to be safe⁹⁻¹¹ could potentially be repurposed to treat 2019-nCoV infection. Several marketed drugs with excellent safety profiles such as metformin, glitazones, fibrates, sartans, and atorvastatin, as well as nutrient supplements and biologics could reduce immunopathology, boost immune responses, and prevent or curb ARDS.⁹⁻¹¹ Zinc and other metal-containing formulations appear to have anti-viral activity,¹² are safe, cheap, and readily available. These formulations could be used as adjuncts to monotherapy or as combinational therapies with cyclosporine, lopinavir-ritonavir, interferon beta-1b, ribavirin, remdesivir, monoclonal antibodies, and anti-viral peptides targeting 2019-nCoV.¹¹ Tocilizumab, a monoclonal antibody that targets the interleukin 6 receptor, has a good safety profile. Monoclonal and polyclonal antibodies to 2019-nCoV could be developed for post-exposure prophylaxis.

Ongoing trials of cellular therapies for treatment of ARDS could be expanded to treatment of seriously ill patients with 2019-nCoV infection. Cellular therapy,¹³ using mesenchymal stromal cells from allogeneic donors, has been shown to reduce non-productive inflammation and affect tissue regeneration and is being evaluated in phase 1/2 trials in patients with ARDS (NCT02804945; NCT03608592). Infection with 2019-nCoV appears to be initially associated with an increased Th2 response,⁴ which might reflect a physiological reaction to curb overt inflammatory responses, a clinical phenomenon that guided the optimal

timing of interferon treatment in patients with sepsis, resulting in increased survival.¹⁴ Interleukin 17 blockade might benefit those patients who have a 2019-nCoV infection and increased plasma concentration of interleukin 17.

The isolation and short-term expansion of anti-viral directed T cells has been proven to be a life-saving procedure in patients after autologous hematopoietic stem-cell transplantation with cytomegalovirus infection.¹⁵ Expansion of anti-2019-nCoV-specific T cells, as cellular drugs, could aid to prepare T-cell products for the adjunct treatment of patients with severe 2019-nCoV infection.

Several unique opportunities to evaluate a range of treatment interventions at the peak of the SARS-CoV and MERS-CoV outbreaks were missed due to avoidable delays and subsequent decline of the numbers of cases, leaving numerous questions about coronavirus pathogenesis unanswered. Disappointingly, treatment trials registered for MERS-CoV are still not complete. As the 2019-nCoV continues to spread and evolve, and the numbers of deaths rise exponentially, advancing new therapeutic development becomes crucial to minimise the number of deaths from 2019-nCoV infection.

We declare no competing interests. AZ is co-principal investigator of the Pan-African Network on Emerging and Re-emerging Infections (PANDORA-ID-NET), funded by the European & Developing Countries Clinical Trials Partnership, supported under Horizon 2020, the EU's Framework Programme for Research and Innovation, and a National Institutes of Health Research senior investigator. MM is a member of the innate immunity advisory group of the Bill & Melinda Gates Foundation, and his work is funded by the Champalimaud Foundation.

*Alimuddin Zumla, David S Hui, Esam I Azhar, Ziad A Memish, *Markus Maeurer*
markus.maeurer@fundacaochampalimaud.pt

Department of Infection, Division of Infection and Immunity, University College London, London, UK (AZ); National Institute for Health Research Biomedical Research Centre, University College London Hospitals NHS Foundation Trust, London,



Published Online
February 5, 2020
[https://doi.org/10.1016/S0140-6736\(20\)30305-6](https://doi.org/10.1016/S0140-6736(20)30305-6)

Submissions should be made via our electronic submission system at <http://ees.elsevier.com/thelancet/>

- UK (AZ); Department of Medicine & Therapeutics, Chinese University of Hong Kong, Prince of Wales Hospital, Shatin, New Territories, Hong Kong, China (DSH); Special Infectious Agents Unit, King Fahd Medical Research Center and Department of Medical Laboratory Technology, Faculty of Applied Medical Sciences, King Abdulaziz University, Jeddah, Saudi Arabia (EIA); Infectious Diseases Division, Department of Medicine and Research Department, Prince Mohamed Bin Abdulaziz Hospital, Ministry of Health, Riyadh, Saudi Arabia (ZAM); College of Medicine, Alfaisal University, Riyadh, Saudi Arabia (ZAM); Immunotherapy Programme, Champalimaud Centre for the Unknown, 1400-038 Lisbon, Portugal (MM); and I Med Clinic, University of Mainz, Mainz, Germany (MM)
- 1 WHO. Novel coronavirus (2019-nCoV) situation report - 14. Feb 3, 2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200203-sitrep-14-ncov.pdf?sfvrsn=f7347413_2 (accessed Feb 3, 2020).
 - 2 Hui DSC, Zumla A. Severe acute respiratory syndrome: historical, epidemiologic, and clinical features. *Infect Dis Clin North Am* 2019; **33**: 869–89.
 - 3 Azhar EI, Hui DSC, Memish ZA, Drosten C, Zumla A. The Middle East respiratory syndrome (MERS). *Infect Dis Clin North Am* 2019; **33**: 891–905.
 - 4 Huang C, Wang Y, Li X. Clinical features of patients infected with 2019 coronavirus in Wuhan, China. *Lancet* 2020; published online Jan 24. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
 - 5 Li G, Fan Y, Lai Y, et al. Coronavirus infections and immune responses. *J Med Virol* 2020; published online Jan 25. DOI:10.1002/jmv.25685.
 - 6 Channappanavar R, Perlman S. Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. *Semin Immunopathol* 2017; **39**: 529–39.
 - 7 Batawi S, Tarazan N, Al-Raddadi R, et al. Quality of life reported by survivors after hospitalization for Middle East respiratory syndrome (MERS). *Health Qual Life Outcomes* 2019; **17**: 101.
 - 8 Ngai JC, Ko FW, Ng SS, To KW, Tong M, Hui DS. The long-term impact of severe acute respiratory syndrome on pulmonary function, exercise capacity and health status. *Respirology* 2010; **15**: 543–50.
 - 9 Zumla A, Chan JF, Azhar EI, Hui DS, Yuen KY. Coronaviruses - drug discovery and therapeutic options. *Nat Rev Drug Discov* 2016; **15**: 327–47.
 - 10 Beigel JH, Nam HH, Adams PL, et al. Advances in respiratory virus therapeutics—a meeting report from the 6th ISIRV antiviral group conference. *Antiviral Res* 2019; **167**: 45–67.
 - 11 Zumla A, Azhar EI, Arabi Y, et al. Host-directed therapies for improving poor treatment outcomes associated with the Middle East respiratory syndrome coronavirus infections. *Int J Infect Dis* 2015; **40**: 71–74.
 - 12 Barnard DL, Wong MH, Bailey K, et al. Effect of oral gavage treatment with ZnAL42 and other metallo-ion formulations on influenza A H5N1 and H1N1 virus infections in mice. *Antivir Chem Chemother* 2007; **18**: 125–32.
 - 13 Horie S, Gonzalez HE, Laffey JG, Masterson CH. Cell therapy in acute respiratory distress syndrome. *J Thorac Dis* 2018; **10**: 5607–20.
 - 14 Davies R, O’Dea K, Gordon A. Immune therapy in sepsis: are we ready to try again? *J Intensive Care Soc* 2018; **19**: 326–44.
 - 15 Lérias JR, Paraschoudi G, Silva I, et al. Clinically relevant immune responses against cytomegalovirus: implications for precision medicine. *Int J Mol Sci* 2019; **20**: 1986.