

Preventive Properties of Ramelteon against Cocaine-Induced Autophagia and Apoptosis: A Hypothetic Role of TNF- α Receptor Involvement and JNK/Bcl-2-Beclin1 or Bcl-2/Bax Signaling Pathway

Cocaine is a behavioral stimulant that has greatly increased its use in recent years, according to some reports that the misuse of this drug induces apoptosis and autophagy through the mechanism of neuroinflammatory and oxidative stress, which ultimately leads to neurodegeneration and neural cell death.^[1,2] It has also been shown that the activation of inflammatory receptors such as the tumor necrosis factor- α (TNF- α) receptor (TNFR) causes the activation of c-Jun N-terminal kinase (JNK) leading to Bcl-2 phosphorylation, which causes the dissociation of Bcl-2 from the Bcl-2/Beclin1 or Bcl-2/Bax complex, causing both apoptosis and autophagia, and thus causing cell death.^[3,4] There is evidence that some parts of cocaine-induced neurodegeneration have been mediated by TNFR, but downstream apoptosis and autophagy have not been approved.^[5,6] The neuroprotective strategy for the management of cocaine-induced neurodegeneration by modulating the neural-inflammatory signaling pathway and the apoptosis and autophagia processes referred to by a novel neuroprotective agent is continually superior to any other therapeutic approach. It is, therefore, necessary to explain and introduce or develop a new neuroprotective agent. Previous studies have shown that ramelteon, as a melatonin agonist, can act as a neuroprotective agent and inhibit the neuroinflammatory process, but its effects on TNFR and the apoptosis and autophagic signaling pathway have not yet been clarified.^[7,8] Therefore, according to the above-mentioned literature, we believed that the use of ramelteon in cocaine-addicted subjects would provide neuroprotection against cocaine-prompted neuro-inflammatory processes and would suggest TNFR-Bcl-2/Beclin1 or TNFR-Bcl-2/Bax signaling pathways and, thus, possibly inhibit cocaine-prompted autophagy and apoptosis, eventually preventing cocaine-prompted cell death [Figure 1]. Nevertheless, further study is needed in order to clarify this suggestion.

Author's contribution as follow

Conception and design of the hypothesis was done by M.M.

Interpretation of data and review of the literature was done by N.K and F.M.

Drafting the work or revising was done by M.M and S.S.

Designing the figure was done by R.D.

**Niyoosha Kandezi, Fatemeh Majdi¹,
Reza Davoudizadeh, Majid Motaghinejad,
Sepideh Safari**




Figure 1: Ramelteon, as a melatonin receptor agonist, can induce inhibition of the effects of cocaine by activating the TNF- α receptor (TNFR)/JNK/Bcl-2-Beclin1 or TNFR/Bcl-2/Bax signaling pathway and cell death caused by it. Melatonin receptor activation by both melatonin and ramelteon also causes mitochondrial biogenesis activation and may cause direct inhibition of JNK/Bcl-2-Beclin1 or JNK/Bcl-2/Bax signaling pathways, which may reduce apoptosis, oxidative stress and inflammation. TNFR: TNF- α receptor, MTR: Melatonin receptor, Ra: Ramelteon, MT: Melatonin

Razi Drug Research Center, Iran University of Medical Sciences, Tehran, Iran, ¹Department of Pharmaceutical Chemistry, Faculty of Pharmaceutical Chemistry, Pharmaceutical Sciences Branch, Islamic Azad University, Tehran, Iran

Address for correspondence:

Dr. Majid Motaghinejad,
Tehran, Sheykhfazloohah Hemmat High Way Iran University of Medical Sciences, P.O. Box: 14496-14525, Iran.
E-mail: Motaghinezhad.m@iums.ac.ir

Received: 30 Nov 19 Accepted: 03 Jan 20

Published: 16 Mar 20

References

- Riezzo I, Fiore C, De Carlo D, Pascale N, Neri M, Turillazzi E, et al. Side effects of cocaine abuse: Multiorgan toxicity and pathological consequences. *Curr Med Chem* 2012;19:5624-46.
- Jorge Dinis-Oliveira R, Carvalho F, Alberto Duarte J, Brandao Proenca J, Santos A, Magalhães T. Clinical and forensic signs related to cocaine abuse. *Curr Drug Abuse Rev* 2012;5:64-83.
- Olmos G, Lladó J. Tumor necrosis factor alpha: A link between neuroinflammation and excitotoxicity. *Mediators Inflamm* 2014;2014:861231.
- Poon HF, Abdullah L, Mullan MA, Mullan MJ, Crawford FC. Cocaine-induced oxidative stress precedes cell death in human neuronal progenitor cells. *Neurochem Int* 2007;50:69-73.
- de Oliveira MR, Jardim FR. Cocaine and mitochondria-related signaling in the brain: A mechanistic view and future directions. *Neurochem Int* 2016;92:58-66.

6. Dey S, Snow DM. Cocaine exposure *in vitro* induces apoptosis in fetal locus coeruleus neurons through TNF- α -mediated induction of Bax and phosphorylated c-Jun NH2-terminal kinase. *J Neurochem* 2007;103:542-56.
7. Srinivasan V, Kaur C, Pandi-Perumal S, Brown GM, Cardinali DP. Melatonin and its agonist ramelteon in Alzheimer's disease: Possible therapeutic value. *Int J Alzheimers Dis* 2011;2011:741974.
8. Miller E, Morel A, Saso L, Saluk J. Melatonin redox activity. Its potential clinical applications in neurodegenerative disorders. *Curr Top Med Chem* 2015;15:163-9.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code: 	Website: www.ijpvmjournal.net/www.ijpm.ir
	DOI: 10.4103/ijpvm.IJPVM_446_19

How to cite this article: Kandezi N, Majdi F, Davoudizadeh R, Motaghinejad M, Safari S. Preventive properties of ramelteon against cocaine-induced autophagia and apoptosis: A hypothetic role of TNF- α receptor involvement and JNK/Bcl-2-Bcl-1 or Bcl-2/Bax signaling pathway. *Int J Prev Med* 2020;11:36.