¹Institute of Health Policy, Management and Evaluation, University of Toronto, ON, Canada; ²Li Ka Shing Knowledge Institute, St. Michael's Hospital, Toronto, ON, Canada; ³Department of Pathology and Laboratory Medicine, University of Toronto, Toronto, ON, Canada; ⁴Mount Sinai Hospital, Laboratory Medicine and Pathobiology, Sinai Health System, University of Toronto, Toronto, ON, Canada; ⁵Lunenfeld-Tanenbaum Research Institute, Sinai Health System, Toronto, ON, Canada; ⁶Laboratory for Molecular Medicine Cambridge, Partners Healthcare for Personalized Medicine, Cambridge, MA, USA; ⁷Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA. Correspondence: Jordan Lerner-Ellis@sinaihealthsystem.ca)

Published online: 2 May 2019

Correction: Evaluation of the cost and effectiveness of diverse recruitment methods for a genetic screening study

Hila Milo Rasouly¹, Julia Wynn², Maddalena Marasa¹, Rachel Reingold¹, Debanjana Chatterjee¹, Sheena Kapoor¹, Stacy Piva¹, Byum Hee Kil¹, Xueru Mu¹, Maria Alvarez¹, Jordan Nestor¹, Karla Mehl¹, Anya Revah-Politi³, Natalie Lippa³, Michelle E. Ernst³, Louise Bier³, Aileen Espinal², Bianca Haser², Anoushka Sinha⁴, Ian Halim⁴, David Fasel⁵, Nicole Cuneo¹, Jacqueline J. Thompson⁴, Miguel Verbitsky¹, Elizabeth G. Cohn⁶, Jill Goldman⁶, Karen Marder⁶, Robert L. Klitzman⁷, Manuela A. Orjuela^{2,8}, Yat S. So⁵, Alex Fedotov⁹, Katherine D. Crew¹, Krzysztof Kiryluk¹, Paul S. Appelbaum⁷, Chunhua Weng⁵, Karolynn Siegel¹⁰, Ali G. Gharavi¹ and Wendy K. Chung^{1,2}

Genetics in Medicine (2019) 21:2407; https://doi.org/10.1038/s41436-019-0528-8

Correction to: Genet Med 2019; https://doi.org/10.1038/s41436-019-0497-y published online 1 April 2019

The original version of this Article contained an error in the undergraduate degree awarded to the author lan Halim, which was incorrectly given as BS. This has now been corrected to BA in both the PDF and HTML versions of the Article.

Published online: 1 May 2019

Open

Correction: Sequencing as a first-line methodology for cystic fibrosis carrier screening

Kyle A. Beauchamp 1,2, Katherine A. Johansen Taber, Peter V. Grauman, Lindsay Spurka, Jeraldine Lim-Harashima, Ashley Svenson 1, James D. Goldberg and Dale Muzzey 1,2

Genetics in Medicine (2019) 21:2407-2408; https://doi.org/10.1038/s41436-019-0543-9

Correction to: Genetics in Medicine; https://doi.org/10.1038/s41436-019-0525-y, published online 30 April 2019

The original version of this Article contained an error in Fig. 3. Specifically, the result "3 (67%) TOP" should read "2 (67%) TOP." This has now been corrected in both the PDF and HTML versions of the Article.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use


¹Department of Medicine, Columbia University Medical Center, New York, NY, USA; ²Department of Pediatrics, Columbia University Medical Center, New York, NY, USA; ³Institute for Genomic Medicine, Columbia University Medical Center, New York, NY, USA; ⁴College of Physician & Surgeons, Columbia University, New York, NY, USA; ⁵Department of Biomedical Informatics, Columbia University Irving Medical Center, New York, NY, USA; ⁶Department of Neurology, Aging & Dementia, Columbia University Medical Center, New York, NY, USA; ⁸Department of Epidemiology, Columbia University Medical Center, New York, NY, USA; ⁸Department of Epidemiology, Columbia University Medical Center, New York, NY, USA; ⁸Department of Epidemiology, Columbia University Medical Center, New York, NY, USA; ¹⁰Department of Sociomedical Sciences, Columbia University Mailman School of Public Health, New York, NY, USA. Correspondence: Wendy K. Chung (wkc15@cumc.columbia.edu) These authors contributed equally: Hila Milo Rasouly and Julia Wynn

is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2019

¹Myriad Women's Health (formerly Counsyl), South San Francisco, CA, USA; ²Myriad Genetics, Salt Lake City, UT, USA; ³Present address: Guardant Health, Redwood City, CA, USA; ⁴Invitae, San Francisco, CA, USA. Correspondence: Dale Muzzey (mwh_research@myriad.com)

Published online: 15 May 2019



⁽a) Actions include IVF with PGT-M, prenatal diagnostic testing if/when pregnancy occurs, adoption, or avoidance of pregnancy

Fig. 3

⁽b) Current and subsequent pregnancies in those screened prenatally, and subsequent pregnancies that occured in those screened preconceptionally.