


STUDY PROTOCOL

Open Access



# Suubi4Stigma study protocol: a pilot cluster randomized controlled trial to address HIV-associated stigma among adolescents living with HIV in Uganda

Proscovia Nabunya<sup>1,2\*</sup> , Fred M. Ssewamala<sup>1,2</sup>, Ozge Sensoy Bahar<sup>1,2</sup>, Lynn T. M. Michalopoulos<sup>3,4</sup>, James Mugisha<sup>5</sup>, Torsten B. Neilands<sup>6</sup>, Jean-Francois Trani<sup>2</sup> and Mary M. McKay<sup>2</sup>

## Abstract

**Background:** Sub-Saharan Africa (SSA) is heavily burdened by HIV, with 85% of the global new infections among adolescents happening in the region. With advances in medication and national policies promoting antiretroviral therapy (ART), children < 15 years living with HIV (CLWH) continue to grow with a chronic, highly stigmatized disease. Unfortunately, the stigma they experience results in much lower quality of life, including poor mental health and treatment outcomes. Family members also experience stigma and shame by virtue of their association with an HIV-infected family member. Yet, stigma-reduction interventions targeting CLWH and their families are very limited. The goal of this study is to address HIV-associated stigma among CLWH and their caregivers in Uganda.

**Methods:** This three-arm cluster randomized control trial, known as *Suubi4Stigma*, will evaluate the feasibility, acceptability, and preliminary impact of two evidence-based interventions: (1) group cognitive behavioral therapy (G-CBT) focused on cognitive restructuring and strengthening coping skills at the individual level and (2) a multiple family group (MFG) intervention that strengthens family relationships to address stigma among CLWH ( $N = 90$ , 10–14 years) and their families (dyads) in Uganda. Nine clinics will be randomized to one of three study arms ( $n = 3$  clinics, 30 child-caregiver dyads each): (1) usual care; (2) G-CBT + usual care; and (3) MFG + usual care. Both treatment and control conditions will be delivered over a 3-month period. Data will be collected at baseline (pre-intervention) and at 3 months and 6 months post-intervention initiation.

**Conclusion:** The primary aim of the proposed project is to address the urgent need for theoretically and empirically informed interventions that seek to reduce HIV-associated stigma and its negative impact on adolescent health and psychosocial well-being. As several countries in SSA grapple with care and support for CLWH, this study will lay the foundation for a larger intervention study investigating how HIV-associated stigma can be reduced to foster healthy child development—especially for CLWH as they transition through adolescence.

**Trial registration:** ClinicalTrials.gov: [NCT04528732](https://clinicaltrials.gov/ct2/show/study/NCT04528732); Registered August 27, 2020

**Keywords:** HIV-associated stigma, Children and adolescents living with HIV, Multiple family group, Group cognitive behavioral therapy, Uganda

\*Correspondence: nabunyp@wustledu

<sup>2</sup> Brown School, Washington University in St. Louis, Campus Box 1196, One Brookings Drive, St. Louis, MO 63130, USA

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Introduction

Globally, an estimated 1.8 million children < 15 years are living with HIV [1, 2]. Sub-Saharan Africa (SSA) is heavily burdened by HIV, with 85% of new infections among adolescents happening in the region [2]. Uganda is one of 7 countries in SSA to achieve the 90-90-90 testing, treatment, and viral suppression targets [2, 3]. However, even with these improvements, HIV prevalence is still high (7.5%) among people between 15 and 49 years [4]. Moreover, close to 150,000 children (ages 0–14) were living with HIV in 2019 [2]. While availability and access to free antiretroviral therapy (ART) have decreased child mortality [5], such accomplishment has resulted in the likelihood that more children living with HIV (CLWH) will transition into adulthood with HIV, a chronic, highly stigmatized illness [6]. Unfortunately, the stigma they experience results in a lower quality of life. However, very few stigma-reduction interventions targeting CLWH and their families exist in SSA [7, 8]. Thus, there is a need for data driven research to address stigma, especially among CLWH as they transition through adolescence into young adulthood.

### HIV-related stigma and associated outcomes

Among people living with HIV (PLWH), stigma is a common experience characterized by public blame and moral condemnation for contracting the infection [9–11]. It perpetuates a culture of silence and fear and prevents individuals from testing and seeking health care [9]. Stigma can be internalized as a result of perceived negative public attitudes. It translates into feelings that the self is reprehensible, damaged and defective; and is associated with depression and post-traumatic stress disorders (PTSD) [12, 13], feelings of loneliness and social isolation [14, 15], poor treatment and adherence to medication [10, 16], poor HIV-related physical health [17], and increased sexual risk-taking behavior [18]. Moreover, internalized stigma increases the risk of loss to treatment follow up [19]. Public stigma is manifested by the general population through negative stereotypes such as those related to sexual behaviors, prejudice (fear, aversion, hatred), and discrimination, all of which create social barriers, including access to healthcare [20]. Moreover, many CLWH live with extended family members after losing their parents to HIV, where stigma is perpetuated through rejection, verbal insults, physical abuse, avoidance and ostracism due to unfounded fears of infection [21].

At the family level, family members are often condemned and stigmatized in similar ways, by virtue of their association with an HIV infected family member [22, 23]. Stigma at the family level may be manifested through gossip, name calling, rejection and

social isolation, loss of social support, and harassment [22, 23]. Specifically, family members are often held accountable for not preventing the perceived “immoral behaviors” of the HIV-infected family member, leading to feelings of failure, anger, guilt, and shame [22]. Such feelings negatively affect family caregiving roles, family functioning, and HIV health outcomes for PLWH, including CLWH. Due to this environment, CLWH may miss developing strong attachment bonds with family members and fail to develop self-esteem, emotional, and behavioral regulations [24]. Such unsupportive social environments increase the risk for mental disorders, including depression, stress, and anxiety [25]. Therefore, it is critical to develop HIV stigma reduction interventions to improve life satisfaction, family functioning, and reduce the potential spread of HIV.

### Potential role of family members in addressing HIV-associated stigma

Adolescence is a period of multiple vulnerabilities marked by the onset of physical and emotional maturation accompanied by the challenges of adapting to social, emotional, and cognitive changes [26]. Hence, children need additional support, including emotional support and acceptance from family and community members. Yet, many CLWH cannot count on the “normal” transition to adolescence due to HIV-associated stigma where community and family members ostracize them for being HIV positive [27], and where family members suffer the same treatment due to having a CLWH in the family [22, 23]. Thus, understanding the role of family members, and involving them in the design and implementation of family level HIV-related programs and interventions for CLWH is essential to their success.

The protective role of families in influencing children’s behavior and mental health outcomes has been well documented [28]. Studies have shown that quality of family relationships predict child mental health functioning and overall adjustment, and that when families are consistently involved in children’s lives, children experience a more positive transition through adolescence [29, 30]. Parental skills have been associated with children’s psychological adjustment, less risky sexual behavior, and less susceptibility to peer pressure [31, 32]. Moreover, more frequent and open communication with parents is associated with better psychological adjustment [33, 34]. Yet, for CLWH, parent-child communication and involvement may be adversely affected by orphanhood and HIV-associated stigma. Therefore, family support strategies may strengthen family functioning and help to address individual and family-level stigma.

### Need for HIV stigma reduction interventions targeting CLWH

While the negative impacts of HIV-associated stigma have been well-documented, systematic reviews have found that stigma reduction interventions for CLWH in SSA are almost non-existent [7, 8]. Existing interventions focus on reducing fear of HIV infection among non-infected populations [8] as they interact with PLWH. For example, of the 48 stigma reduction interventions, only 3 aimed to reduce stigma among PLWH in SSA [35, 36]. None targeted CLWH, nor assessed the impact of stigma reduction on HIV-related outcomes. To address these gaps, this pilot trial, entitled “*Suubi4Stigma*” (also known as *Hope for Stigma* in Luganda local language), seeks to address the urgent need for innovative, theoretically, and empirically informed interventions to reduce HIV-associated stigma and its negative impact on adolescent health and psychosocial well-being. This study examines two evidence-based interventions used in mental health settings, schools, and communities: (1) group cognitive behavior therapy (G-CBT) focuses on cognitive restructuring and strengthening coping skills at the individual level; and (2) a multiple family group (MFG) intervention that strengthens family relationships to address stigma among CLWH and their families. The specific aims are the following:

#### Aim 1

Pilot test the feasibility, acceptability, and preliminary impact of G-CBT and MFG on reducing HIV-associated (internalized and family level) stigma, and its impact on adolescent and family outcomes (trauma symptoms, depression, sexual risk, family/social support and adherence to medication) compared to: (a) usual care vs G-CBT; (b) usual care vs MFG; (c) G-CBT vs. MFG.

#### Aim 2

Qualitatively examine participants’ and facilitators’ intervention experiences and identify individual, family, and

institutional-level facilitators and barriers to G-CBT and MFG intervention implementation and participation.

The *Suubi4Stigma* study has been designed as a three-arm cluster randomized control trial (RCT), consisting of a control arm and two treatment arms. Assessments will be conducted at baseline, 3 months, and 6 months post-intervention initiation.


### Theoretical framework

This study is informed by the HIV stigma framework [37, 38] suggesting that HIV stigma impacts PLWH via distinct HIV stigma mechanisms of internalized, anticipated, and enacted HIV stigma. Anticipated and enacted HIV stigma involve experiences with others [38]. Internalized stigma—the focus of this study—involves endorsing negative feelings and beliefs associated with HIV and applying them to the self. In addition, family members of PLWH are also subjected to and experience stigma by association via similar mechanisms. Within this framework, MFG provides opportunities for caregivers and children to communicate in a safe setting. It focuses on addressing internalized and family-level stigma by normalizing shared experiences with other families, fostering peer support and family communication, facilitating optimism and morale, and enhancing interpersonal and coping skills [39].

On the other hand, G-CBT addresses internalized stigma through the core components of psychoeducation, cognitive restructuring, and skill-building to increase adaptive coping mechanisms [40]. These mechanisms may impact a range of psychological, behavioral, and health outcomes for CLWH and their families (Fig. 1).

### Methods

The *Suubi4Stigma* study is a three-arm cluster RCT evaluating the feasibility, acceptability, and preliminary impact of G-CBT versus MFG interventions among, 90 CLWH (10–14 years) and their caregivers (dyads). Nine clinics will be randomized to one of three study arms



**Fig. 1** Conceptual model

(*n* = 3 clinics, 30 child-caregiver dyads each): (1) usual care; (2) G-CBT + usual care; and (3) MFG + usual care. Both treatment and control arms will be delivered over a 3-month period. Data will be collected at baseline (pre-intervention), 3 months, and 6 months post-intervention initiation (Fig. 2).

**Study setting**

The study will be implemented in the greater Masaka Region of Uganda. The region is composed of seven political districts: Rakai, Masaka, Lwengo, Kalungu, Lyantonde, Kyotera, and Bukomansimbi, and has the highest HIV prevalence compared to the national average [3, 4]. Child-caregiver dyads will be recruited from health clinics where the Washington University’s International Center for Child Health and Development (ICHAD) that houses the study and Reach the Youth (RTY), our implementation partner, operate. For a health clinic to be included in the study, it will have to be credited by the Uganda Ministry of Health to provide ART and have adolescent friendly services (e.g., adolescent days).


**Study participants**

A total of 90 CLWH and their caregivers (dyads) enrolled in care at a health clinic that has partnered with ICHAD and RTY will be recruited into the study. Child inclusion

criteria are: (1) living with HIV and know their status, (2) prescribed ART, (3) living within a family (defined broadly—not necessarily with biological parents), and (4) between 10 and 14 years. All eligible children from a particular household will be enrolled in the study and will be assigned to the same study condition. In addition, caregivers of children who agree to participate in the study will be enrolled. If the caregiver chooses to withdraw from the study, the child will also be asked to withdraw from the study. Children and caregivers will be excluded if they are unable to comprehend the informed consent/ assent process and study expectations or unwilling to complete study activities. A complete timeline of project activities is presented in Table 1.

**Clinic randomization**

Stratified random sampling will be utilized to assign health care clinics to two strata based on health clinic level (health center III vs health center IV) to ensure overall clinic balance across study arms. Per the 2019 Uganda Ministry of Health’s Guidelines for Health Center Health Unit Management Committees (<http://library.health.go.ug/publications/health-infrastructure>), Health Center III is amid level care with provisions for basic laboratory services for diagnosis, maternity care and inpatient care with first level referral cover to the subcounty. It is usually



**Fig. 2** Suubi4Stigma study design

**Table 1** Study timeline

Activities	1-3	4-6	7-9	10-12	13-15	16-18	19-21	21-24
Study preparations (hiring & training of research staff, refinement/adaptation of MFG and G-CBT interventions, preparation/refinement of assessment instruments, development of interview protocol, IRB approval)	X							
Participant recruitment and baseline assessments; intervention training for facilitators (MFG and G-CBT)		X						
Intervention delivery (MFG and G-CBT)			X					
3-month assessment (post-intervention), including qualitative interviews				X				
6-month assessment (follow-up)					X			
Quantitative data entry		X	X	X	X	X	X	
Qualitative data transcription and translation				X	X			
Data analysis, report writing, and results dissemination				X	X	X	X	X

staffed by nurse aids, qualified nurses and clinical officers. Health Center IV on the other hand, is a higher-level primary care facility below the district hospital, staffed with a medical doctor, with provisions for operating theater, inpatient and laboratory services and receives referrals from lower-level health facilities. All clinics offer HIV services. Each clinic will be randomly assigned to one of the three study arms, such that all selected child-caregiver dyads in the same clinic receive the same intervention, to reduce contamination. Randomization of clinics will be done by an independent research associate based at Washington University in St. Louis, using STATA software. Each health clinic's assignment to a study condition will remain blinded to other study participants and facilitators, with the exception of research assistants, Project Coordinator and MPIs.

### **Recruitment, retention, and attrition**

Recruitment procedures tested in our previous studies will be utilized [41]. Children will be identified and recruited from health clinics associated with RTY and ICHAD. Patients are seen at least annually and each patient on ART must have prescriptions filled monthly at each of the health clinics. Although appointment days (not times) are provided, most patients arrive early in the morning on days that are convenient for them and wait for several hours before they are seen, providing an opportunity for recruitment through medical staff. A list of all eligible families will be created from medical records by one of the medical staff. Each chart contains data on the patient's HIV status, age, and family. A clinic staff member will review the daily schedule, noting the eligibility of patients. Providers will present the project to adult caregivers of eligible children during appointments. If caregivers are interested, verbal consent to be contacted by research staff (who will be on-site during clinic days) will be requested. After speaking with the research staff one-on-one about the study, interested caregivers will provide written consent for themselves and for their child to participate. Children will be asked to provide written assent separately to avoid coercion.

The Suubi4Stigma study will take place in a highly stable region where geographical moves are rare, facilitating the ability to track and retain the sample. Retention procedures used in our previous Suubi studies will be utilized [41–43]. Specifically, we will ask participants to give their telephone number (if they have one), names, addresses, and contact information for three people who will always know how to reach them. Participants will be told that if we contact the people listed, we will not discuss any details about them or their study participation. We will use these records to help track their location only if we have lost contact. In addition, we will have monthly

contact with children in both treatment conditions during MFG and G-CBT intervention delivery. This frequent contact will enable the research team to continually engage all participants and minimize loss to follow-up. Based on prior studies in the same region, we conservatively expect attrition by end of the 2-year study to be no more than 10% [43, 44].

### **Suubi4Stigma study arms**

#### **Control arm (usual care)**

All participants in both control and treatment arms will receive the traditional clinic intervention that focuses on testing services, as well as medical and treatment support for PLWH, including children and adolescents [45]. Currently, patients coming to the clinics receive testing and ART treatment as well as information about disease management. Both children and caregivers will receive this information. All participants in both control and treatment arms will receive medical and psychosocial support as part of usual care.

#### **Treatment arm 1 (G-CBT)**

In addition to usual care, participants in this arm will receive 10 sessions of G-CBT for HIV-associated stigma. Within G-CBT for stigma, we will utilize core components of CBT, such as psychoeducation, cognitive restructuring, and skill-building to increase adaptive coping mechanisms [40]: (1) exploration of HIV's role and impact of stigma in adolescents' lives; (2) use of cognitive restructuring to identify and address the negative stigma-associated beliefs, loss of self-esteem, and self-blame; and (3) skills-building around stress management and emotion-focused coping strategies to address negative feelings (e.g., assertiveness, relaxation skills and problem solving skills) [46]. G-CBT will be facilitated by trained para-counselors with experience in mental health support and working with children and adolescents living with HIV in the study region. Caregivers will not participate in G-CBT. Sessions will be delivered twice a week, outside of school hours and will be delivered in the Luganda local language. G-CBT is likely to offer more opportunities for normalization, positive peer modeling, reinforcements, social support, and exposure to social situations and feedback sources, given the context of shared experiences [47].

#### **Treatment arm 2 (MFG)**

The MFG intervention is rooted in family systems theory, structural family theory and social learning theory with elements of psychoeducation and social group work. MFG is a family-centered, group-delivered, evidence-informed, strength-based 10-session intervention for adolescents whose families struggle with poverty and

associated stressors [39]. MFG integrates components of existing evidence-based practices that successfully improve parental management, mental health promoting family processes, and family strengthening [48, 49]. Specific session content will draw on the current interventions implemented by ICHAD [42, 50]. Sessions will focus on the core components of MFG, also known as 4Rs and 2S's (rules, responsibility, relationships, respectful communication, stress, and social support). Families (children and caregivers) will be combined into groups of no more than 10 families each to promote communication and support within and among families. Sessions will be delivered in the Luganda local language, will last approximately 1 h and will be delivered twice a week, outside of school hours. Given the significant and protective role families play in children and adolescents' health and mental health, we expect that strengthening family functioning and dialogue by involving caregivers through MFG will lead to better child outcomes, including addressing HIV-associated stigma.

#### **Intervention adaptation, delivery, monitoring, and fidelity**

We will engage stakeholders, including implementing partners, community health care workers and peer parents already trained in the delivery on MFG sessions [42, 50], as well as in-country mental health workers and community leaders, and para-counselors, who will deliver G-CBT sessions. In Uganda, para-counselors are trained to assist with the psychological needs of individuals, including those related to HIV/AIDS and mental health [51, 52]. Specifically, while counselors undergo both generic and specialty training, para-counselors acquire basic general counseling and can offer basic help as they organize for a referral to the professional. They will participate in a 5-day training and receive monthly group supervision. Adaptation of G-CBT content, including cultural adaptation, will be ensured by intensive consultative meetings with the aforementioned stakeholders. During the consultative meetings, stakeholders will be introduced to the study and the core components of G-CBT for stigma, review manual/session content developed by the research team (based on a review of existing literature and CBT manuals utilized in Uganda and other settings), provide feedback on session content and cultural relevance, and make final revisions to the manual. The research team will ensure: (1) content adaptation of the materials to address stigma associated with mental illness to the context of HIV/AIDS-associated stigma, (2) developmental and cognitive adaptation of the content for children between 10 and 14 years, and (3) cultural adaptation of the G-CBT to local context. Content will be tailored to match children's ability to comprehend and implement the therapeutic techniques, including

age-appropriate activities, child friendly materials, simplified language and visuals (e.g., cartoons) [53], less complex behavioral techniques, and more support, structure and feedback [46]. In addition, the cultural adaptation is intended to enhance its treatment relevance, credibility, efficacy, and effectiveness by aligning it to the socioeconomic situation, cultural beliefs, family, political, and health systems in the region [54, 55].

For MFG, parent peers ( $n = 6$ ) and community health workers ( $n = 6$ ) already trained in MFG delivery focused on child behavioral health [42, 43, 50] will be invited to participate and receive a refresher training focused on Suubi4Stigma's new content. Sessions focused on HIV and stigma will be adapted from curriculum used in our previous HIV-focused studies and resources from the Ministry of Health [41, 53]. The protocols have been designed to provide opportunities during each session to directly apply content to the realities of family life, emergent cultural and values perspectives, as well as tailor messages to age of child [43, 56]. These will include group activities, role plays, sharing experiences, and family take home activities. During MFG implementation, facilitators will receive 2 h per month of group supervision across sites.

#### **Implementation process evaluation**

Process measures will be used to monitor fidelity and intervention implementation. For intervention delivery fidelity, independent observations (by trained research staff) using fidelity checklists will be made with a random sample of 60% of intervention sessions for both G-CBT and MFG. These data will be used to assess: (1) the relationship between planned and actual implementation; (2) integrity of implementation; and (3) how sessions were altered to maximize effectiveness and acceptability. In addition, having two facilitators per group will allow for detailed process notes on the implementation process and balancing "fidelity" and "fit" to local culture. Group participants will report monthly on factors that affect their participation using an implementation checklist that assesses satisfaction and obstacles to program delivery: (1) factors interfering with child or family participation (e.g., time, other priorities, stigma); (2) concrete obstacles (e.g., weather, transportation); and (3) site and staffing obstacles (e.g., time constraints).

#### **Quantitative data collection**

Data will be collected at baseline, 3 months, and 6 months post-intervention initiation from both children and caregivers (see Table 2 for assessment measures). Interviewer administered interviews will be conducted by a trained research assistant at each clinic (in a private location) and take approximately 60–90 min with breaks

**Table 2** Assessment table

Variable	Measurement
<b>Respondent: children</b>	
Demographics	Socio-demographic questionnaire
Moderators	Economic/household income, gender, rural/urban/semi-urban
Mental health functioning	Adapted Child Depression Inventory [57] Adapted Tennessee Self-Concept Scale (TSC-2) [58] Beck Hopelessness Scale [59]
Post-traumatic stress disorder	Child PTSD Reaction Index [60]
Family support	Family Environment Scale [61] Family Assessment Measure [62] Adapted Social Support Behaviors Scale (SS-B) [63]
Social support	Friendship Qualities Scale [64]
Stigma and shame	Shame Questionnaire [65] Stigma-by-association scale for children [66] HIV Stigma mechanism measure [67]
Loneliness and social isolation	UCLA Loneliness Scale Version 3 [68]
Sexual risk taking	Adapted Youth AIDS Prevention Project used in CHAMP [69] Questions adapted from Auslander et al. [70] and Slonim-Nevo et al. [71]
Medication adherence	Youth-self-reports [72], viral load, pill counting, pharmacy records
Intervention feedback	Semi structured interviews
<b>Respondent: caregivers</b>	
Stigma and shame	AIDS-related stigma scale [73] Adapted Stigma-by-association measure [66]
Family relations	Family Environment Scale [61] Family Assessment Measure [62]
Mental health functioning	Adapted Brief Symptoms Inventory [74] Parenting Stress Index [75]
Intervention feedback	Semi-structured interviews

and will be administered in Luganda, the local language widely spoken in the study region. Assessments will be translated and back translated into the local language by a certified translator.

#### Qualitative data collection

Our qualitative methods are informed by an interpretive approach that examines the experiences and practices of participants and their subjective meaning making processes [76]. Two sets of qualitative interviews with participants (children and caregivers) in the two treatment arms (G-CBT and MFG) will be conducted. The first set will be conducted at baseline to explore participants' experiences with HIV-related stigma and its perceived impact on their lives and family wellbeing. The second interview will be conducted upon intervention completion to obtain feedback on intervention acceptability and relevance. Specifically, these interviews will (1) explore participants' intervention experiences and perceived impact of the intervention on change mechanisms and (2) identify multi-level facilitators and barriers to G-CBT

and MFG intervention, implementation, and participation. Our interview guides will be informed by the HIV stigma framework and the acceptability and appropriateness concepts developed by Proctor et al. to explore perceptions of intervention acceptability and impact [76]. The interviews will provide rich data on processes and experiences with the program; processes behind key outcomes, mechanisms of change, and mediating variables; and potential individual, family, contextual, programmatic, and structural factors affecting their experiences. Dyads with the highest, medium, and lowest scores on the stigma measures [66, 67, 73] in two treatment arms (6 clinics) will be identified at following baseline. Two families within each of these groups in the 6 treatment clinics will be invited to participate in qualitative interviews ( $2 \times 3 \times 6 = 36$  dyads). This sampling method will ensure that participants with varying experiences are represented. This will allow us to identify common patterns and variations in participants' experiences. The sample size will be sufficient for theoretical saturation [77, 78], and will allow for identification of common patterns and/

or variations across participant experiences. In addition, all facilitators will be interviewed to gain a deeper understanding of implementation patterns and processes, including their views on sustainability. Interviews will be conducted in English or Luganda. Questions will be translated/back translated by two team members fluent in both languages. Interviews will last about 1 h and will be audio taped.

## Quantitative measures and analyses

### Feasibility

We will monitor recruitment rates and staff level of effort, number of screenings conducted, as well as proportion eligible and agreed to enroll. Enrollment of 70% or higher will be considered feasible [79]. We will record the number of rescheduled, cancelled and missed G-CBT and MFG sessions, as well as assessments to inform estimation of staffing needs and retention protocols for a subsequent trial. We will also monitor attrition at each data collection point.

### Acceptability

We will adapt satisfaction surveys, e.g., the Client Satisfaction Questionnaire (CSQ-8), to assess acceptability [80]. Some of the items include: “How satisfied were you with the program?”, “How helpful was the program in addressing HIV-associated stigma?” and “How likely are you to recommend this program to other families with CLWH?” Given the modest sample size, quantitative analyses of intervention data will be largely descriptive and concentrate on tabulating and summarizing satisfaction outcomes.

### Preliminary impact

We expect that: H1: Following the intervention, relative to the control arm, participants in both treatment arms (G-CBT and MFG) will have: H1a: lower mean count of children reporting HIV-associated stigma; H1b: lower mean levels of reported HIV-associated stigma and improved child psychosocial functioning; H1c: lower mean levels of reported child mental health challenges; and H1d: higher mean levels of treatment adherence. H2: Following intervention, relative to the G-CBT intervention arm, participants in the MFG intervention arm will have: H2a: lower mean count of children reporting HIV-associated stigma; H2b: lower mean levels of reported HIV-associated stigma and improved child psychosocial functioning; H2c: lower mean levels of reported child mental health challenges; and H2d: higher mean levels of treatment adherence. Children will be the unit of analysis for these primary preliminary analyses.

Linear mixed models (LMMs) will be used to evaluate the proposed hypotheses. In addition, we will plot means

by group over time to describe overall patterns of change. We will fit three-level LMMs including random intercepts for clinic membership and random intercepts and slopes for subjects. Due to the modest sample size, significance testing will be de-emphasized. Similarly, although the modest sample size precludes formal investigation of moderation, we will apply the same LMM approach described above to compare children across study arms over time on the moderators listed in Table 2. These moderation analyses will be secondary exploratory analyses. Additional exploratory analyses will study caregivers and children jointly as the unit of analysis via dyadic analysis methods such as actor-partner and means-and-deviation models to quantify caregiver vs. child stigma effects and between- vs. within-dyad effects on mental health outcomes [81, 82].

### Sample size and power calculation

Although the study purpose is to determine feasibility and acceptability, rather than conduct formal hypothesis tests, we conducted several power analyses using NCSS PASS to supply additional information. Assuming  $\alpha = .05$  and power = .80, 81 participants would be retained at the final time point following 10% estimated attrition, and a clinic-based conservative unconditional ICC of 9.3% based on our previous Suubi study implemented among children orphaned by HIV/AIDS in the study region [83]. Therefore, our sample size of 90 dyads provides adequate statistical power to test our primary aim of feasibility. Additionally, for the target enrollment proportion of 70% to assess feasibility, the width to the limit of the confidence interval is 27.9% (standardized distance .32). For continuous standard normal variables to assess acceptability, the distance from the mean to the confidence limit is .30. These distances to confidence limits are between small (.20) and medium (.50) effect sizes. For preliminary efficacy exploratory analyses with two time points and paired comparisons of two out of the three groups at 81/3 = 27 participants per group ( $N = 54$  per comparison), minimum detectable standardized mean differences for continuous outcomes (stigma, mental health, and social support measures as described in Table 2) ranged from .79 to .97 for within-subject correlations ( $r$ ) ranging from .20 to .80. Overall, our study is powered to detect small-medium distances to confidence limits for descriptive statistics and large longitudinal analysis effects.

### Qualitative data analysis

All interviews will be audio-taped, transcribed, and uploaded to QSR NVivo12 analytic software. Transcripts will be reviewed by the research team to develop a broad understanding of the content and identify topics of discussion and observation. Analytic induction techniques



[84] will be used for coding. For initial coding, 10 transcripts will be randomly selected and read multiple times and independently coded by the team using a priori (from the interview guide) or emergent themes (open coding) [85]. Broader themes will be broken down into smaller, more specific units until no further subcategory is necessary. Analytic memos will be written to further develop categories, themes, and subthemes, and to integrate the ideas that emerge from the data [85, 86]. The codes and the inclusion/exclusion criteria for assigning a specific code [87] will be discussed as a team to create the final codebook in NVivo. Each transcript will then be coded independently by two team members using the codebook to establish inter-coder reliability. A level of agreement between 66 and 97% indicates good reliability [88]. Disagreements will be resolved through discussions in team meetings. The secondary analysis will compare themes within (including children versus caregivers) and across the two treatment groups to identify patterns, differences, and relationships among findings. Facilitators' data will be analyzed using the same procedures and will be compared and contrasted to participant data. To further ensure rigor, member checking to explore the opinions, beliefs and attitudes of participants, data audit trail, and analytic memos will be used [78, 89].

#### Data integration

Although qualitative and quantitative data analyses will be done separately, findings will be integrated at the interpretation and discussion stages. Conclusions and inferences will be synthesized for a more contextualized and thorough understanding of change mechanisms and the preliminary impact of each intervention arm. Specifically, qualitative and quantitative data will serve two purposes: (1) complementarity and (2) expansion [90, 91]. Qualitative findings will be connected to quantitative findings where the former will provide explanations and context for findings produced by the latter. Moreover, qualitative findings will complement our understanding of attendance and participant satisfaction for each treatment arm.

#### Data safety and monitoring

A Data and Safety Monitoring Board (DSMB) will not be needed for this study because of the following: (1) *Suubi4Stigma* is a pilot intervention study with children and their caregivers in a single site; (2) We are using two evidence-based interventions, with no known adverse effect on their recipients; and (3) the interventions are psychosocial in nature and poses no more than minimal risk to participants. We believe that the multiple principal investigators (MPIs) (Nabunya and Ssewamala), In-country-PI (Mugisha), and Institutional Review Boards (IRBs)—both

in Uganda and at Washington University in St. Louis, will be sufficient to monitor the trial. The MPIs have put in place a detailed Data and Safety Monitoring Plan (DSMP, details below) to ensure data safety.

To protect the integrity of participants' data, the following procedures will be followed: first, all data collected from the study participants will be used only for the purpose of research. All data will be kept confidential. We will not share any information or answers we get from participants with their families, classmates, friends, or teachers. In the same way, we will not share any information or answers we get from the primary caregivers with their children, other relatives, friends, teachers, or public officials. Second, all participants (adolescents and caregivers) in the study will be assigned a random code number by the in-country study coordinator under the guidance of the MPIs. This code number is used on all information collected from participants, including questionnaires. We will maintain lists of participants with links between identifying information and code numbers to facilitate participant follow-up. Only the MPIs and in-country study coordinator will have access to these lists, which are kept in locked files. Other study personnel have access on an as needed basis to individual participants' names and code numbers in order to adequately perform their duties, for example, interviewers must label the questionnaires with the correct code number of the participant whom they are interviewing.

All study personnel must complete certain levels of training before they are granted access to this identifying information. They must complete the Human Subjects Training. Personnel also sign confidentiality statements that specify that if the participants' confidentiality is breached unintentionally that personnel will follow the procedures for reporting this breach to the MPIs. The confidentiality statements also state that unintentional or deliberate violations of participants' confidentiality may result in demotion or termination depending upon the severity of the event. The project personnel also participate in training with the MPIs, the in-country study coordinator and the in-country PI regarding data safety, confidentiality of participants, limits of confidentiality, and proper administration of the study protocol.

All hard copies of data will be stored in locked cabinets to which only the MPIs and the in-country study coordinator will have access. After completion of an interview with a study participant, data with code numbers is placed in a separate locked file cabinet while waiting for entry. Once data is entered into computer files and password protected, only the MPIs, the in-country study coordinator and data entry assistants will have access to these files. All requests, current and future, to use the data are reviewed by the MPIs. Any data files that are

provided to other individuals are stripped of identifiers and contain only code numbers so that data across multiple assessment waves can be matched.

Within the informed consent/assent, participants are notified of the above procedures. Participants are also informed of the limits of confidentiality. Specifically, participants are warned that Ugandan law mandates reporting of abuse and/or neglect of children, and that threat of harm to self or others requires intervention by clinical staff. Participants will be informed of these limits during the consent process and at the beginning of the interview process. To make sure that interviewers have accurate knowledge of what does and does not constitute reportable child abuse and/or neglect, interviewers will receive training on the Ugandan laws regarding child abuse and/or neglect. Interviewers who suspect child abuse and/or neglect will be instructed to contact the in-country study coordinator, the in-country PI, and/or MPIs, rather than contact the Ugandan local authorities themselves. Prior to making a decision of whether to make a report of child abuse and/or neglect, the case will be discussed among a group consisting of the interviewer, study coordinator, and the MPIs. It is important to note that all three MPIs are social workers by training and are knowledgeable about what types of child abuse and/or neglect rise to the level required for a report to be made. These procedures are intended to protect the safety of children, and at the same time reduce the risk that erroneous reports are made. If we determine that a report must be made, we will inform the caregiver of our intention to report, and the reasons why a report must be made, unless we think that doing so would pose an immediate risk to the youth. In case further counseling is needed, the project staff will make the referrals.

#### **Monitoring and responding to adverse events**

All study personnel based in Uganda will be trained in identifying indicators of conditions that may jeopardize the welfare of participants and the limits of confidentiality. This training, conducted by the MPIs, includes reviewing possible scenarios and knowledge of key questions used to assess risk. Interview staff are trained to err on the side of caution and told to contact the study coordinator, who will always be available, by telephone, in the event of the need to break confidentiality due to mandatory reporting or ethical concerns. Under the guidance of in-country PI, research staff are trained either to contact the police to ensure safety of participants, or if appropriate, to have emergency personnel take the youth or caregiver to the nearest hospital.

Reporting of adverse events will occur according to a project protocol. For this study, safety and monitoring will be overseen by the in-country PI (Dr. Mugisha), and

the in-country Co-I (Dr. Mwebembezi)—both of whom will be stationed in Uganda during the study, and the other two MPIs (Drs. Nabunya and Ssewamala, based in the USA). In the case of an adverse event, staff will inform the in-country PI (Dr. Mugisha) and Co-I, Dr. Mwebembezi immediately and then the MPIs Drs. Nabunya and Ssewamala within 24 h of the presence of a possible unanticipated adverse event. Any presence of a possible unanticipated adverse event will be immediately reported and brought to the attention of the Washington University Institutional Review Board (along with the Ethics Committee at Uganda Virus Research Institute, and Uganda National Council of Science and Technology). The IRBs will determine whether it is appropriate to stop the study protocol temporarily or will provide suggestions and/or modifications to the study procedures. Possible modifications may include adding new risks to the consent form and re-consenting all study participants.

Preliminary outcomes data will be examined quarterly by the MPIs, in-country PI, and the co-investigators. If preliminary outcome data indicates harmful impact of the program to the study participants (the children and/or their caregivers), Washington University IRB committee, as well as the Ethic Committee at Uganda Virus Research Institute and Uganda National Council of Science and Technology IRB will be notified and it is possible that the study will be discontinued immediately. However, we do not anticipate any negative effects of

#### **Data management and quality assurance**

To collect data, Qualtrics, an electronic data capture system will be used. All tablets/iPads containing Qualtrics will be password protected. However, the option of completing a paper interview will remain available in case of technical difficulties. The interviewer will submit the tablet/iPad to their supervisor who will store the tablet/iPad in a locked file box until return from the field. Responses from Qualtrics will be downloaded daily by the data manager, onto a state-of-the-art secured dedicated server. All completed surveys will be printed out upon interviewers' return from the field. These hard copies will be stored as back up of the data in a double-locked environment. Trained, supervised research assistants will enter the data onto secure, encrypted computers. The entered data will then be double-checked for completeness and accuracy. To address concerns of breach of confidentiality, all study documents and data will be maintained in password-protected computer files. Confidentiality of study documents will be maintained by assigning unique study IDs and using these rather than participant names on all study related materials. Paper copies of documents will be maintained in locked file cabinets in respective countries. Participant consent forms and ID logs will be

kept in a separate locked cabinet. Tracking information also will be kept on a password-protected computer. The master list will only be used to coordinate data collection, and all staff will be required to receive training on both human subjects' protections, as well as maintaining the confidentiality of participants. The MPIs and the in-country study coordinator will oversee the quality assurance monitoring.

#### **Missing data**

Research assistants and study coordinator will be trained to check assessments for missing items before leaving the participant. If any missing items are identified, participants will be asked to answer those items—unless they chose not to answer those particular items. The study coordinator will also review assessments within 5 working days of completion. If missing items are identified in the baseline assessment (time point 1), research assistants will follow-up with the participants before the first session of the intervention. If missing items are identified at follow-up time points (3 and 6 months post-intervention initiation), research assistants will follow-up with the participants within 7 days of the original date of the assessment.

#### **Plan for record keeping**

The study will maintain records of adverse events, any referrals for counseling, as well as copies of the consent and assent forms. All records will be maintained in a locked filing cabinet at the Uganda-project office accessible only by the research team. The MPIs will be responsible for data security and record keeping. The data sets that will be used for analysis will not contain any identifying information—specifically, names and addresses of the participants.

#### **Plan for disposition of identifiers at the end of the study**

Identifiers for the participants will be disposed of following local IRB guidelines. To protect the participants' confidentiality, identifiers only will be accessible by the MPIs and Co-Is (Uganda- and US-based) and the study coordinator and will be kept separated from others with the participants' responses.

## **Discussion**

This study seeks to address the urgent need for innovative, theoretically, and empirically informed interventions to reduce HIV/AIDS-associated stigma and its negative impact on adolescent health and psychosocial well-being. More specifically, the study will examine two evidence-based interventions used in mental health settings, schools, and communities: G-CBT focused on cognitive restructuring and strengthening

coping skills at the individual level; and MFG intervention that strengthens family relationships to address stigma among CLWH and their families. Existing interventions focus on reducing fear of HIV infection among non-infected individuals as they interact with PLWH. This study offers an opportunity to develop a culturally acceptable and effective family-level intervention to address HIV/AIDS-associated stigma and its impact on CLWH's wellbeing in SSA.

This study innovates in the following ways: (1) HIV stigma-reduction interventions targeting CLWH in SSA are almost non-existent. This study will generate data driven knowledge to address HIV-associated stigma among CLWH and within their families. (2) The study will apply and compare two evidence-based, theoretically guided interventions, G-CBT vs MFG, to address HIV-associated stigma. (3) The MFG approach is culturally consistent with SSA's collective approach of families raising children "together," which strengthens its appeal to communities and its likelihood of success in addressing both individual-and family-level stigma. (4) Delivery of G-CBT, which will be facilitated by trained para-counselors, is an approach that has not been tested in this context and with this specific population. In Uganda, para-counselors are trained to assist with the psychological needs of individuals, including those related to HIV/AIDS and mental health. (5) Partnering with local institutions, including health clinics and community organizations, grounds the project with a practical understanding of the needs of CLWH in Masaka, a region hardest hit by the HIV/AIDS epidemic.

Study findings will be disseminated through local, national, and global meetings/workshops/conferences and peer-reviewed publications. The research team will also hold meetings to share study results with community stakeholders in the study region. This study will lay the foundation for a larger intervention study investigating how HIV/AIDS-associated stigma can be reduced to foster healthy child development, especially for CLWH as they transition to adolescence. The long-term goal is to develop culturally appropriate, feasible, acceptable, and effective interventions that address HIV/AIDS-associated stigma and its impact on CLWH's wellbeing and their families in SSA.

#### **Abbreviations**

ART: Antiretroviral therapy; CLWH: Children living with HIV; CSQ: Client Satisfaction Questionnaire; DSMB: Data and Safety Monitoring Board; DSMP: Data Safety and Monitoring Plan; G-CBT: Group Cognitive Behavioral Therapy; HIV: Human Immunodeficiency Viruses; ICHAD: International Center for Child Health and Development; IRB: Institutional Review Board; LMM: Linear mixed models; MFG: Multiple family groups; MPI: Multiple principal investigator; PI: Principal investigator; PLWH: People living with HIV; PTSD: Post-traumatic stress disorders; RA: Research assistant; RCT: Randomized controlled trial; RTY: Reach the Youth; SSA: Sub-Saharan Africa.

### Acknowledgements

We are grateful to our staff and volunteers at the International Center for Child Health and Development (ICHAD) field offices in Masaka, Uganda, and Reach the Youth (RTY) Uganda, for their respective contributions to study implementation. In addition, we are grateful to the health care clinics that have agreed to work with children and their families, as well as facilitators and counselors who will deliver the interventions. Our special thanks go to all children and their families who have agreed to participate in the Suubi4Stigma study.

### Authors' contributions

PN and FS are the Multiple Principal Investigators (MPIs) responsible for the conceptualization, design, implementation and administration of the study. JM is the in-country PI responsible for overseeing field activities. OSB contributed to the conceptualization and design of the study, and she is a Co-Investigator on the study. TN is the statistical consultant. LTMM, TJ-F and MMM contributed to the initial review and edits to the study proposal. All authors have reviewed and commented on drafts and have approved the final manuscript prior to submission.

### Funding

The study outlined in this protocol is supported by the National Institute of Mental Health (NIMH) under Award Number R21MH121141 (MPIs: Proscovia Nabunya, PhD; Fred M. Ssewamala, PhD). NIMH has no role in the study design, data collection, analysis, interpretation of findings and manuscript preparation. The content is solely the responsibility of the authors and does not necessarily represent the official views of NIMH or the National Institutes of Health.

### Availability of data and materials

N/A

### Declarations

#### Ethics approval and consent to participate

Voluntary written informed assent and consent will be obtained from children and their caregivers, respectively. In addition, all study procedures were approved by the Washington University in St. Louis Review Board (#202009185) and by in-country local IRBs in Uganda: Uganda Virus Research Institute (UVRI – GC/127/20/10/792), and the Uganda National Council for Science and Technology (UNCST – #SS632ES).

#### Consent for publication

N/A

#### Competing interests

The authors declare that they have no competing interests.

#### Author details

<sup>1</sup>International Center for Child Health and Development (ICHAD), St. Louis, USA. <sup>2</sup>Brown School, Washington University in St. Louis, Campus Box 1196, One Brookings Drive, St. Louis, MO 63130, USA. <sup>3</sup>The Moving Well Project International, Inc., Silver Spring, USA. <sup>4</sup>School of Social Work, Columbia University, New York City, USA. <sup>5</sup>College of Health Sciences, Makerere University, Kampala, Uganda. <sup>6</sup>Division of Prevention Science, School of Medicine, University of California San Francisco, San Francisco, USA.

Received: 6 December 2020 Accepted: 14 April 2022

Published online: 29 April 2022

### References

- UNAIDS. Global HIV and AIDS Statistics–2020 fact sheet. Available from: [https://www.unaids.org/sites/default/files/media\\_asset/UNAIDS\\_Factsheet\\_en.pdf](https://www.unaids.org/sites/default/files/media_asset/UNAIDS_Factsheet_en.pdf). Accessed 23 Aug 2020.
- UNAIDS. UNAIDS data 2020. Available from: [https://www.unaids.org/sites/default/files/media\\_asset/2020\\_aids-data-book\\_en.pdf](https://www.unaids.org/sites/default/files/media_asset/2020_aids-data-book_en.pdf). Accessed 23 Aug 2020.
- Uganda AIDS Commission. 12<sup>th</sup> annual joint AIDS review. Final report July 2018–June 2019. Available from: <https://uac.go.ug/sites/default/files/Reports/Final%20%2012th%20Annual%20Joint%20AIDS%20Review%20Report%20.pdf>. Accessed 23 Aug 2020.
- UPHIA. Uganda population-based HIV impact assessment 2016–2017. Available from: <https://www.afro.who.int/sites/default/files/2017-08/UPHIA%20Uganda%20factsheet.pdf>. Accessed 23 Aug 2020.
- Domek GJ. Social consequences of antiretroviral therapy: preparing for the unexpected futures of HIV-positive children. *Lancet*. 2006;367(9519):1367–9.
- Gortmaker SL, Hughes M, Cervia J, Brady M, Johnson GM, Seage GR III, et al. Effect of combination therapy including protease inhibitors on mortality among children and adolescents infected with HIV-1. *N Engl J Med*. 2001;345(21):1522–8.
- Sengupta S, Banks B, Jonas D, Miles MS, Smith GC. HIV interventions to reduce HIV/AIDS stigma: a systematic review. *AIDS Behav*. 2011;15(6):1075–87.
- Stangl AL, Lloyd JK, Brady LM, Holland CE, Baral S. A systematic review of interventions to reduce HIV-related stigma and discrimination from 2002 to 2013: how far have we come? *J Int AIDS Soc*. 2013;16:18734.
- Bennett DS, Traub K, Mace L, Juarascio A, O'Hayer CV. Shame among people living with HIV: a literature review. *AIDS Care*. 2016;28(1):87–91.
- Katz IT, Ryu AE, Onuegbu AG, Psaros C, Weiser SD, Bangsberg DR, et al. Impact of HIV-related stigma on treatment adherence: systematic review and meta-synthesis. *J Int AIDS Soc*. 2013;16:18640.
- Lowther K, Selman L, Harding R, Higginson IJ. Experience of persistent psychological symptoms and perceived stigma among people with HIV on antiretroviral therapy (ART): a systematic review. *Int J Nurs Stud*. 2014;51(8):1171–89.
- Abas M, Ali GC, Nakimuli-Mpungu E, Chibanda D. Depression in people living with HIV in sub-Saharan Africa: time to act. *Trop Med Int Health*. 2014;19(12):1392–6.
- Breuer E, Myer L, Struthers H, Joska JA. HIV/AIDS and mental health research in sub-Saharan Africa: a systematic review. *Afr J AIDS Res*. 2011;10(2):101–22.
- Fekete EM, Williams SL, Skinta MD. Internalised HIV-stigma, loneliness, depressive symptoms and sleep quality in people living with HIV. *Psychol Health*. 2018;33(3):398–415.
- Nachega JB, Morroni C, Zuniga JM, Sherer R, Beyrer C, Solomon S, et al. HIV-related stigma, isolation, discrimination, and serostatus disclosure: a global survey of 2035 HIV-infected adults. *J Int Assoc Phys AIDS Care*. 2012;11(3):172–8.
- Luisa Zúñiga M, Blanco E, Martínez P, Strathdee SA, Gifford AL. Perceptions of barriers and facilitators to participation in clinical trials in HIV-positive Latinas: a pilot study. *J Womens Health*. 2007;16(9):1322–30.
- Dickerson SS, Gruenewald TL, Kemeny ME. When the social self is threatened: shame, physiology, and health. *J Pers*. 2004;72(6):1191–216.
- Owe-Larsson MB, Säll L, Salamon E, Allgulander C. HIV infection and psychiatric illness. *Afr J Psychiatry*. 2009;12(2):115–28.
- Wolf HT, Halpern-Felsher BL, Bukusi EA, Agot KE, Cohen CR, Auerswald CL. "It is all about the fear of being discriminated [against]... the person suffering from HIV will not be accepted": a qualitative study exploring the reasons for loss to follow-up among HIV-positive youth in Kisumu, Kenya. *BMC Public Health*. 2014;14(1):1154.
- Karim QA, Meyer-Weitz A, Mboyi L, Carrara H, Mahlase G, Frohlich JA, et al. The influence of AIDS stigma and discrimination and social cohesion on HIV testing and willingness to disclose HIV in rural KwaZulu-Natal, South Africa. *Glob Public Health*. 2008;3(4):351–65.
- Bogart LM, Cowgill BO, Kennedy D, Ryan G, Murphy DA, Elijah J, et al. HIV-related stigma among people with HIV and their families: a qualitative analysis. *AIDS Behav*. 2008;12(2):244–54.
- Powell-Cope GM, Brown MA. Going public as an AIDS family caregiver. *Soc Sci Med*. 1992;34(5):571–80.
- Mwinitoo PP, Mill JE. Stigma associated with Ghanaian caregivers of AIDS patients. *West J Nurs Res*. 2006;28(4):369–82.
- DuBois DL, Burk-Braxton C, Swenson LP, Tevendale HD, Lockerd EM, Moran BL. Getting by with a little help from self and others: self-esteem and social support as resources during early adolescence. *Dev Psychol*. 2002;38(5):822.
- Du H, Li X, Chi P, Zhao J, Zhao G. Relational self-esteem, psychological well-being, and social support in children affected by HIV. *J Health Psychol*. 2015;20(12):1568–78.

26. Hamburg BA. Life skills training: preventive interventions for young adolescents. New York: Carnegie Council on Adolescent Development; 1990.
27. Krizek BL, Mugisha J, Osafo J, Kinyanda E. Growing up HIV-positive in Uganda: "psychological immunodeficiency"? A qualitative study. *BMC Psychol*. 2017;5(1):1–0.
28. Rohner RP, Britner PA. Worldwide mental health correlates of parental acceptance-rejection: review of cross-cultural and intracultural evidence. *Cross-Cult Res*. 2002;36(1):16–47.
29. Day RD, Padilla-Walker LM. Mother and father connectedness and involvement during early adolescence. *J Fam Psychol*. 2009;23(6):900.
30. Steinberg L. We know some things: parent-adolescent relationships in retrospect and prospect. *J Res Adolesc*. 2001;11(1):1–9.
31. Hauser ST. Understanding resilient outcomes: adolescent lives across time and generations. *J Res Adolesc*. 1999;9(1):1–24.
32. McNeely C, Shew ML, Behring T, Sieving R, Miller BC, Blum RW. Mothers' influence on the timing of their first sex among 14- and 15-year-olds. *J Adolesc Health*. 2002;31(3):256–65.
33. Amerikaner M, Monks G, Wolfe P, Thomas S. Family interaction and individual psychological health. *J Couns Dev*. 1994;72(6):614–20.
34. Walker CE, Roberts MC. *Handbook of clinical child psychology*. Hoboken: Wiley; 2001.
35. Neema S, Atuyambe LM, Otolok-Tanga B, Twijukye C, Kambugu A, Thayer L, et al. Using a clinic-based creativity initiative to reduce HIV related stigma at the Infectious Diseases Institute, Mulago National Referral Hospital, Uganda. *Afr Health Sci*. 2012;12(2):231–9.
36. Uys L, Chirwa M, Kohi T, Greeff M, Naidoo J, Makoae L, et al. Evaluation of a health setting-based stigma intervention in five African countries. *AIDS Patient Care STDs*. 2009;23(12):1059–66.
37. Earnshaw VA, Chaudoir SR. From conceptualizing to measuring HIV stigma: a review of HIV stigma mechanism measures. *AIDS Behav*. 2009;13(6):1160.
38. Earnshaw VA, Smith LR, Chaudoir SR, Amico KR, Copenhaver MM. HIV stigma mechanisms and well-being among PLWH: a test of the HIV stigma framework. *AIDS Behav*. 2013;17(5):1785–95.
39. McKay MM, Gonzales JJ, Stone S, Ryland D, Kohner K. Multiple family therapy groups: a responsive intervention model for inner city families. *Soc Work Groups*. 1995;18(4):41–56.
40. Beck AT. Cognitive therapy: past, present, and future. *J Consult Clin Psychol*. 1993;61(2):194.
41. Ssewamala FM, Byansi W, Bahar OS, Nabunya P, Neilands TB, Mellins C, et al. Suubi+ Adherence study protocol: a family economic empowerment intervention addressing HIV treatment adherence for perinatally infected adolescents. *Contemp Clin Trials Commun*. 2019;16:100463.
42. Ssewamala FM, Bermudez LG, Neilands TB, Mellins CA, McKay MM, Garfinkel I, et al. Suubi4Her: a study protocol to examine the impact and cost associated with a combination intervention to prevent HIV risk behavior and improve mental health functioning among adolescent girls in Uganda. *BMC Public Health*. 2018;18(1):693.
43. Sensoy Bahar O, Byansi W, Kivumbi A, Namatovu P, Kiyingi J, Ssewamala FM, et al. From "4Rs and 2Ss" to "Amaka Amasanyufu" (happy families): adapting a US-based evidence-based intervention to the Uganda context. *Fam Process*. 2020;59(4):1928–45.
44. Ssewamala FM, Dvalishvili D, Mellins CA, Geng EH, Makumbi F, Neilands TB, et al. The long-term effects of a family based economic empowerment intervention (Suubi+ Adherence) on suppression of HIV viral loads among adolescents living with HIV in southern Uganda: findings from 5-year cluster randomized trial. *PLoS One*. 2020;15(2):e0228370.
45. The Republic of Uganda. National HIV and AIDS strategic plan 2015/2016–2019/2020. Available from: [https://www.ilo.org/wcmsp5/groups/public/-ed\\_protect/-protrav/-ilo\\_aids/documents/policy/wcms\\_455260.pdf](https://www.ilo.org/wcmsp5/groups/public/-ed_protect/-protrav/-ilo_aids/documents/policy/wcms_455260.pdf). Accessed 23 Aug 2020.
46. Clauss-Ehlers CS, editor. *Encyclopedia of cross-cultural school psychology*: Springer Science & Business Media; 2010.
47. Nardi B, Massei M, Arimatea E, Moltedo-Perfetti A. Effectiveness of group CBT in treating adolescents with depression symptoms: a critical review. *Int J Adolesc Med Health*. 2017;29(3):20150080.
48. Sexton TL, Alexander JF. Family-based empirically supported interventions. *Couns Psychol*. 2002;30(2):238–61.
49. Jones TL, Prinz RJ. Potential roles of parental self-efficacy in parent and child adjustment: a review. *Clin Psychol Rev*. 2005;25(3):341–63.
50. Ssewamala FM, Bahar OS, McKay MM, Hoagwood K, Huang KY, Pringle B. Strengthening mental health and research training in Sub-Saharan Africa (SMART Africa): Uganda study protocol. *Trials*. 2018;19(1):423.
51. Nseroko ND. The evolution of mental health understanding and practice in Uganda. *Int J Emerg Ment Health Hum Resilience*. 2017;19(1):354–60.
52. Hohenshil TH, Amundson NE, Niles SG, editors. *Counseling around the world: an international handbook*: Wiley; 2015.
53. Ssewamala FM, Mukasa MN, Kivumbi A, Mwebembezi A, Byansi W, Nabunya P, et al. Suubi+Adherence Cartoon Intervention: utilizing cartoons to address challenges and share experiences from HIV positive youth in Uganda. New York: Masaka Diocese, Reach the Youth Uganda and Columbia University, International Center for Child Health and Asset Development; 2020.
54. Verdelli H, Clougherty K, Bolton P, Speelman L, Lincoln N, Bass J, et al. Adapting group interpersonal psychotherapy for a developing country: experience in rural Uganda. *World Psychiatry*. 2003;2(2):114.
55. Pappas RK, Sidle JE, Martino S, Baliddawa JB, Songole R, Omolo OE, et al. Systematic cultural adaptation of cognitive-behavioral therapy to reduce alcohol use among HIV-infected outpatients in western Kenya. *AIDS Behav*. 2010;14(3):669–78.
56. Bhana A, Mellins CA, Petersen I, Alicea S, Myeza N, Holst H, et al. The VUKA family program: piloting a family-based psychosocial intervention to promote health and mental health among HIV infected early adolescents in South Africa. *AIDS Care*. 2014;26(1):1–1.
57. Kovacs M. *Children's depression inventory: manual*. North Tonawanda: Multi-Health Systems; 1992.
58. Fitts WH, Warren WL. *Tennessee self-concept scale: TSCS-2*. Los Angeles: Western Psychological Services; 1996.
59. Beck AT, Weissman A, Lester D, Trexler L. The measurement of pessimism: the hopelessness scale. *J Consult Clin Psychol*. 1974;42(6):861.
60. Frederick C, Pynoos RS, Nader KO. *Childhood post-traumatic stress reaction index (CPTSS-RI)*. Los Angeles: Two Suns Measures; 1992.
61. Moos RH, Moos BS. *Family environment scale manual: development, applications, research*. 3rd ed. Palo Alto: Consulting Psychologist Press; 1994.
62. Skinner HA, Steinhauer PD, Santa-Barbara J. The family assessment measure. *Can J Commun Ment Health*. 1983;2(2):91–103.
63. Vaux A, Riedel S, Stewart D. Modes of social support: the social support behaviors (SS-B) scale. *Am. J. Community Psychol*. 1987;15(2):209–32.
64. Bukowski WM, Hoza B, Boivin M. Measuring friendship quality during pre- and early adolescence: the development and psychometric properties of the Friendship Qualities Scale. *J Soc Pers Relat*. 1994;11(3):471–84.
65. Michalopoulos LM, Meinhardt M, Barton SM, Kuhn J, Mukasa MN, Namuwonge F, et al. Adaptation and validation of the shame questionnaire among Ugandan youth living with HIV. *Child Indic Res*. 2019;12(3):1023–42.
66. Boyes ME, Mason SJ, Cluver LD. Validation of a brief stigma-by-association scale for use with HIV/AIDS-affected youth in South Africa. *AIDS Care*. 2013;25(2):215–22.
67. Berger BE, Ferrans CE, Lashley FR. Measuring stigma in people with HIV: psychometric assessment of the HIV stigma scale. *Res Nurs Health*. 2001;24(6):518–29.
68. Russell DW. UCLA Loneliness Scale (Version 3): reliability, validity, and factor structure. *J Pers Assess*. 1996;66(1):20–40.
69. McKay MM, Alicea S, Elwyn L, McClain ZR, Parker G, Small LA, et al. The development and implementation of theory-driven programs capable of addressing poverty-impacted children's health, mental health, and prevention needs: CHAMP and CHAMP+, evidence-informed, family-based interventions to address HIV risk and care. *J Clin Child Adolesc Psychol*. 2014;43(3):428–41.
70. Auslander WF, Slonim-Nevo V, Ozawa M, Shepard S, Gehlert S. Comparison of skills training, discussion, and control groups to prevent AIDS among abused and delinquent youth. In: VIII International Conference on AIDS/III STD World Congress, Amsterdam, The Netherlands; 1992. p. 19–24.
71. Slonim-Nevo V, Auslander WF, Ozawa MN. Educational options and AIDS-related behaviors among troubled adolescents. *J Pediatr Psychol*. 1995;20(1):41–60.
72. Wilson IB, Lee Y, Michaud J, Fowler FJ, Rogers WH. Validation of a new three-item self-report measure for medication adherence. *AIDS Behav*. 2016;20(11):2700–8.

73. Genberg BL, Kawichai S, Chingono A, Sendah M, Chariyalertsak S, Konda KA, et al. Assessing HIV/AIDS stigma and discrimination in developing countries. *AIDS Behav.* 2008;12(5):772–80.
74. Derogatis LR. BSI brief symptom inventory. Administration, scoring, and procedures manual; 1993.
75. Abidin RR, Parenting Stress Index-Manual Charlottesville, Pediatric Psychology Press, (ERIC Document Reproduction Service No. ED 238 896), Beckman P. Comparison of mothers' and fathers' perceptions of the effect of young children with and without disabilities. *Am J Ment Retard.* 1991;95:585–95.
76. Tesch R. Qualitative research: analysis types and software tools. Bedford: Routledge.
77. Guest G, Bunce A, Johnson L. How many interviews are enough? An experiment with data saturation and variability. *Field Methods.* 2006;18(1):59–82.
78. Padgett DK. Qualitative methods in social work research: Sage publications; 2016.
79. Crepaz N, Lyles CM, Wolitski RJ, Passin WF, Rama SM, Herbst JH, et al. Do prevention interventions reduce HIV risk behaviours among people living with HIV? A meta-analytic review of controlled trials. *Aids.* 2006;20(2):143–57.
80. Client Satisfaction Questionnaire (CSWQ-3). Available at <http://www.csqscales.com/>. Accessed 23 Aug 2020.
81. Kenny DA, Kashy DA, Cook WL, Simpson JA. Dyadic data analysis (*Methodology in the social sciences*); 2020. p. 335–7.
82. Neuhaus JM, Kalbfleisch JD. Between-and within-cluster covariate effects in the analysis of clustered data. *Biometrics.* 1998;54(2):638–45.
83. Ssewamala FM, Neilands TB, Waldfogel J, Ismayilova L. The impact of a comprehensive microfinance intervention on depression levels of AIDS-orphaned children in Uganda. *J Adolesc Health.* 2012;50(4):346–52.
84. Lincoln YS, Guba EG. *Naturalistic inquiry*. Newbury Park: Sage Publications; 1985.
85. Corbin J, Strauss A. *Basics of qualitative research: techniques and procedures for developing grounded theory*: Sage publications; 2014.
86. Charmaz K. Grounded theory: objectivist and constructivist methods. In: Denzin NK, Lincoln YS, editors. *Strategies for qualitative inquiry*; 2003. p. 249–91.
87. Miles MB, Huberman AM. *Qualitative data analysis: an expanded sourcebook*: Sage; 1994.
88. Boyatzis RE. *Transforming qualitative information: thematic analysis and code development*: Sage; 1998.
89. Birt L, Scott S, Cavers D, Campbell C, Walter F. Member checking: a tool to enhance trustworthiness or merely a nod to validation? *Qual Health Res.* 2016;26(13):1802–11.
90. Greene JC, Caracelli VJ, Graham WF. Toward a conceptual framework for mixed-method evaluation designs. *Educ Eval Policy Anal.* 1989;11(3):255–74.
91. Greene JC, Caracelli VJ. *Advances in mixed-method evaluation: the challenges and benefits of integrating diverse paradigms*. San Francisco: Jossey-Bass; 2020.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

