

## Original Article

# Elucidating the role of ApxI in hemolysis and cellular damage by using a novel *apxIA* mutant of *Actinobacillus pleuropneumoniae* serotype 10

Nai-Yun Chang<sup>1</sup>, Zeng-Weng Chen<sup>3</sup>, Ter-Hsin Chen<sup>1,2</sup>, Jiunn-Wang Liao<sup>1</sup>, Cheng-Chung Lin<sup>1</sup>,  
Maw-Sheng Chien<sup>1</sup>, Wei-Cheng Lee<sup>1</sup>, Jiunn-Horng Lin<sup>3</sup>, Shih-Ling Hsuan<sup>1,\*</sup>

Graduate Institutes of <sup>1</sup>Veterinary Pathobiology, and <sup>2</sup>Microbiology and Public Health, College of Veterinary Medicine, National Chung Hsing University, Taichung 402, Taiwan, R.O.C.

<sup>3</sup>Division of Animal Medicine, Animal Technology Institute Taiwan, Miaoli, 350, Taiwan, R.O.C.

Exotoxins produced by *Actinobacillus (A.) pleuropneumoniae* (Apx) play major roles in the pathogenesis of pleuropneumonia in swine. This study investigated the role of ApxI in hemolysis and cellular damage using a novel *apxIA* mutant, ApxIA336, which was developed from the parental strain *A. pleuropneumoniae* serotype 10 that produces only ApxI *in vitro*. The genotype of ApxIA336 was confirmed by PCR, Southern blotting, and gene sequencing. Exotoxin preparation derived from ApxIA336 was analyzed for its bioactivity towards porcine erythrocytes and alveolar macrophages. Analysis results indicated that ApxIA336 contained a kanamycin-resistant cassette inserted immediately after 1005 bp of the *apxIA* gene. Phenotype analysis of ApxIA336 revealed no difference in the growth rate as compared to the parental strain. Meanwhile, ApxI production was abolished in the bacterial culture supernatant, *i.e.* exotoxin preparation. The inability of ApxIA336 to produce ApxI corresponded to the loss of hemolytic and cytotoxic bioactivity in exotoxin preparation, as demonstrated by hemolysis, lactate dehydrogenase release, mitochondrial activity, and apoptosis assays. Additionally, the virulence of ApxIA336 appeared to be attenuated by 15-fold in BALB/c mice. Collectively, ApxI, but not other components in the exotoxin preparation of *A. pleuropneumoniae* serotype 10, was responsible for the hemolytic and cytotoxic effects on porcine erythrocytes and alveolar macrophages.

**Keywords:** *Actinobacillus pleuropneumoniae*, ApxI, attenuation, cytotoxicity, mutant

## Introduction

*Actinobacillus (A.) pleuropneumoniae*, a Gram-negative bacterium, is the causative agent of necrotizing and

hemorrhagic pleuropneumonia in pigs [8,13]. Fifteen serotypes of *A. pleuropneumoniae* have been identified based on antigenic differences of the capsular polysaccharide [7]. *A. pleuropneumoniae* exotoxin (Apx) is one of the important virulence factors contributing to the pathogenesis of porcine pleuropneumonia [13]. Given their structural feature, Apx toxins belong to the family of repeats in toxin (RTX) [24]. Other members of the RTX family include hemolysin produced by *Escherichia (E.) coli* and leukotoxin (LktA) from *Mannheimia haemolytica* [19,23]. ApxI to IV toxins cause pulmonary damage in pigs [1,15,22]. Among these toxins, ApxI exerts strong hemolytic and cytotoxic effects on a variety of cells including porcine neutrophils, lymphocytes, porcine alveolar macrophages (PAMs), and pulmonary endothelial cells [9,11,14,21,22]. Additionally, ApxI induces apoptosis and pro-inflammatory cytokine expression in PAMs through a mitogen-activated protein kinase pathway [2,3,25].

Our previous study investigated how a single species of exotoxin affects PAMs by using the crude cultural supernatant derived from *A. pleuropneumoniae* serotype 10 as the source of native ApxI. It was necessary to use either heat-inactivated or antibody-neutralized exotoxin or add lipopolysaccharide (LPS) inhibitor polymyxin B to demonstrate the specificity of apoptotic effect caused by ApxI [2,4,25]. The purpose of the present study was to construct an *apxIA* mutant using a strain of *A. pleuropneumoniae* serotype 10 bacterium [10] by inserting a kanamycin-resistant gene cassette into the *apxIA* gene via allelic exchange. The culture supernatant of this mutant designated ApxIA336 was evaluated for biological activity namely, hemolytic activity; cytotoxicity; apoptotic effects; and virulence to mice.

\*Corresponding author: Tel: +886-4-2284-0368; Fax: +886-4-2285-2186; E-mail: hsuan@nchu.edu.tw

## Materials and Methods

### Antibodies and chemicals


Restriction enzymes were purchased from New England Biolabs (USA). Anti-digoxigenin (DIG) Fab fragment conjugated with horseradish peroxidase (HRP), a DIG DNA labeling kit, and cytotoxic detection kit (LDH) were obtained from Roche (Switzerland). Hoechst 33258, nicotinamide adenine dinucleotide (NAD), poly-D-lysine (PDL), polymyxin B, and 2,3-bis (2-methoxy-4-nitro-5-sulfophenyl)-2H-tetrazolium-5-carboxanilide inner salt (XTT) were obtained from Sigma-Aldrich (USA). Bradford reagent and  $\gamma$ -globulin standard were purchased from Bio-Rad Laboratories (USA). A limulus amoebocyte lysate (LAL) assay kit was obtained from Cambrex Bio Science (USA). Medium, antibiotics and supplementary components for cell culture were purchased from Gibco (USA). Bacterial culture materials were purchased from Becton Dickinson (USA).

### Cell culture

Primary porcine alveolar macrophages (PAMs) were prepared from 4 to 6 weeks old healthy piglets as described elsewhere [25]. The piglets were cared for and euthanized based on a protocol approved by the Institutional Animal Care and Use Committee (IACUC) of National Chung Hsing University (Taiwan, R.O.C.). PAMs were obtained through lavage and stored in liquid nitrogen as previously described [4]. PAMs were thawed and cultured overnight in RPMI 1640 supplemented with 10% fetal bovine serum (FBS), 2 mM L-glutamine, 100 U/mL penicillin, and 100  $\mu$ g/mL streptomycin at 37°C in 5% CO<sub>2</sub> before the experiments.

### Construction of the *apxIA* insertion mutant

An *apxIA* mutant was constructed by introducing a kanamycin-resistant gene cassette into the *apxIA* gene of *A. pleuropneumoniae* serotype 10 (strain 13039, Denmark), a kind gift from the Animal Health Research Institute, Council of Agriculture, Republic of China, using a molecular cloning approach (Fig. 1). Briefly, *A. pleuropneumoniae* serotype 10 was cultured overnight in brain heart infusion (BHI) broth containing 10  $\mu$ g/mL NAD (NAD/BHI) at 37°C. Genomic DNA was extracted from the bacterial culture according to the manual provided by QIAamp DNA mini kit (QIAGEN, Germany). Full-length *apxIA* DNA was amplified by PCR using primers ApxI-F1 and ApxI-R1 (Table 1): initial denaturing of DNA at 94°C for 5 min, followed by 35 cycles of denature at 94°C for 45



**Fig. 1.** Schematic diagram of the *apxIA* mutant construction. A kanamycin-resistant cassette (*kan<sup>R</sup>*) was inserted into the *apxIA* gene. Primer locations (arrows) and restriction enzyme cleavage sites are indicated.

**Table 1.** Plasmids and primers used in this study

Plasmids	Characteristics	Source		
pCRZAp <sub>x</sub> IA	Kan <sup>R</sup> , Zeo <sup>R</sup> cloning vector; 3246 bp <i>apxIA</i> cloned into pCR-XL-TOPO	This work		
pCRZAp <sub>x</sub> IA	Zeo <sup>R</sup> cloning vector; modified pCRZAp <sub>x</sub> IA by deleting kanamycin-resistant gene	This work		
pCRZAp <sub>x</sub> IAK	Kan <sup>R</sup> , Zeo <sup>R</sup> cloning vector; 1252 bp kanamycin-resistant gene inserted into <i>apxIA</i> gene of pCRZAp <sub>x</sub> IA	This work		
Primers	Sequence (5' → 3')*	Amplified gene	Position	Accession number
ApxI-F1	TTACAACATGGCTAACTCTCAGCTC	<i>apxIA</i>	(-5)-17	X52899
ApxI-R1	TAGCAATCCTACACACCTAATATAT		(-36)-(-60)	
ApxI-F2	GCGGCAGGCTTATCCACAAC		876-896	
ApxI-R2	CGATAGATGAACCGGATGAAAGA		1859-1881	
ApxIV-F	GTGCCCATGGGCACAAAATTAAGTATGCAAGATGT	<i>apxIVA</i>	17-39	AF021919
ApxIV-R	AAGT <u>AAGCTT</u> ACTTTTTTAAGTACTTTTTTAACGGCG		1492-1513	
Kana-F	CAAGTCAGCGTAATGCTCTGCCAGTGTTAC	<i>kan<sup>R</sup></i>	(-50)-(-21)	X06404
Kana-R	TATGCCTCTTCCGACCATCAAGCATTTTTAT		553-582	

\*Restriction enzyme sites are underlined.

sec, annealing at 53°C for 45 sec, and extension at 72°C for 2 min, then a final extension at 72°C for 7 min. The amplified PCR product was then purified by QIAquick PCR purification kit (QIAGEN, Germany) and ligated to a pCR-XL-TOPO vector (Invitrogen, USA) according to the manufacturer's instructions. The construct was used to transform into One Shot TOP10 *E. coli* and cultured in Luria-Bertani (LB) broth supplemented with 25 µg/mL of zeocine. pCR-XL-TOPO plasmid containing *apxI* was referred to as pCRKZApXIA. Next, a suicide vector construct was created by removing the kanamycin-resistant gene of pCRKZApXIA via restriction enzyme digestion with *SphI* followed by self-ligation of the plasmid. The resulting plasmid DNA was called pCRZApXIA. The pCRZApXIA plasmid was digested with *SspI* to create a blunt end in the *apxIA* gene fragment and ligated to a kanamycin-resistant cassette gene from the pUC4K plasmid digested with *HincII*. The resulting suicide vector used for allelic exchange was named pCRZApXIAK.

Allelic exchange in *A. pleuropneumoniae* was performed by diluting the overnight culture of *A. pleuropneumoniae* serotype 10 at a ratio of 1 : 20 in LB broth supplemented with 10 µg/mL NAD and culturing at 37°C for 3 h with constant agitation. Ten mL of the bacterial culture were then centrifuged at 16,000 × g for 1 min at 4°C, and the resulting pellet was washed three times with and resuspended in 400 µL ice-cold electroporation buffer [27 mM sucrose, 15% glycerol, and 3 mM of potassium phosphate (pH 7.4)] [12]. Five to ten µg of pCRZApXIAK plasmid DNA were added to the bacterial suspension, transferred to a 2-mm electroporation cuvette (Thermo Fisher Scientific, USA), and pulsed at 2.5 kV, 800 Ω, and 25 µF (Bio-Rad Laboratories). Next, 1 mL of NAD/BHI broth was added and the cells were cultured at 37°C for 3 h. The transformants were further cultured overnight at 37°C on a NAD/BHI agar plate supplemented with 25 µg/mL kanamycin. Finally, bacterial colonies were analyzed by PCR with different primer pairs (Table 1). DNA amplified with primers ApxI-F1/R1 and kana-F/R was initially denatured at 94°C for 5 min, followed by 35 cycles of denature at 94°C for 45 sec, annealing at 53°C for 45 sec, extension at 72°C for 2 min, and a final extension at 72°C for 7 min. DNA amplified with primers ApxIV-F/R was initially denatured at 94°C for 5 min, followed by 35 cycles of denature at 94°C for 30 sec, annealing at 62°C for 30 sec, extension at 72°C for 3 min, and a final extension at 72°C for 10 min.

### Southern blot analysis

Genomic DNA of parental and mutant strains were extracted and digested with *PvuI*. Five µg of digested DNA were separated by electrophoresis in 0.7% agarose and transferred onto a nylon membrane (Amersham Biosciences, Sweden). The DNA was crosslinked to the

membrane by exposure to ultraviolet light for 2 min. ApxI-F2/R2 primers and DIG-labeled dNTP were then used to generate the *apxIA* probe by using *A. pleuropneumoniae* serotype 10 genomic DNA as a PCR template. Next, pUC4K plasmid DNA was amplified using Kana-F/R primers to produce a DIG-labeled kanamycin-resistant cassette probe. The nylon membranes were hybridized overnight with 25 ng/mL of DIG-labeled *apxIA* or kanamycin-resistant cassette probes at 42°C. The membranes were subsequently washed with buffer containing 100 mM maleic acid, 150 mM NaCl, and 3% Tween 20 (pH 7.5) and probed with an anti-DIG Fab fragment conjugated with horseradish peroxidase at room temperature for 1 h. Probe hybridization was detected using chemiluminescence reagent plus (PerkinElmer, USA).

### Sequencing analysis

PCR was performed with genomic DNA from mutant strain using the primer pair ApxI-F1/R1. The PCR product was sequenced by Tri-I Biotech (Taiwan, R.O.C.).

### Bacterial growth curve determination and measurement of kanamycin-resistant cassette insertion stability

A single colony of parental or mutant strain was transferred to 10 mL NAD/BHI broth and cultured at 37°C with constant agitation for 16 h. The culture was mixed with fresh NAD/BHI broth at a 1 : 100 ratio and incubated at 37°C for 7 h. At 1-h intervals, 100 µL of bacterial suspension was sampled, 10-fold serial-diluted and spread onto NAD/BHI agar plates. Following incubation at 37°C overnight, the colonies on the plates were counted, and the number of colony forming units (CFUs) per mL was calculated.

Stability of the kanamycin-resistant cassette in mutant strain was also evaluated by growing bacteria in NAD/BHI broth for 10 serial passages. Genomic DNA was extracted from the bacteria at each passage. PCR amplification of the kanamycin-resistant cassette was then performed.

### Exotoxin preparation, protein concentration determination, and measurement of lipopolysaccharide (LPS) content

*A. pleuropneumoniae* exotoxin was prepared from a cultural supernatant as previously described [4]. Protein and LPS concentrations in the exotoxin preparation were measured with a Bradford protein assay and limulus amoebocyte lysate (LAL) assay, respectively. Exotoxin derived from the parental strain contained 1.56 mg/mL of protein and  $1.02 \times 10^6$  endotoxin units (EU)/mL of LPS. Exotoxin from the mutant strain contained 1.50 mg/mL of protein and  $9.99 \times 10^5$  EU/mL of LPS.

### SDS-PAGE and Western blot analysis

Twenty-five  $\mu\text{g}$  of total protein from *A. pleuropneumoniae* exotoxin was separated by SDS-PAGE and stained with Coomassie Brilliant Blue R-250 (Sigma-Aldrich). For Western blot analysis, exotoxin with 2.5  $\mu\text{g}$  protein was subjected to SDS-PAGE and transfer to a polyvinylidene difluoride (PVDF) membrane (PerkinElmer). The proteins were detected with a rabbit antiserum raised against recombinant subunit ApxI [4].

### Hemolytic assay

Hemolytic activity of the exotoxin was evaluated according to a protocol described elsewhere with minor modifications [9]. Heparinized whole blood was collected from a pig. Red blood cells (RBCs) were washed with Tris-saline buffer [TS buffer; 10 mM Tris and 0.8% NaCl (pH 7.5)] and resuspended in TS buffer supplemented with 1 mM of  $\text{CaCl}_2$  (TS/ $\text{CaCl}_2$ ) at a ratio of 1 : 100. Exotoxin was 2-fold serially diluted with TS/ $\text{CaCl}_2$  buffer and incubated with washed RBCs at 37°C for 2 h. Percentage of hemolysis was calculated by measuring absorbance at 550 nm using a plate reader (Dynex Technologies, USA).

### Lactate dehydrogenase (LDH) release and mitochondrial activity assays

Membrane-damaging effects of the exotoxin were evaluated in PAMs using an LDH release assay (Roche, Switzerland) based on the manufacturer's instructions. Briefly,  $2.5 \times 10^4$  PAMs were incubated with various concentrations of exotoxin at 37°C for 4 h. Levels of LDH in the culture medium were determined with the LDH assay. Effects of the exotoxin on mitochondrial activity were evaluated in the PAMs with an XTT assay as previously described [4].

### Apoptosis scoring

PAMs were plated onto a 12-mm coverslip coated with 25  $\mu\text{g}/\text{mL}$  PDL at a density of  $2.5 \times 10^5$  cells per well and stimulated with bacterial cultural supernatants with protein concentrations of 1, 2.5, or 5  $\mu\text{g}/\text{mL}$  at 37°C for 6 h. Apoptosis was then monitored by evaluating nuclear condensation and/or fragmentation after Hoechst 33258 staining as previously described [4].

### Fifty percent lethal dose ( $\text{LD}_{50}$ ) in mice

Six-week-old BALB/c mice were purchased from the National Laboratory Animal Center (Taiwan, R.O.C.). The mice were cared for and handled according to the guidelines of the IACUC of National Chung Hsing University. The mice were intraperitoneally injected with 500  $\mu\text{L}$  of 10-fold serial-diluted bacterial suspension. Mortality was recorded 7 days post-inoculation, and the  $\text{LD}_{50}$  was calculated according to the method of Reed and Muench [20].


### Statistical analysis

Data were obtained from three independent experiments with at least duplicate determinations. Differences were analyzed using a one-way analysis of variance (ANOVA) with Microsoft Office Excel 2003 (Microsoft, USA). Results are expressed as the mean  $\pm$  standard error of the mean (SE).

## Results

### Identification of the *apxIA* insertion mutant

An *apxIA* insertion mutant was constructed through homologous recombination in *A. pleuropneumoniae* serotype 10. Genotype of a candidate mutant strain was characterized using PCR, Southern blot analysis, and DNA sequencing. Results of PCR using primers targeting *apxIA* indicated that the parental strain of *A. pleuropneumoniae* serotype 10 contained the full-length *apxIA* gene with a




**Fig. 2.** PCR results and Southern blot analysis of the *apxIA* insertion mutant. Genomic DNA of parental (P) or ApxIA336 mutant (M), or de-ionized water (W) was amplified by PCR with primers specific for the *apxIA* (A) or *apxIVA* (B) gene. Genomic DNA was also digested with the restriction enzyme *PvuI* and subjected to Southern blot analysis using DIG-labeled probe specific to *apxIA* (C) or kanamycin-resistant (D) gene.

size of 3137 bp. PCR product generated from the mutant strain was 4389 bp, indicating that a kanamycin-resistant cassette had been inserted into the *apxIA* gene (Fig. 2A). PCR amplification of *apxIVA* was also performed to confirm that both the mutant and parental strains were *A. pleuropneumoniae* (Fig. 2B).

Insertion of the kanamycin-resistant gene cassette into *apxIA* introduced a *PvuI* restriction enzyme cleavage site within the genomic DNA of the mutant strain (Fig. 1). Southern blot analysis confirmed that only one DNA restriction fragment hybridized with the DIG-labeled *apxIA* probe in the parental strain while two hybridized restriction fragments were observed for the mutant strain (Fig. 2C). Additionally, a probe specific for the kanamycin-resistant gene hybridized with two DNA fragments from the mutant strain but not the parental strain (Fig. 2D).

Finally, sequencing analysis revealed that the kanamycin-resistant cassette had been inserted into a region immediately after 1005 bp of the *apxIA* gene in the mutant strain (data not shown). The insertion created a frame-shift mutation in the *apxIA* gene. The mutant strain was designated as ApxIA336.




**Fig. 3.** Growth curve analysis and stability of the kanamycin-resistant gene in the ApxIA336 mutant. (A) Growth rate of ApxIA336 or the parental strain was monitored for 7 h and the number of colony forming units (CFUs) was determined. Data are from three independent experiments. Error bars represent the standard error (SE). (B) Presence of the kanamycin-resistant gene in ApxIA336 10 serial passages was detected by PCR. De-ionized water (W) served as a negative control. The arrow indicates the amplified fragment corresponding to the kanamycin-resistant cassette (633 bp). NS: no significant difference between the two strains.

### Insertion of the kanamycin-resistant cassette does not affect ApxIA336 growth


In this study, we also attempted to determine whether insertion of the kanamycin-resistant cassette into the *apxIA* gene affected bacterial growth. For this, the mutant and parental strains were cultured, and the growth curves were monitored. Experimental results indicated that the growth rates of both strains were similar and reached a plateau at 5 h post-inoculation (Fig. 3A). Stability of the insertion mutation was further evaluated by identifying the kanamycin-resistant gene by PCR in ApxIA336 cultures subjected to 10 serial passages. A fragment of expected size (633 bp) was observed in bacteria from all 10 passages, and provided evidence of the stability of the insertion in ApxIA336 (Fig. 3B).

### ApxIA336 does not produce ApxI

Disruption of ApxI production in the ApxIA336 mutant was demonstrated using SDS-PAGE and Western blotting. SDS-PAGE analysis of bacterial cultural supernatants revealed multiple proteins produced by both the parental and mutant strains (Fig. 4A). Western blot analysis indicated that when applying an antiserum raised against the recombinant subunit ApxI [4], only a single band was observed in the parental strain, but not in the mutant strain (Fig. 4B). This finding confirmed that ApxIA336 failed to produce exotoxin ApxI. Additionally, the levels of protein and LPS in cultural supernatants from the mutant and parental strains appeared to be comparable (see Materials and Methods).



**Fig. 4.** ApxIA336 failed to produce ApxI. Cultural supernatant from the parental (P) or ApxIA336 mutant (M) strain was subjected to SDS-PAGE (A) and Western blot analysis (B). Arrows indicate the band corresponding to ApxI at 105 kDa.



**Fig. 5.** ApxIA336 cultural supernatant failed to induce hemolysis or PAM damage. Porcine RBCs were treated with different concentrations of the cultural supernatants derived from ApxIA336 or parental strain in hemolytic assay (A). Effects of ApxIA336 or parental cultural supernatant on PAMs were evaluated in LDH release (B), mitochondrial activity (C), and apoptosis (D) assays. Results are from three independent experiments of at least duplicate determinations. Error bars represent the SE. \* $p < 0.05$ , \*\* $p < 0.01$ , and \*\*\* $p < 0.001$ .

### Cultural supernatant from ApxIA336 fails to induce hemolysis

Hemolytic activity of cultural supernatants from both the mutant and parental strains was assessed using porcine RBCs. Notably, ApxIA336 cultural supernatant did not induce hemolysis even at a protein concentration of 200 µg/mL. Conversely, the parental culture supernatant containing 3.1 to 200 µg/mL of protein caused 10 to 100% of the RBCs undergo hemolysis (Fig. 5A). These results indicated that without the production of ApxI, cultural supernatant from ApxIA336 had no lytic effect on porcine erythrocytes.

### Cultural supernatant from ApxIA336 does not affect cell membrane integrity or mitochondrial activity in PAMs

Damaging effects of the bacterial cultural supernatants on cell membrane integrity and mitochondrial activity were

evaluated. For this, LDH release and XTT assays, respectively, were performed with PAMs as the target cells. Experimental results indicated that cultural supernatant from ApxIA336 did not adversely affect cell membrane integrity or mitochondrial activity even at protein concentrations of 200 µg/mL, the highest concentration tested (Figs. 5B and C).

In contrast, the level of LDH release was elevated with increasing protein concentrations of the parental cultural supernatant (Fig. 5B). At 50 µg/mL, the parental supernatant induced 100% LDH release, indicative of extensive membrane damage to the PAMs. Treatment with the parental supernatant also impaired PAM mitochondrial activity in a concentration-dependent manner. Moreover, parental supernatant containing 12.5 to 200 µg/mL of protein caused a 25 to 100% of decrease in mitochondrial activity (Fig. 5C).

**Table 2.** LD<sub>50</sub> of the parental or ApxIA336 strain in BALB/c mice

Bacterial strain	Challenge dose (CFUs/mouse)	Dead/Total	LD <sub>50</sub> (CFUs)
Parental	1.3 × 10 <sup>6</sup>	0/5	1.9 × 10 <sup>7</sup>
	1.3 × 10 <sup>7</sup>	2/5	
	1.3 × 10 <sup>8</sup>	5/5	
ApxIA336	1.2 × 10 <sup>7</sup>	0/5	2.8 × 10 <sup>8</sup>
	1.2 × 10 <sup>8</sup>	1/5	
	1.2 × 10 <sup>9</sup>	5/5	

### Cultural supernatant of ApxIA336 fails to induce apoptosis in PAMs

The ability of ApxI in cultural supernatant to induce apoptosis was determined by treating PAMs with various concentrations of bacterial cultural supernatants and scoring for apoptosis. Cultural supernatant derived from ApxIA336 with protein concentrations ranging from 1 to 5 µg/mL induced apoptosis at a rate of 3.5 to 4.9%, which did not significantly differ from that of the untreated control group (Fig. 5D). PAMs treated with parental cultural supernatant containing 1, 2.5, or 5 µg/mL of protein induced apoptosis at a rate of 6.2%, 17.1%, and 41.3%, respectively (Fig. 5D). These results indicated that cultural supernatant from ApxIA336 failed to induce apoptosis in PAMs.

### The virulence of ApxIA336 is attenuated in mice

Bacterial virulence was evaluated by intraperitoneally injecting BALB/c mice with various concentrations of the mutant or parental strain. The LD<sub>50</sub> values of the mutant and parental strains were 2.8 × 10<sup>8</sup> and 1.9 × 10<sup>7</sup> CFUs, respectively (Table 2). In the mice, the mutant strain ApxIA336 was 15 times less virulent than the parental strain.

## Discussion

Apx exotoxins produced by *A. pleuropneumoniae* are virulence factors that profoundly impact the pathogenesis of porcine pleuropneumonia [13,21]. Given the heat-labile nature of the exotoxins [13], bioactive native Apx is not easily obtained after purification [5,6]. Our previous study investigated how a single species of exotoxin affects PAMs using crude cultural supernatant derived from *A. pleuropneumoniae* serotype 10 as the source of native ApxI. Specificity of the apoptotic effect exerted by ApxI has been demonstrated by using several controls, including neutralization with antibody, heat-inactivation of the toxin, and the addition of polymyxin B [2,4,25]. Alternatively, an *apxIA*-insertion mutant derived from *A. pleuropneumoniae* serotype 10 would be valuable for further elucidating the effects of ApxI on target cells.

In the present study, ApxIA336 mutant strain was constructed through homologous recombination that resulted in the insertion of a kanamycin-resistant gene cassette into the *apxIA* gene, leading to the disruption of gene expression. A full-length *apxIA* gene consists of 3069 bp. The insertion occurred immediately after 1005 bp, which was approximately one-third of the length from the 5'-end of the *apxIA* gene and caused a frame-shift in the *apxIA* open reading frame. The disruption of ApxI production in mutant strain was confirmed by Western blot analysis in which rabbit antiserum raised against amino acids 37-626 of ApxI [4] failed to recognize any protein in the ApxIA336 cultural supernatant (Fig. 4B).

The observation that ApxIA336 had a similar growth rate as that of the parental strain indicates that the insertion mutation did not affect overall bacterial growth. This finding is similar to previous data from Xu *et al.* [26] who constructed a mutant strain of *A. pleuropneumoniae* serotype 10 by replacing the 1.8-kb gene fragment at the C-terminal of *apxIA* with a chloramphenicol-resistant cassette. The resulting mutant strain produced a truncated form of ApxI protein, while its growth rate was comparable to that of the parental strain. This same study demonstrated that the mutant strain did not induce hemolysis when cultured on blood agar with *Staphylococcus epidermidis*. To the best of our knowledge, the cytotoxic effects of exotoxin derived from the mutant strain [26] have not been previously addressed.

The present investigation demonstrated that ApxIA336 did not produce ApxI. However, multiple protein bands were observed in cultural supernatants from both the parental and mutant strains by SDS-PAGE. Additionally, a similar amount of LPS was found in the cultural supernatants of both parental and mutant strains. However, without ApxI, a mixture of multiple proteins and LPS in the cultural supernatant apparently did not cause hemolysis, damage cell membranes, impair mitochondrial activity, or induce the apoptosis of PAMs. These findings support the notion that ApxI, rather than LPS or other components produced by the bacteria, is largely responsible for the hemolytic and cytotoxic effects of crude exotoxin preparations. It is still unclear whether LPS potentiates the effects of ApxI. Nevertheless, a previous study determined that LPS enhances β2 integrin expression on bovine neutrophils, subsequently increasing the cytotoxic effect of *M. haemolytica* LktA [17].

The ApxIA336 mutant constructed in our study was subjected to virulence analysis in mice. According to the results, virulence of the mutant strain was attenuated compared to the parental strain. The reduction of virulence in mice was similar to the findings from a study by Xu *et al.* [26] in which 80% of the mice survived after being challenged with 2 × 10<sup>8</sup> CFUs of a mutant strain while only 40% of the mice survived a challenge with 2 × 10<sup>6</sup> CFUs of

the parental strain. Notably, in both studies, the parental strain exhibited a higher virulence in mice than the *apxIA*-mutant strains, indicating a potential role of ApxI in bacterial virulence, despite the fact that mice are not a natural host of *A. pleuropneumoniae*. Nevertheless, the mouse model has been extensively adopted in evaluating the vaccine efficacy against *A. pleuropneumoniae* [3,16,18]. Whether the virulence of ApxIA336 is attenuated in pigs warrants further study.

In conclusion, disruption of ApxI production corresponds to the loss of hemolytic and ability to damage cells of the culture supernatant from ApxIA336, indicating a pivotal role of ApxI in these effects. Without ApxI, LPS in conjunction with other components in bacterial cultural supernatant failed to elicit hemolytic or cytotoxic effects towards RBCs or PAMs, respectively. Mutant ApxIA336 is highly promising for use in future studies designed to elucidate how ApxI affects target cells.

## Acknowledgments

The authors would like to thank Dr. Kuang-Sheng Yeh (Department of Veterinary Medicine, National Taiwan University) for kindly donating a kanamycin-resistant cassette gene from the pUC4K plasmid. Ted Knoy is appreciated for his editorial assistance. This work was supported in part by a grant (Aiming for Top University Plan) from National Chung Hsing University and by a grant NSC101-3114-Y-466-002 from the National Science Council of the Republic of China.

## References

1. **Bandara AB, Lawrence ML, Veit HP, Inzana TJ.** Association of *Actinobacillus pleuropneumoniae* capsular polysaccharide with virulence in pigs. *Infect Immun* 2003, **71**, 3320-3328.
2. **Chen ZW, Chien MS, Chang NY, Chen TH, Wu CM, Huang C, Lee WC, Hsuan SL.** Mechanisms underlying *Actinobacillus pleuropneumoniae* exotoxin ApxI induced expression of IL-1 $\beta$ , IL-8 and TNF- $\alpha$  in porcine alveolar macrophages. *Vet Res* 2011, **42**, 25.
3. **Chiang CH, Huang WF, Huang LP, Lin SF, Yang WJ.** Immunogenicity and protective efficacy of ApxIA and ApxIIA DNA vaccine against *Actinobacillus pleuropneumoniae* lethal challenge in murine model. *Vaccine* 2009, **27**, 4565-4570.
4. **Chien MS, Chan YY, Chen ZW, Wu CM, Liao JW, Chen TH, Lee WC, Yeh KS, Hsuan SL.** *Actinobacillus pleuropneumoniae* serotype 10 derived ApxI induces apoptosis in porcine alveolar macrophages. *Vet Microbiol* 2009, **135**, 327-333.
5. **Czuprynski CJ, Welch RA.** Biological effects of RTX toxins: the possible role of lipopolysaccharide. *Trends Microbiol* 1995, **3**, 480-483.
6. **Devenish J, Rosendal S.** Identification of the heat-labile hemolysin of *Actinobacillus pleuropneumoniae* serotype 1. *Can J Vet Res* 1989, **53**, 251-254.
7. **Dubreuil JD, Jacques M, Mittal KR, Gottschalk M.** *Actinobacillus pleuropneumoniae* surface polysaccharides: their role in diagnosis and immunogenicity. *Anim Health Res Rev* 2000, **1**, 73-93.
8. **Frey J.** Virulence in *Actinobacillus pleuropneumoniae* and RTX toxins. *Trends Microbiol* 1995, **3**, 257-261.
9. **Frey J, Nicolet J.** Hemolysin patterns of *Actinobacillus pleuropneumoniae*. *J Clin Microbiol* 1990, **28**, 232-236.
10. **Frey J, Beck M, Stucki U, Nicolet J.** Analysis of hemolysin operons in *Actinobacillus pleuropneumoniae*. *Gene* 1993, **123**, 51-58.
11. **Frey J, Bosse JT, Chang YF, Cullen JM, Fenwick B, Gerlach GF, Gygi D, Haesebrouck F, Inzana TJ, Jansen R, Kamp EM, Macdonald J, MacInnes JI, Mittal KR, Nicolet J, Rycroft AN, Segers RPAM, Smits MA, Stenbaek E, Struck DK, van den Bosch JF, Willson PJ, Young R.** *Actinobacillus pleuropneumoniae* RTX-toxins: uniform designation of haemolysins, cytolysins, pleurotoxin and their genes. *J Gen Microbiol* 1993, **139**, 1723-1728.
12. **Jansen R, Briaire J, Smith HE, Dom P, Haesebrouck F, Kamp EM, Gielkens ALJ, Smits MA.** Knockout mutants of *Actinobacillus pleuropneumoniae* serotype 1 that are devoid of RTX toxins do not activate or kill porcine neutrophils. *Infect Immun* 1995, **63**, 27-37.
13. **Kamp EM, van Leengoed LAMG.** Serotype-related differences in production and type of heat-labile hemolysin and heat-labile cytotoxin of *Actinobacillus (Haemophilus) pleuropneumoniae*. *J Clin Microbiol* 1989, **27**, 1187-1191.
14. **Kamp EM, Popma JK, Anakotta J, Smits MA.** Identification of hemolytic and cytotoxic proteins of *Actinobacillus pleuropneumoniae* by use of monoclonal antibodies. *Infect Immun* 1991, **59**, 3079-3085.
15. **Kamp EM, Stockhofe-Zurwieden N, van Leengoed LAMG, Smits MA.** Endobronchial inoculation with Apx toxins of *Actinobacillus pleuropneumoniae* leads to pleuropneumonia in pigs. *Infect Immun* 1997, **65**, 4350-4354.
16. **Kim JM, Jung DI, Eom YJ, Park SM, Yoo HS, Jang YS, Yang MS, Kim DH.** Surface-displayed expression of a neutralizing epitope of ApxIIA exotoxin in *Saccharomyces cerevisiae* and oral administration of it for protective immune responses against challenge by *Actinobacillus pleuropneumoniae*. *Biosci Biotechnol Biochem* 2010, **74**, 1362-1367.
17. **Leite F, Gyles S, Atapattu D, Maheswaran SK, Czuprynski CJ.** Prior exposure to *Mannheimia haemolytica* leukotoxin or LPS enhances  $\beta_2$ -integrin expression by bovine neutrophils and augments LKT cytotoxicity. *Microb Pathog* 2003, **34**, 267-275.
18. **Liao CW, Chiou HY, Yeh KS, Chen JR, Weng CN.** Oral immunization using formalin-inactivated *Actinobacillus pleuropneumoniae* antigens entrapped in microspheres with aqueous dispersion polymers prepared using a co-spray drying process. *Prev Vet Med* 2003, **61**, 1-15.
19. **Maier E, Reinhard N, Benz R, Frey J.** Channel-forming activity and channel size of the RTX toxins ApxI, ApxII, and ApxIII of *Actinobacillus pleuropneumoniae*. *Infect Immun* 1996, **64**, 4415-4423.



20. **Reed LJ, Muench H.** A simple method of estimating fifty percent endpoints. *Am J Hyg* 1938, **27**, 493-497.
21. **Rosendal S, Devenish J, MacInnes JI, Lumsden JH, Watson S, Xun H.** Evaluation of heat-sensitive, neutrophil-toxic, and hemolytic activity of *Haemophilus (Actinobacillus) pleuropneumoniae*. *Am J Vet Res* 1988, **49**, 1053-1058.
22. **Schaller A, Kuhn R, Kuhnert P, Nicolet J, Anderson TJ, MacInnes JI, Segers RPAM, Frey J.** Characterization of *apxIVA*, a new RTX determinant of *Actinobacillus pleuropneumoniae*. *Microbiology* 1999, **145**, 2105-2116.
23. **Strathdee CA, Lo RYC.** Extensive homology between the leukotoxin of *Pasteurella haemolytica* A1 and the alpha-hemolysin of *Escherichia coli*. *Infect Immun* 1987, **55**, 3233-3236.
24. **Welch RA.** Pore-forming cytolysins of gram-negative bacteria. *Mol Microbiol* 1991, **5**, 521-528.
25. **Wu CM, Chen ZW, Chen TH, Liao JW, Lin CC, Chien MS, Lee WC, Hsuan SL.** Mitogen-activated protein kinases p38 and JNK mediate *Actinobacillus pleuropneumoniae* exotoxin ApxI-induced apoptosis in porcine alveolar macrophages. *Vet Microbiol* 2011, **151**, 372-378.
26. **Xu F, Chen X, Shi A, Yang B, Wang J, Li Y, Guo X, Blackall PJ, Yang H.** Characterization and immunogenicity of an *apxIA* mutant of *Actinobacillus pleuropneumoniae*. *Vet Microbiol* 2006, **118**, 230-239.