


RESEARCH PAPER

 OPEN ACCESS 

MiR-455 targeting SOCS3 improve liver lipid disorders in diabetic mice

Shu Fang^{a*}, Jie Feng^{b*}, Hongbin Zhang^c, Ping Li^a, Yudan Zhang ^a, Yanmei Zeng^a, Yingying Cai^{a,d}, Xiaochun Lin^a, Yaoming Xue^a, and Meiping Guan^a

^aDepartment of Endocrinology & Metabolism, Nanfang Hospital, Southern Medical University, Guangzhou, Guangdong, China; ^bDepartment of Medical Imaging Center, Nanfang Hospital, Southern Medical University, Guangzhou, Guangdong, China; ^cDepartment of Biomedical Sciences, University of Copenhagen, Copenhagen, Denmark; ^dWomen and Children's Hospital, School of Medicine, Xiamen University, Fujian, China

ABSTRACT

MiR-455 has been verified a key regulator of brown adipose tissue and adipose tissue-specific overexpression of miR-455 (ap2-miR-455) mice could combat high-fat-diet-induced obesity. This study is to verify overexpression of miR-455 could ameliorate the lipid accumulation and metabolism in the liver of db/db diabetic mice and explore the potential mechanisms. Diabetic mice (db/db) and control mice (db/m) were randomly divided into four groups. After overexpression of miR-455 in the liver of db/db mice, the triglycerides level in both serum and liver decreased, the lipid deposit in liver was improved, the expression of fatty acid synthase, stearoyl-CoA desaturase 1, sterol regulatory element binding protein 1c (SREBP-1c) and acetyl-CoA carboxylase (ACCα) was also significantly down-regulated. TargetScan indicated that suppressor of cytokine signalling 3 (SOCS3) is predicated to target miR-455 and the protein of SOCS3 in the liver of db/db mice after intervention was significantly decreased. The dual luciferase reporter assay showed that SOCS3 was target gene of miR-455. In vitro, in Palmitate (PA)-stimulated human normal liver (LO2) cells, transfected miR-455 mimic could significantly inhibit the expression of SOCS3, while transfected miR-455 inhibitor could up-regulate the expression of SOCS3. Transfecting LO2 cells with siRNA of SOCS3 could significantly down-regulate the protein expression of SREBP-1c and ACCα. Our study showed that overexpression of miR-455 in the liver could improve lipid metabolism in diabetic mice by down-regulating its target gene SOCS3.

ARTICLE HISTORY

Received 1 December 2019
Revised 20 March 2020
Accepted 23 March 2020

KEYWORDS

Diabetes; NAFLD; miR-455; lipid metabolism; SOCS3



Introduction

Non-alcoholic fatty liver disease (NAFLD) is closely related to diabetes. The estimated global prevalence of NAFLD is 25%, which is as high as 30% in some countries [1]. Recent clinical researches have revealed that diabetes plays an important role in the progression of NAFLD [2,3] and has become a strong predictor of non-alcoholic steatohepatitis (NASH) and liver fibrosis [4]. However, there is still a lack of effective treatment for NAFLD [5].

MicroRNA (miRNA) is a non-coding single-stranded RNA molecule of approximately 20–24 nucleotides in length, regulates gene expression at the post-transcriptional level by binding to the mRNA of the target gene [6]. MiRNA can regulate a variety of biological processes such as developmental processes, tumorigenesis and cell proliferation and apoptosis, not only participates in the process of multiple diseases, but also serves as a marker molecule for diagnosing diseases [7]. MiRNAs have been used as the clinical diagnostic markers for NAFLD/NASH, and even the potential predictors of primary liver cancer

[8]. Our previous study shows that miR-455 is a key regulator of brown adipose tissue that promotes differentiation of brown adipocytes by targeting *Necdin*, *Runx1t1* and *HIF1an* [9]. Recent studies indicate that miR-455 could improve the progression of different cancers [10,11], while there is still lack of the research focused on the regulation of miR-455 on liver metabolism.

In this study, we observed the changes in liver lipid metabolism in diabetic mice (db/db) after the overexpression of miR-455 in the liver. Furthermore, we found that miR-455 could improve the liver metabolism by inhibiting suppressor of cytokine signalling 3 (SOCS3), as a member of SOCS family, negatively regulates the transduction of cytokine signalling [12]. At the same time, SOCS3 also plays an important role in the regulation of metabolism. The overexpression of miR-19a [13] and miR-125a [14] could improve the metabolism of glucose and lipid through targeting SOCS3. We found that miR-455 ameliorated the lipid metabolic disorder of liver in db/db mice and this improvement may be caused by inhibition of SOCS3.

CONTACT Meiping Guan  mpguan@163.com  Department of Endocrinology & Metabolism, Nanfang Hospital, Southern Medical University, #1838 Guangzhou Avenue North, Guangzhou, Guangdong 510515, China

*These authors contributed equally to this work.

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Methods

Animal experiments

All animal experiments were approved by the Animal Care and Use Committee of the Southern Medical University. 20 male C57BLKS/J diabetic mice (db/db; 6 weeks) and 20 male C57BLKS/J normal mice (db/m; 6 weeks) were purchased from the Nanjing Model Animal Centre. Mice were raised in the specific pathogen free barrier facility at the Southern Medical University under the controlled environment (temperature 22°C; 12 hours light/dark cycle). All mice were fed a standard chow diet (Guangzhou, China) and water.

After 1 week of adaptive culture, the mice were randomly divided into four groups: db/db-455 mice ($n = 10$), db/db-con mice ($n = 10$), db/m-455 mice ($n = 10$) and db/m-con mice ($n = 10$). The db/db-455 and db/m-455 mice were injected via tail vein with a plasmid construct expressing miR-455 (pCMV-miR-455)-miR-455 (Obio Technology, Shanghai) diluted with phosphate buffered saline and the db/db-con and db/m-con mice were given a pCMV (Obio Technology, Shanghai) injection. All mice received weekly injection for 4 weeks. When the experiment was over, the blood and liver tissue were collected after the mice were anesthetized by 1% pentobarbital sodium.

Metabolic measurement

The body weight (BW) and the random blood glucose (RBG) were recorded every 2 weeks. The serum and liver triglyceride (TG) content were detected by the Triglyceride detection kit (Jiancheng, Nanjing).

Cell culture and transfection

Human normal liver cell line (LO2) and human embryonic kidney 293 cell line (HEK293) were obtained from the Cell Bank of the Chinese Academy of Sciences (Shanghai, China). All cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) medium (Gibco, USA) supplemented with 10% foetal bovine serum (FBS) (Gibco, USA) and 1% penicillin-streptomycin in a 5% CO₂ atmosphere at 37°C. Palmitate (PA) (Sigma, USA) was used to mimic the hyperlipidaemia in cells. After stimulated with PA for 24 hours, the cells were transfected with miR-455 mimics, miR-455 inhibitors, small interfering RNAs (siRNAs) and negative control (RiboBio, Guangzhou) respectively for 48 hours, and the transfection was performed by Lipofectamine 3000 reagent (Invitrogen, MA).

Bioinformatics prediction and dual luciferase reporter assay

TargetScan (<http://www.targetscan.org/>) was used to predict the potential targets of miR-455, the target site of miR-455-5p in SOCS3 3'-UTR was shown in Figure 4(g). Luciferase reporter plasmid wide-type (wt) or mutated-type (mut) were co-transfected into the 293 T cells with miR-455 mimics, respectively. The luciferase activity of the cells was measured by Luciferase Reporter Assay System (Promega Corporation) after 48 hours of transfection.

Quantitative real-time PCR (qRT-PCR)

Total RNA was extracted from the cells and liver tissue by Trizol (Invitrogen, MA) and cDNA was synthesized by PrimeScript RT Master Mix (Takara, Japan). The cDNA of miR-455 and U6 was synthesized by MiRNA cDNA First Strand Synthesis Kit (TianGen, China). The qRT-PCR was performed with SYBR Green real-time Kit (Takara, Japan) and qRT-PCR of miR-455 and U6 was performed with SYBR Green real-time Kit (TianGen, China) on Roche LightCycler 480 Real-Time PCR System. β -actin was served as housekeeping gene in mouse liver and LO2 cells. U6 were used as housekeeping gene in compared with miR-455. Collect CT values of genes, and then use the $2^{-\Delta\Delta Ct}$ method to analyse the expression of all genes. All primers were listed in Tables 1 & 2.

Western blot analysis

The protein of cells and liver tissue was extracted by RIPA lysis buffer (KeyGen, China) and the concentration of protein was quantified by BCA Protein Assay Kit (Takara, Japan). The denatured protein (20 μ g) was separated by 10% sodium dodecyl sulphate-polyacrylamide gel electrophoresis and transferred onto the polyvinylidene difluoride membrane. After being blocked with 5% skim milk for 1 hour at ambient temperature, the membrane was incubated with primary antibodies of sterol regulatory element binding protein 1c (SREBP-1c) (1:1000, Abclone, China), acetyl-CoA carboxylase (ACC α) (1:1000, Abclone, China), fatty acid synthase (FAS) (1:1000, Abclone, China), NADPH (nicotinamide adenine dinucleotide phosphate oxidase) oxidase 4 (NOX4) (1:1000, Abclone, China), SOCS3 (1:1000, Abclone, China), IL-6 (1:1000, CST, USA) and β -actin (1:1000, Zhongshan, China) at 4°C overnight. Then the membrane was incubated with horseradish peroxidase labelled secondary antibody (1:5000) for 1 hour after being washed three times with TBST (tris-buffered saline Tween-20). Enhanced chemiluminescence reagent (Millipore, USA) was used to detect

Table 1. Primer sequences (mouse).

Gene	Forward prime (5'-3')	Reverse prime (5'-3')
β-actin	GTCCACCCCGGGGAAGGTGA	AGGCCTCAGACCTGGGCCATT
PPAR γ	ATTCTGGCCACCAACTTCGG	CTGCGAAACTTCAGGAAATG
FAS	GGCCAGTGTATGCTGAGAT	CACCTCTGGAGACATCGAAAC
CD36	ATGGGCTGTGATCGGAACCTG	TTGCCACGTCATCTGGGTTT
SCD-1	CTGGCATTGTTCGGTTCT	CGGGATTGAATGTTCTGTCTG
SREBP-1c	TGGAAGCCTGATGCTTTATCCCCA	GGTTCGGAATGCTATCCAGG
ACC α	TATTCGGCTGAAGCTGGTGTAC	TTCTTGCATACACTCTGGTGC
TNF- α	CATCTTCTAAAATTCGAGTGACAA	TGGGAGTAGACAAGGTACAACCC
MCP-1	GTTAACGCCCACTCACCTG	GGGCCGGGTATGTAACCTCA
NOX4	TGCTGTCTCATTTGGCTGT	CCGGCACATAGGTAAAAGGATG
RUN6	TGGCCCCTGCGCAAGGATG	/
MiR-455-5p	TATGTGCCTTTGGACTACATCG	/

Table 2. Primer sequences (human).

Gene	Forward prime (5'-3')	Reverse prime (5'-3')
β-actin	CATGTACGTTGCTATCCAGGC	CTCCTTAATGTCACGCACGAT
PPAR γ	GGGATCAGCTCCGTGGATCT	TGCACCTTGGTACTCTTGAAGTT
FAS	ATGTCAGTCACTGGGCATTA	CATCTGGACCCTCTACCTCT
ACC α	ATGCTGGCTTGCACCTAGTA	CCCCAAAGCGAGTAACAAATCT
SREBP-1c	CGGAACCATCTTGGCAACAGT	CGCTTCTCAATGGCGTGT
PPAR α	CTGGCATTGTTTCTGTTCTTT	CTCCTCGGTGACTTATCTGT
CPT-1	TCCAGTTGGCTTATCGTGGTG	TCCAGAGTCCGATTGATTTTTGC
IL-6	ACTCACCTCTTCAAGCAACGAATTG	CCATCTTGGAAAGGTTCAAGTTG
TNF- α	CCTCTCTAATCAGCCCTCTG	GAGGACCTGGGAGTAGATGAG
NOX4	CAGATGTTGGGCTAGGATTG	GAGTGTTCGGCACATGGGTA

immunoreactive bands and the Image J software was used to analyse the bands.

Histological staining

The liver tissue was first fixed with 4% paraformaldehyde, then dehydrated and embedded in paraffin. The tissue embedded in the paraffin was subjected to H&E staining, and the freshly fixed tissue was stained with oil red O. Olympus B \times 20 upright light microscope (Olympus, Tokyo, Japan) was used to detect all sections.

Statistical Analysis

Data were expressed as the mean \pm SD. Two-tailed Student's *t* test was used to compare two independent groups, and one-way ANOVA was used to examine the differences among four independent groups. Statistical analysis was performed by using SPSS 20.0. *P* < 0.05 was considered statistically significant.

Results

The changes of basal metabolic indicator in mice after the intervention of miR-455

Db/db mice showed obvious obesity after 6 weeks and polydipsia, polyphagia and polyuria. The weight (Figure 1(a)) and RBG (Figure 1(b)) of db/db mice were also

significantly higher than those of db/m mice, indicating the establishment of diabetes model. After intervention, the expression of miR-455 was significantly increased in the liver of db/db-455 mice and db/m-455 mice compared to db/db-con mice and db/m-con mice respectively (Figure 1(c)). During the whole experiment, the BW (Figure 1(d)) and RBG (Figure 1(e)) of db/db-con and db/db-455 mice were higher than those of db/m mice. Although there was no obvious improvement in the levels of BW and blood glucose in db/db-455 mice, the overexpression of miR-455 in liver significantly reduced serum TG levels (Figure 1(f)) and liver TG content (Figure 1(g)) compared with db/db-con mice.

Overexpression of miR-455 improved the liver lipid metabolism

Compared with db/m-con mice, the mRNA expression of peroxisome proliferator-activated receptor γ (PPAR γ), FAS, fatty acid transporter (CD36) and stearyl-CoA desaturase 1 (SCD-1) was notable up-regulated in db/db-con mice, the mRNA expression of ACC α and SREBP-1c also increased although without statistical significant (Figure 2(a)). However, the mRNA expression levels of FAS, SCD-1, SREBP-1c and ACC α in db/db-455 mice were significantly lower than those in db/db-con group (Figure 2(a)). In addition, we detected the protein expression of SREBP-1c, ACC α and FAS in the liver of db/db-455 and db/db-con mice, the results were consistent with the expression trend of mRNA (Figure 3(b)). H&E staining showed liver structural disorder, vacuolar degeneration of hepatocytes and inflammatory infiltration in db/db-con mice (Figure 2(d)) compared with db/m-con mice (Figure 2(b)), but the overexpression of miR-455 in liver ameliorated these changes in db/db-455 mice (Figure 2(e)). Simultaneously, the oil red O staining revealed that lipid deposition in liver of db/db-con mice (Figure 2(h)) was severe than db/m-con mice (Figure 2(f)), while the overexpression of miR-455 significantly reduced the number and size of liver lipid droplets in db/db-455 mice (Figure 2(i)).




Figure 1. The changes of basic metabolism indicator of mice after intervention by miR-455. The BW (g) (a) and RBG (mmol/L) (b) of mice before intervention. (c) The expression of miR-455 in liver of mice. The BW (d) and the RBG (e) of mice after intervention. The serum (f) and liver TG (g) content of mice. $n = 6-8$ per group.

Overexpression of miR-455 ameliorated the liver inflammation and oxidative stress

We then detected the liver mRNA expression of tumour necrosis factor- α (TNF- α), monocyte chemoattractant protein-1 (MCP-1) and NADPH oxidase 4 (NOX4), as the indicators of inflammation, the high expression of TNF- α and MCP-1 in db/db-con mice was significantly reduced after miR-455 overexpres-

sion in the liver of db/db-455 mice (Figure 3(a)). Overexpression of miR-455 in liver also tremendously inhibited the mRNA expression of NOX4 in db/db-455 mice, an indicator of oxidative stress (Figure 3(a)). Also, the protein expression level of SREBP-1c, ACC α , FAS and NOX4 was significantly decreased in the liver of db/db-455 mice compared with db/db-con mice (Figure 3(b)).




Figure 2. Overexpression of miR-455 improved the liver lipid metabolism. (a) The mRNA expression of PPAR γ , FAS, CD36, SCD-1, SREBP-1c and ACC α in the liver of mice in four groups. H&E (b–e) and oil red O staining (f–i) of liver in four groups. Db/m-con group (b + f), db/m-455 group (c + g), db/db-con group (d + h), db/db-455 group (e + i). Magnification $\times 400$. $n = 6$ –8 per group. $**P < 0.01$ and $***P < 0.001$ vs db/m-con group; $\#P < 0.05$ vs db/db-con group.

Cytokine signal transduction inhibitor 3 (SOCS3) was the target gene of miR-455 in liver

Since the overexpression of miR-455 could improve the lipid metabolism, inflammation and oxidative stress of db/db mice, we predicted its possible target genes using

TargetScan software in order to investigate its underlying mechanism. We found that SOCS3 may be one of the target genes of miR-455, and the detail bioinformatic analysis for the target site of miR-455-5p in SOCS3 3'-UTR was shown in [Figure 4\(g\)](#). Next, we detected the protein




Figure 3. Overexpression of miR-455 ameliorated the liver inflammation and oxidative stress. (a): The mRNA expression of TNF- α , MCP-1 and NOX4 of liver in four groups. (b) The protein levels of SREBP-1c, ACC, FAS and NOX4 in liver of db/db-con group and db/db-455 group, the quantitative analysis results of protein levels. $n = 6-8$ per group. * $P < 0.05$ and ** $P < 0.01$ vs db/m-con group; # $P < 0.05$ and ## $P < 0.01$ vs db/db-con group.

levels of SOCS3 in mice. The result indicated that the overexpression of miR-455 in liver obviously inhibited the expression of SOCS3 in db/db-455 mice (Figure 4(a)).

To mimic the hyperlipidaemia in vitro, we incubated the LO2 cells with PA. Lipid metabolism-related gene expression levels were significantly higher at 0.4 mM PA stimulation than 0.2 mM PA (Figure 4(b)), so we choose a stimulus concentration of 0.4 mM and the oil red O staining also indicated the 0.4 mM was a suitable concentration (Figure 4(c,d)). On the basis of PA stimulation, cells were transfected with mimic and inhibitor of miR-455 respectively. The expression of miR-455 in LO2 cells increased significantly after treatment with miR-455 mimic (Figure 4(e)), also the expression of miR-455 decreased after intervention with miR-455 inhibitor (Figure 4(f)). The protein level of SOCS3 was remarkably down-regulation after intervention by miR-455 mimic in LO2 cells (Figure 4(e)). On the contrary, the protein level of SOCS3 was significantly up-regulated after intervention by miR-455 inhibitor (Figure 4(f)). Then we constructed the SOCS3 wt plasmids and mut plasmids, which were co-transfected into 293 T cells with miR-455 mimic. The luciferase assay showed that the luciferase activity was significantly decreased after co-transfection of miR-455 mimic and SOCS3 wt plasmids compared to co-

transfection with miR-455 mimic and SOCS3 mut plasmids (Figure 4(g)). These results demonstrated that SOCS3 was a target gene of miR-455 in liver.

The effect of knocking down the expression of SOCS3 in LO2 cells

In order to investigate whether knocking down SOCS3 could affect the metabolism of LO2 cells, we treated the LO2 cells with siRNA-SOCS3 on the basis of PA. The mRNA (Figure 5(a)) and protein (Figure 5(b)) expression of SOCS3 was tremendously down-regulation after siRNA-SOCS3 intervention. SiRNA intervention significantly reduced mRNA expression of FAS in cells (Figure 5(c)). Despite the mRNA expression of SREBP-1c, ACC α , TNF- α and interleukin-6 (IL-6) had a decreasing trend, it had not reached a significant difference. Nevertheless, siRNA intervention of SOCS3 significantly decreased the protein expression of SREBP-1c, ACC α and IL-6 (Figure 5(d)).

Discussion

The synthesis of hepatic TGs exceeds the rate of catabolism could lead to the accumulation of TGs. And excessive




Figure 4. SOCS3 was the target gene of miR-455 in liver. (a) The protein level of SOCS3 in liver in four groups and the quantitative analysis result ($n = 6-8$ per group). (b) The mRNA expression of PPAR γ , FAS, SREBP-1c, ACC α , PPAR α and CPT-1 in LO2 cells. The oil red O staining of cells stimulated with BSA (bovine serum albumin) (c) and 0.4 mM PA (d). (e) The expression of miR-455 and SOCS3 in LO2 cells after treatment of miR-455 mimic. (f) The expression of miR-455 and SOCS3 in LO2 cells after treatment of miR-455 inhibitor. (g) The binding sites of miR-455 and SOCS3. Luciferase reporter assay of the interaction between miR-455 and SOCS3 ($n = 3$ per group). * $P < 0.05$ vs db/m-con group; ### $P < 0.001$ vs db/db-con group. + $P < 0.05$ and ++ $P < 0.01$ vs BSA group. & $P < 0.05$ and &&& $P < 0.001$ vs mimic control group. ^ $P < 0.05$ vs inhibitor control group. * $P < 0.05$ and *** $P < 0.001$.

accumulation of TGs in hepatocytes contribute to the occurrence of NAFLD [15]. In our study, we observed obvious obesity and lipid deposition in liver of db/db mice consistent with the result of published studies [16,17]. MicroRNAs play an important role in a variety of diseases [18–20] including the lipid metabolic disorder [21,22], because they can regulate a wide range of growth and development processes [7]. Recent studies indicate that

miR-27a [23], miR-194 [24] and miR-146b [25] can target different genes to regulate the development of NAFLD. According to our previous research, miR-455 is the key regulator of BAT in the process of differentiation [9] and may have an effect on the regulation of lipid synthesis. Then, we achieved the overexpression of miR-455 to figure out whether it could improve the lipid metabolism of liver in db/db mice. The results showed that the serum and liver




Figure 5. The effect of knocking down the expression of SOCS3 in LO2 cells. The mRNA (a) and protein (b) expression of SOCS3 in LO2 cells after incubated with PA and siRNA-SOCS3. (c) The mRNA expression of SREBP-1c, ACC α , FAS, IL-6, TNF- α and NOX4 in LO2 cells after incubated with PA and siRNA-SOCS3. (d) The protein expression of SREBP-1c, ACC α and IL-6 in cells after c results. $n = 3$ per group. $\blacklozenge\blacklozenge P < 0.05$ and $\blacklozenge\blacklozenge\blacklozenge P < 0.01$ vs PA-control group.

TG content of db/db-455 mice was significantly decreased. Histological staining results also suggested that there was improvement in the structural disorder and lipid deposition in the liver of db/db-455 mice.

Liver is a central regulator of lipid metabolism in human [26]. Nowadays, a lot of researches have already confirmed that plenty of molecules are involved in the lipid metabolism, including PPAR γ , SREBP-1c, FAS and ACC α [26–28]. As the crucial genes for the development of NAFLD, these genes were significantly up-regulated in the liver of db/db mice. There was also a significant increase in CD36 expression, which involved in fatty acid transport [29]. However, increased expression of these genes was conspicuously inhibited after the overexpression of miR-455 in liver. SREBP-1c has been reported as a upstream signal to directly regulate ACC α and FAS

[30]. In our study, overexpression of miR-455 suppressed the protein levels of SREBP-1c, ACC α and FAS. Thus, miR-455 could ameliorate the lipid deposition in db/db mice by downregulating SREBP-1c and its downstream signal pathway.

SOCS3 has been reported involving in the progress of various diseases such as autoimmune diseases [31], cardiovascular diseases [32], metabolic diseases [33] and tumours [34], which is responsive to an increase in IL-6 and plays a negative regulatory role in cytokine signalling [12]. It has been shown that miR-19a-3p can promote cell proliferation and insulin secretion by targeting SOCS3 [13], overexpression of miR-125a-5p also inhibit the expression of SOCS3 to improve glycolipid metabolism in the liver [14]. In our study, miR-455 can significantly reduce serum and liver TG level without affecting BW

and blood glucose, the probable reason is that the cycle of the experiment is not long enough to significantly improve blood glucose. In the end of the animal experiment, we found that the RBG level of db/db-455 group has been decreasing, but there was no significant difference compared with db/db-con group. We confirmed that SOCS3 was a target gene of miR-455. And Inhibition of SOCS3 could down-regulate SREBP-1c, ACC α and FAS in LO2 cells incubated with PA. Studies have shown that overexpression of SOCS3 enhances the promoter activity of SREBP-1c and then promotes the transcription of SREBP-1c [35]. Therefore, combined with our in vitro and in vivo experiments, we considered that overexpression of miR-455 can regulate lipid synthesis genes such as SREBP-1c by inhibiting its target gene SOCS3, thereby improve liver lipid metabolism.

Liver inflammation and oxidative stress are very important mechanisms of NAFLD [36]. In our study, we observed a significant increase of TNF- α , MCP-1 and NOX4 in db/db mice, which were significantly decreased by overexpression of miR-455 in db/db mice. In vitro, IL-6 was further decreased by transfecting siRNA of SOCS3 in LO2 cells. It has been reported that SOCS3 could promote the expression of inflammatory factors by inhibiting the phosphorylation of JAK2/STAT3 signalling pathway in adipocytes [37].

In conclude, we found that overexpression of miR-455 in db/db mice could improve the lipid metabolism, inflammation and oxidative stress in liver by targeting SOCS3. MiR-455/SOCS3 pathway may be a potential target for the treatment of abnormal liver lipid metabolism in type 2 diabetes.

Disclosure statement

The authors declare that they have no competing interests with the contents of this article.

Data availability statement

The analysed data sets generated during this study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

All animal experiments were approved by the Animal Care and Use Committee of the Southern Medical University.

Funding

This work was supported by grants from the National Natural Science Foundation of China [No. 81870612, No.

81628004 and No. 81470047] and Guangdong Science and Technology Project [2019A1515011997.

ORCID

Yudan Zhang  <http://orcid.org/0000-0003-3311-789X>

References

- [1] Cohen JC, Horton JD, Hobbs HH. Human fatty liver disease: old questions and new insights[J]. *Science*. 2011;332(6037):1519–1523.
- [2] Wang W, Xu AL, Li ZC, et al. Combination of probiotics and salvia miltiorrhiza polysaccharide alleviates hepatic steatosis via gut microbiota modulation and insulin resistance improvement in high fat-induced NAFLD mice[J]. *Diabetes Metab J*. 2019;43:e69.
- [3] Gan L, Meng ZJ, Xiong RB, et al. Green tea polyphenol epigallocatechin-3-gallate ameliorates insulin resistance in non-alcoholic fatty liver disease mice[J]. *Acta Pharmacol Sin*. 2015;36(5):597–605.
- [4] Mcpherson S, Hardy T, Henderson E, et al. Evidence of NAFLD progression from steatosis to fibrosing-steatohepatitis using paired biopsies: implications for prognosis & clinical management.[J]. *J Hepatol*. 2015;62(5):1148–1155.
- [5] Yoon H. Pathogenesis and therapeutic approaches for non-alcoholic fatty liver disease[J]. *World J Hepatol*. 2014;6(11):800.
- [6] Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function[J]. *Cell*. 2004;116(2):281–297.
- [7] Gebert L, Macrae IJ. Regulation of microRNA function in animals[J]. *Nat Rev Mol Cell Biol*. 2019;20(1):21–37.
- [8] Lee CH, Kim JH, Lee SW. The role of MicroRNA in pathogenesis and as markers of HCV chronic infection[J]. *Curr Drug Targets*. 2017;18(7):756–765.
- [9] Zhang H, Guan M, Townsend KL, et al. MicroRNA-455 regulates brown adipogenesis via a novel HIF1 α -AMPK-PGC1 α signaling network[J]. *EMBO Rep*. 2015;16(10):1378–1393.
- [10] Li YJ, Ping C, Tang J, et al. MicroRNA-455 suppresses non-small cell lung cancer through targeting ZEB1[J]. *Cell Biol Int*. 2016;40(6):621–628.
- [11] Mao QD, Zhang W, Zhao K, et al. MicroRNA-455 suppresses the oncogenic function of HDAC2 in human colorectal cancer[J]. *Braz J Med Biol Res*. 2017;50(6):e6103.
- [12] Xiang S, Dong NG, Liu JP, et al. Inhibitory effects of suppressor of cytokine signaling 3 on inflammatory cytokine expression and migration and proliferation of IL-6/IFN- γ -induced vascular smooth muscle cells[J]. *J Huazhong Univ Sci Technolog Med Sci*. 2013;33(5):615–622.
- [13] Li Y, Luo T, Wang L, et al. MicroRNA-19a-3p enhances the proliferation and insulin secretion, while it inhibits the apoptosis of pancreatic beta cells via the inhibition of SOCS3[J]. *Int J Mol Med*. 2016;38(5):1515–1524.
- [14] Xu L, Li Y, Yin L, et al. miR-125a-5p ameliorates hepatic glycolipid metabolism disorder in type 2

- diabetes mellitus through targeting of STAT3[J]. *Theranostics*. 2018;8(20):5593–5609.
- [15] Birkenfeld AL, Shulman GI. Nonalcoholic fatty liver disease, hepatic insulin resistance, and type 2 diabetes[J]. *Hepatology*. 2014;59(2):713–723.
- [16] Wang B, Chandrasekera PC, Pippin JJ. Leptin- and leptin receptor-deficient rodent models: relevance for human type 2 diabetes[J]. *Curr Diabetes Rev*. 2014;10(2):131–145.
- [17] Sanches SC, Ramalho LN, Augusto MJ, et al. Nonalcoholic steatohepatitis: a search for factual animal models[J]. *Biomed Res Int*. 2015;2015:574832.
- [18] Garzon R, Calin GA, Croce CM. MicroRNAs in cancer[J]. *Annu Rev Med*. 2009;60(1):167–179.
- [19] Wojciechowska A, Braniewska A, Kozar-Kamińska K. MicroRNA in cardiovascular biology and disease[J]. *Adv Clin Exp Med*. 2017;26(5):865–874.
- [20] Rottiers V, Näär AM. MicroRNAs in metabolism and metabolic disorders[J]. *Nat Rev Mol Cell Biol*. 2012;13(4):239–250.
- [21] Huang Y, Yan Y, Xv W, et al. A new insight into the roles of MiRNAs in metabolic syndrome[J]. *Biomed Res Int*. 2018;2018:1–15.
- [22] Zhang H, Feng Z, Huang R, et al. MicroRNA-449 suppresses proliferation of hepatoma cell lines through blockade lipid metabolic pathway related to SIRT1[J]. *Int J Oncol*. 2014;45(5):2143–2152.
- [23] Zhang M, Sun W, Zhou M, et al. MicroRNA-27a regulates hepatic lipid metabolism and alleviates NAFLD via repressing FAS and SCD1[J]. *Sci Rep*. 2017;7(1):14493.
- [24] Nie H, Song C, Wang D, et al. MicroRNA-194 inhibition improves dietary-induced non-alcoholic fatty liver disease in mice through targeting on FXR[J]. *Biochim Biophys Acta (BBA) - Mol Basis Dis*. 2017;1863(12):3087–3094.
- [25] He S, Guo W, Deng F, et al. Targeted delivery of microRNA 146b mimic to hepatocytes by lactosylated PDMAEMA nanoparticles for the treatment of NAFLD[J]. *Artif Cells Nanomed Biotechnol*. 2018;46(sup2):217–228.
- [26] Lakhani HV, Sharma D, Dodrill MW, et al. Phenotypic alteration of hepatocytes in non-alcoholic fatty liver disease[J]. *Int J Med Sci*. 2018;15(14):1591–1599.
- [27] Vidal-Puig A, Jimenez-Li An M, Lowell BB, et al. Regulation of PPAR gamma gene expression by nutrition and obesity in rodents[J]. *J Clin Investig*. 1996;97(11):2553.
- [28] Shimano H, Horton JD, Shimomura I, et al. Isoform 1c of sterol regulatory element binding protein is less active than isoform 1a in livers of transgenic mice and in cultured cells [J]. *J Clin Investig*. 1997;99(5):846–854.
- [29] Koo SH. Nonalcoholic fatty liver disease: molecular mechanisms for the hepatic steatosis[J]. *Clin Mol Hepatol*. 2013;19(3):210–215.
- [30] X D L, X M L, Gu JW, et al. MiR-155 regulates lymphoma cell proliferation and apoptosis through targeting SOCS3/JAK-STAT3 signaling pathway[J]. *Eur Rev Med Pharmacol Sci*. 2017;21(22):5153.
- [31] Shouda T, Yoshida T, Hanada T, et al. Induction of the cytokine signal regulator SOCS3/CIS3 as a therapeutic strategy for treating inflammatory arthritis.[J]. *J Clin Investig*. 2001;108(12):1781–1788.
- [32] Soraya T, MéliSSa R, Bhama R, et al. Loss of SOCS3 expression in T cells reveals a regulatory role for interleukin-17 in atherosclerosis.[J]. *J Exp Med*. 2009;206(10):2067–2077.
- [33] Paul K, Howard JK, Badman MK, et al. Enhanced leptin sensitivity and improved glucose homeostasis in mice lacking suppressor of cytokine signaling-3 in POMC-expressing cells[J]. *Cell Metab*. 2006;4(2):123–132.
- [34] Marina L, Kurkowski MU, Katharina L, et al. Stat3/Socs3 activation by IL-6 transsignaling promotes progression of pancreatic intraepithelial neoplasia and development of pancreatic cancer[J]. *Cancer Cell*. 2011;19(4):441–455.
- [35] Kohjiro U, Tatsuya K, Yu-Hua T, et al. Central role of suppressors of cytokine signaling proteins in hepatic steatosis, insulin resistance, and the metabolic syndrome in the mouse[J]. *Proc Natl Acad Sci U S A*. 2004;101(28):10422–10427
- [36] Pan X, Wang P, Luo J, et al. Adipogenic changes of hepatocytes in a high-fat diet-induced fatty liver mice model and non-alcoholic fatty liver disease patients[J]. *Endocrine*. 2015;48(3):834–847.
- [37] Liu Z, Gan L, Zhou Z, et al. SOCS3 promotes inflammation and apoptosis via inhibiting JAK2/STAT3 signaling pathway in 3T3-L1 adipocyte[J]. *Immunobiology*. 2015;220(8):947–953.