

Response to letter to editor regarding ECEIM consensus statement on equine metabolic syndrome

Dear Editors,

We would like to offer the following explanation and clarification regarding the points raised by Ms McLeod concerning the recent ECEIM consensus statement on equine metabolic syndrome.

Specific guidance on representative figures for diets high or low in non-structural carbohydrate (NSC) is offered in section 7.1 of the article, with figures of <10% and >18% NSC, respectively.

Regarding the classification of carrots and apples as high NSC feed items, it is both logical and customary in animal nutrition to compare feeds based on their dry matter content and as such, carrots and apples have frequently more than half of their dry matter represented by NSC, and invariably far more than 18%. Comparison of feeds "as fed," such as in Ms McLeod's letter, can lead to confusing paradoxes. For example, 100 g of pure glucose would be regarded as a low NSC feed if given along with a liter of water. We are aware that the actual glycaemic load of a single carrot may be small, but we still regard the feeding of even small quantities of high-NSC feeds as suboptimal practice that demonstrates poor discipline of the horse carer as well as offering no nutritional benefit to a well-balanced diet as described in the statement.

Ms McLeod suggests that the article fails to mention forage energy content, although in section 7.1 we recommend total digestible energy intake to be 64%-94% of maintenance requirements. In cases where digestible energy content of forage is known, then a simple calculation can be made based on total dry matter fed to result in the total recommended forage digestible energy intake. We feel that it is self-evident that feeding a forage with lower digestible energy content should follow failure to observe weight loss.

We would also like to clarify the basis for the recommendation that haylage should not be fed to EMS cases. This view is entirely consistent with current best evidence and we did not feel able to speculate on what might or might not be the case with lower NSC haylages whether fed alone or mixed with straw. It remains unclear exactly what the insulinemic components of equine feeds are. Evidence in other species indicates that it is not only glucose that stimulates insulin release, with other important factors including aminoacids, fatty acids, and incretins, for example. The concerning finding in the study referenced was that haylage appeared to have a disproportionately high insulinemic effect compared to hay with a similar NSC content and raises the possibility of further insulinemic factors associated with

haylage.¹ This, along with the generally higher palatability of haylages inevitably, leads to caution against haylage feeding in EMS cases pending any further evidence.

Regarding hay soaking, Ms McLeod's question misrepresents what is actually written in the article. Section 7.1 of the manuscript details the beneficial effect of soaking forage for 7-16 hours. Because of concerns about microbial growth in warm water,² the article then goes on to recommend that in warm conditions the soaking time is limited to 1-2 hours. Although, clearly longer soaking times will have the greatest effect on reducing water soluble carbohydrates (WSC) in forage, soaking hay for as little as 15 minutes has been shown to have a significant effect on WSC in hay.^{3,4}

Andy E. Durham¹

Nicholas Frank²

Cathy M. McGowan³

Nicola J. Menzies-Gow⁴

Ellen Roelfsema⁵

Ingrid Vervuert⁶

Karsten Feige⁷

Kersten Fey⁸

¹Liphook Equine Hospital, United Kingdom

²Department of Clinical Sciences,
Cummings School of Veterinary Medicine at Tufts University,
North Grafton, Massachusetts

³Institute of Veterinary Science,
University of Liverpool,
United Kingdom

⁴Department of clinical sciences and services,
Royal Veterinary College,
Herts, United Kingdom

⁵Utrecht University, Department of Equine Sciences,
Utrecht, The Netherlands

⁶Institute of Animal Nutrition,
Nutrition Diseases and Dietetics,
Faculty of Veterinary Medicine,

University of Leipzig, Leipzig, Germany

⁷Clinic for Horses, University of Veterinary Medicine Hannover,
Hanover, Germany

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2019 The Authors. *Journal of Veterinary Internal Medicine* published by Wiley Periodicals, Inc. on behalf of the American College of Veterinary Internal Medicine.

⁸*Equine Clinic, Internal Medicine, Faculty of Veterinary Medicine,
Justus-Liebig-University of Giessen, Giessen, Germany*

REFERENCES

1. Carslake HB, Argo CM, Pinchbeck GL, Dugdale AHA, McGowan CM. Insulinaemic and glycaemic responses to three forages in ponies. *Vet J.* 2018;235:83-89.
2. Müller CE, Nostell K, Bröjer J. Microbial counts in forages for horses—effect of storage time and of water soaking before feeding. *J Equine Vet.* 2015;35(7):622-627.
3. Warr E, Petch J. Effects of soaking hay on its nutritional quality. *Equine Vet Educ.* 1993;5(3):169-171.
4. Martinson K, Jung H, Hathaway M, Sheaffer C. The effect of soaking on carbohydrate removal and dry matter loss in orchardgrass and alfalfa hays. *J Equine Vet.* 2012;32(6):332-338.