

# Clarithromycin vs. Gemifloxacin in Quadruple Therapy Regimens for Empiric Primary Treatment of *Helicobacter pylori* Infection: A Randomized Clinical Trial

Mohsen Masoodi<sup>1</sup>, Mahshid Talebi-Taher<sup>2\*</sup>, Khadijeh Tabatabaie<sup>3</sup>, Siamak Khaleghi<sup>4</sup>,  
Amir-Hossein Faghihi<sup>1</sup>, Shahram Agah<sup>1</sup>, Reyhaneh Asadi<sup>3</sup>

1. Associate professor, Colorectal Research Center, Rasoul-e-Akram General Hospital, Iran University of Medical Sciences, Tehran, Iran
2. Associate professor, Antimicrobial Resistance Research Center, Infectious Diseases Department, Rasoul-e-Akram General Hospital, Iran University of Medical Sciences, Tehran, Iran
3. Researcher, Rasoul-e-Akram General Hospital, Iran University of Medical Sciences, Tehran, Iran
4. Assistant professor, Colorectal Research Center, Rasoul-e-Akram General Hospital, Iran University of Medical Sciences, Tehran, Iran

**\*Corresponding Author:**

Mahshid Talebi-Taher, MD, MPH  
Sattarkhan St., Niayesh, Rasoul-e-Akram  
General Hospital, Tehran, Iran  
Tel: +98 21 66507056  
Fax: +98 21 66506864  
Email: mtalebitaher2000@yahoo.com  
Received: 28 Dec. 2014  
Accepted: 10 Mar. 2015

## ABSTRACT

### BACKGROUND

Eradication of *Helicobacter pylori* infection plays a crucial role in the treatment of peptic ulcer. Clarithromycin resistance is a major cause of treatment failure. This randomized clinical trial aimed at evaluating the efficacy of a clarithromycin versus gemifloxacin containing quadruple therapy regimen in eradication of *H.pylori* infection.

### METHODS

In this randomized double blind clinical trial (RCT 2012102011054N2), a total of 120 patients were randomized to two groups of 60 patients each. Patients with proven *H.pylori* infection were consecutively assigned into two groups to receive OBAG or OBAC in gastroenterology clinic in Rasoul-e-Akram General Hospital in Tehran, Iran. The patients in the OBAG group received omeprazole (20 mg) twice daily, bismuth subcitrate (240 mg) twice daily, amoxicillin (1 gr) twice daily, and gemifloxacin (320 mg) once daily, and those in the OBAC group received omeprazole (20 mg) twice daily, 240 mg of bismuth subcitrate twice daily, amoxicillin (1 gr) twice daily, and clarithromycin (500 mg) twice daily for 10 days.

### RESULTS

Five patients from each group were excluded from the study because of poor compliance, so 110 patients completed the study. The intention-to-treat eradication rate was 61.6% and 66.6% for the OBAC and OBAG groups, respectively. According to the per protocol analysis, the success rates of eradication of *H.pylori* infection were 67.2% and 72.7% for OBAC and OBAG groups, respectively ( $p=0.568$ ).

### CONCLUSION

The results of this study suggest that gemifloxacin containing regimen is at least as effective as clarithromycin regimen; hence, this new treatment could be considered as an alternative for the patients who cannot tolerate clarithromycin.

### KEYWORDS

*H.pylori*; Gemifloxacin; Clarithromycin; Iran

Please cite this paper as:

Masoodi M, Talebi-Taher M, Tabatabaie K, Khaleghi S, Faghihi AH, Agah S, Asadi R. Clarithromycin vs. Gemifloxacin in Quadruple Therapy Regimens for Empiric Primary Treatment of *Helicobacter pylori* Infection: A Randomized Clinical Trial. *Middle East J Dig Dis* 2015;7:88-93.

## INTRODUCTION

*Helicobacter pylori* infection is not only a major cause of gastritis,

peptic ulcer, gastric cancer, and MALT (mucosa-associated lymphoid tissue) lymphoma but also plays a leading role in creation of iron deficiency anemia, vitamin B12 deficiency, and idiopathic thrombocytopenic purpura.<sup>1</sup>

Eradication rates in the first-line *H.pylori* therapy have been declining over the years, essentially due to increasing resistance against the recommended antibiotics of metronidazole and clarithromycin. Resistance of *H.pylori* to clarithromycin is an important reason for the treatment failure.<sup>2,3</sup>

Fluoroquinolones are active against gram-negative bacteria such as *H.pylori* and have a synergistic effect with proton pump inhibitors (PPIs).<sup>4,5</sup> Some data showed that levofloxacin-based therapy was an effective treatment with 70-80% eradication rate.<sup>6,7</sup> Minehart evaluated the susceptibility of *H.pylori* to fluoroquinolones, and found that gemifloxacin was the most active agent followed by gatifloxacin, ciprofloxacin, levofloxacin, and moxifloxacin.<sup>8</sup>

Some studies in Iran showed that the resistance rate to clarithromycin was in the range of 5% to 45.2%.<sup>9,10</sup> Shokrzadeh and colleagues reported the resistance rates to metronidazole, ciprofloxacin, clarithromycin, and tetracycline as 40.5%, 2.4%, 14.3%, and 4.8%, respectively.<sup>11</sup> This alarming finding indicates an urgent need for introduction of new antibiotics in Iran.

The present study aimed at evaluating the efficacy of a clarithromycin versus gemifloxacin containing quadruple therapy regimen in eradication of *H.pylori* infection.

## MATERIALS AND METHODS

This double blind randomized clinical trial (RCT 2012102011054N2) was conducted in Rasoul-e-Akram Medical University Hospital (Tehran, Iran) during 2012-2013. The participants were chosen from those patients with the complaint of dyspepsia who referred to the Gastroenterology Clinic of the hospital. They were enrolled for this study after giving informed consent.

A total of 120 patients were included to make two groups of 60 patients. The sample size was calculated assuming eradication of *H.pylori* in at least

85% of treated patients, aiming to detect a difference of 10% based on a 0.80 power to detect significant difference ( $p=0.05$ ).

Exclusion criteria were: receiving *H.pylori* treatment, pregnancy, age lower than 18 years, history of chronic renal failure, congestive heart failure, decompensated liver cirrhosis, and gastric surgery.

The presence of *H.pylori* was defined as positive <sup>13</sup>C-urea breath test (UBT), positive pathology and/or rapid urease test (RUT). All biopsy samples were stained with hematoxylin & eosin and giemsa in Pathology Department by a pathologist, who was blinded to the treatment arm. The result of RUT was defined positive if the color of the gel turned pink or red 6 hours after examination. Of the 120 patients, 77 underwent endoscopy and were diagnosed by using RUT and/or histological evaluation (64.1%), and 43 were diagnosed by UBT (35.9%). Endoscopy was performed by the discretion of the managing physician.

A trained interviewer obtained demographic data and filled a standardized questionnaire. The participants were randomly assigned into two groups to receive OBAG or OBAC. The former group received omeprazole (20 mg) twice daily, bismuth subcitrate (240 mg) twice daily, amoxicillin (1 gr) twice daily, and gemifloxacin (320 mg) once daily for 10 days, and the latter group received omeprazole (20 mg) twice daily, bismuth subcitrate (240 mg) twice daily, amoxicillin (1 gr) twice daily, and clarithromycin (500 mg) twice daily for 10 days.

The participants were requested to refer again to the clinic during the second week to evaluate their compliance, as well as adverse drug effects. Compliance was acceptable when over 80% of the total medications were taken. The adverse effects included anorexia, nausea, vomiting, headache, skin rash, and bitter mouth. All the participants underwent UBT 8 weeks after completion of treatment to confirm *H.pylori* eradication.

Gender distribution as well as the efficacy and frequency of side effects in the two groups were compared by Chi-square test. Data analysis was performed on both per protocol (PP) and intention-to-treat (ITT) bases. A  $p$ -value < 0.05 was considered

as statistically significant. Statistical analyses were performed using the SPSS software (version 18; SPSS Inc.).

This study was approved by both the Deputy Research of Iran University of Medical Sciences and Colorectal Research Center at Rasoul-e-Akram General Hospital, Tehran, Iran.

## RESULTS

A total of 120 patients were enrolled in the study, and randomly assigned into OBAG (n=60) or OBAC (n=60) groups. Five patients from each group were excluded from the study because of poor compliance. Both groups had similar compliance according to the number of pills used (OBAG=98.3%, OBAC=97.1%,  $p=0.903$ ).

Finally, 110 patients completed their treatment and follow-up. They had a mean age of  $40.57 \pm 12.74$  years (range, 19-81 years). Fifty four (49.09%) patients were men. ITT and PP analyses showed similar eradication rate in both groups. According to the PP analysis, the success rates of eradication of *H.pylori* infection were 67.2% and 72.7% for OBAC and OBAG groups, respectively ( $p=0.568$ , table 1).

There was significant statistical difference in the eradication rate of *H.pylori* infection according to the pathological findings between the two groups ( $p=0.001$ , table 2). In logistic regression analysis, after adjusting for endoscopic and pathological findings, there was no significant difference between the two groups in response to the treatment ( $p=0.732$ ).

Adverse drug effects were reported in 37 (61.66%) and 19 (31.66%) patients in the clarithromycin and gemifloxacin groups, respectively. Among those, only bitter mouth was significantly more common in the clarithromycin group ( $p=0.001$ , table 3).

## DISCUSSION

Our findings in the present study showed that gemifloxacin is as effective as clarithromycin in eradication of *H.pylori* infection. 5-45% of *H.pylori* isolates from Iran show some degrees of resistance

to clarithromycin.<sup>9,10</sup> To tackle this problem, an alternative drug that is effective, safe, and easy to use has been searched.

The eradication rate has got a decreasing trend by using clarithromycin from 2000 to 2012 in Iran. Fakheri and colleagues showed that the ITT eradication rate was 85% for clarithromycin in quadruple therapy regimens in 2000.<sup>12</sup> However, similar to the present study, Minakari and co-workers found the eradication rates of ITT and PP as 64.5% and 74.7%, respectively, by using amoxicillin, clarithromycin, bismuth and omeprazole as the second-line therapy.<sup>13</sup>

The main risk factor for clarithromycin resistance is previous consumption of macrolides. Although clarithromycin had not been introduced in Iran, some studies has shown 17% resistance rate to this drug. Erythromycin was used instead of it in Iran.<sup>14,15</sup>

Several authors have reported an increasing trend in clarithromycin resistance rates in other areas of the world such as Latin America and Turkey.<sup>16,17</sup> In contrast, there was no change in the trend of *H.pylori* eradication rate by using PPI, clarithromycin and amoxicillin over the recent 11 years in Korea.<sup>18</sup>

Unfortunately, inappropriate use of antibiotics, especially macrolides in Iran has led to the emergence and spread of resistant bacteria, and plans to reduce the over-prescription of antibiotics by physicians should be arranged.

It is important to keep in mind that clarithromycin containing triple therapy is not an appropriate regimen when resistance is between 7% and 10%.<sup>19</sup> So identifying alternative treatments that are more effective than clarithromycin against *H.pylori* is crucial.

Some randomized trials have shown that a levofloxacin containing therapy is more effective compared with a clarithromycin containing regimen.<sup>6,20</sup> There are reports showing that levofloxacin had significant activity against *H.pylori* in vitro but no cross-resistance to B-lactams and macrolides. Moreover, it had a synergistic effect with PPIs.<sup>4-5</sup>

The eradication rate of levofloxacin-containing triple therapy ranged from 72% to 96%. Some

**Table 1: Demographic data and treatment results in patients receiving OBAC and/or OBAG**

Variables	Gemifloxacin	Clarithromycin	p-value
Mean age	41.20±14.00	39.93±11.98	0.646
Male	22 (40%)	34 (58.3%)	0.028
Smoking	6 (13.3%)	6 (13.3%)	0.990
Endoscopy	38(69.09%)	39(70.90%)	0.890
Endoscopy	Non-erosive gastritis	3	0.870
	Erosive gastritis	14	
	Peptic ulcer	21	
Treatment result	ITT <sup>1</sup> (n=60)	40 (66.6%)	0.568
	PP <sup>2</sup> (n=55)	40 (72.7%)	

<sup>1</sup>ITT= intention to treat,  
<sup>2</sup>PP=per protocol

**Table 2: Eradication rate of *H. pylori* infection in the two groups receiving OBAC and/or OBAG treatment**

Endoscopic findings OBAC group	n	Eradication rate(%)	p-value
PUD <sup>1</sup>	23	16 (73.91)	0.001
Erosive gastritis	14	8 (57.14)	
Non-erosive gastritis	2	2 (100)	
<b>Endoscopic findings OBAG group</b>			
PUD <sup>1</sup>	21	11 (52.38)	
Erosive gastritis	14	10 (71.42)	
Non-erosive gastritis	3	1 (33.33)	

<sup>1</sup>Peptic ulcer disease

**Table 3: Adverse effects reported by the patients during the treatment**

Adverse effect	OBAC (n=55)	OBAG (n=55)
Nausea	3	2
Vomiting	1	1
Diarrhea	3	3
Abdominal pain	2	3
Headache	2	2
Skin rash	1	6
Bitter mouth*	25	2

\*p<0.001

authors suggest that this regimen is reasonable in populations with clarithromycin resistance greater than 15-20% and fluoroquinolones resistance less than 10%.<sup>21</sup> But it might not be as the first-line therapy because over-consumption of fluoroquinolones would likely lead to the emergence of more quinolone-resistant pathogens responsible for other organs infections.<sup>19</sup>

A first-line eradication regimen should be based on the prevalence of antimicrobial resistance in Iran. Primary resistance of *H.pylori* isolates in Iran was to clarithromycin (34%), levofloxacin (5.3%), and moxifloxacin (4.6%).<sup>22</sup> So we need more studies to identify the best regimen for *H.pylori* infection therapy.

Gemifloxacin has shown an spectrum of activity against Gram-positive and Gram-negative bacteria. Its mechanism of action focuses on inhibiting DNA gyrase and topoisomerase, thus preventing cellular

replication.<sup>23</sup> Some studies have shown that gemifloxacin is superior to levofloxacin in antimicrobial activity against *H. pylori*, and even overcomes some levofloxacin resistance.<sup>8,24</sup>

Graham suggested that in empiric therapies that did not reliably yield 90% or greater cure, ITT should not be prescribed.<sup>25</sup> It has been reported that standard triple therapy is not a suitable regimen in the areas with clarithromycin resistance over 20% because the PP eradication rate of that regimen is often less than 85%, and the ITT eradication rate is usually less than 80%.<sup>26</sup>

To overcome the problem of the growing clarithromycin resistance in the Middle East, several methods of treatment have been proposed, including the extension of the treatment duration to 14 days, and using bismuth-containing quadruple, sequential, and concomitant treatments.<sup>19</sup> Nasa and colleagues reported that sequential therapy was

better than standard therapy for eradicating *H.pylori* infection.<sup>27</sup> More studies are needed to find the best regimen for eradication of *H.pylori* in Iran.

The results of this study suggest that gemifloxacin containing regimen is at least as effective as clarithromycin regimen; hence, this new treatment could be considered as an alternative choice for the patients who cannot tolerate clarithromycin. Non-optimal result of the two regimens may be due to the emergence of resistant strains of *H.pylori* in Iran. So empirical use of these regimens without susceptibility testing may not be appropriate.

#### ACKNOWLEDGMENT

This study was supported by a grant from the Research Council of Iran University of Medical Sciences, and Colorectal Research Center at Rasoul-e-Akram Hospital (grant No. 18790). We would like to thank the staff in the Division of Gastroenterology at Rasoul-e-Akram General Teaching Hospital, Tehran, Iran.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest related to this work.

#### REFERENCES

- Franceschi F, Zuccala G, Roccarina D, Gasbarrini A. Clinical effects of *Helicobacter pylori* outside the stomach. *Nat Rev Gastroenterol Hepatol* 2014;**11**:234-42.
- Krasz S, Miehlke S, Berning M, Morgner A, Labenz J. [Current value of quinolones in *Helicobacter pylori* therapy]. *Z Gastroenterol* 2011;**49**:989-96.
- Ducons JA, Santolaria S, Guirao R, Ferrero M, Montoro M, Gomollon F. Impact of clarithromycin resistance on the effectiveness of a regimen for *Helicobacter pylori*: A prospective study of 1-week lansoprazole, amoxicillin and clarithromycin in active peptic ulcer. *Aliment Pharmacol Ther* 1999;**13**:775-80.
- Sanchez JE, Saenz NG, Rincon MR, Martin IT, Sanchez EG, Martinez MJ. Susceptibility of *Helicobacter pylori* to mupirocin, oxazolidones, quinupristin/dalfopristin and new quinolones. *J Antimicrob Chemother* 2000;**46**:283-5.
- Tanaka M, Isogai E, Isogai H, Hayashi S, Sugiyama T, Sato K. Synergic effect of quinolone antibacterial agents and proton pump inhibitors on *Helicobacter pylori*. *J Antimicrob Chemother* 2002;**49**:1039-40.
- Kuo CH, Hu HM, Kuo FC, Hsu PI, Chen A, Yu FJ, et al. Efficacy of levofloxacin-based rescue therapy for *Helicobacter pylori* infection after standard triple therapy: A randomized control trial. *J Antimicrob Chemother* 2009;**63**:1017-24.
- Gisbert JP, Fernandez-Bermejo M, Molina-Infante J, Perez-Gallardo B, Prieto-Bermejo AB, Mateos-Rodriguez JM, et al. First-line triple therapy with levofloxacin for *Helicobacter pylori* eradication. *Aliment Pharmacol Ther* 2007;**26**:495-500.
- Minehart HW, Chalker AF. In vitro activity of gemifloxacin against *Helicobacter pylori*. *J Antimicrob Chemother* 2001;**47**:360-1.
- Farshad S, Alborzi A, Japoni A, Ranjbar R, Hosseini ASI K, Badiee P, et al. Antimicrobial susceptibility of *Helicobacter pylori* strains isolated from patients in Shiraz, Southern Iran. *World J Gastroenterol* 2010;**16**:5746-51.
- Abadi AT, Taghavi T, Mobarez AM, Carpenter BM. Frequency of antibiotic resistance in *Helicobacter pylori* strains isolated from the northern population of Iran. *J Microbio* 2011;**49**:987-93.
- Shokrzadeh L, Jafari F, Dabiri H, Baghaei K, Zojaii H, Alizadeh AH, et al. Antibiotic susceptibility profile of *Helicobacter pylori* isolated from the dyspepsia patients in Tehran, Iran. *Saudi J Gastroentero* 2011;**17**:261-4.
- Fakher H, Malekzadeh R, Merat S, Khatibian M, Fazel A, Alizadeh BZ, et al. Clarithromycin vs. furazolidone in quadruple therapy regimens for the treatment of *Helicobacter pylori* in a population with a high metronidazole resistance rate. *Aliment. Pharmacol Ther* 2001;**15**:411-6.
- Minakari M, Davarpanah Jazi AH, Shavakhi A, Moghareabed N, Fatahi F. A randomized controlled trial: Efficacy and safety of azithromycin, ofloxacin, bismuth, and omeprazole compared with amoxicillin, clarithromycin, bismuth, and omeprazole as second-line therapy in patients with *Helicobacter pylori* infection. *Helicobacter* 2010;**15**:154-9.
- Mohammadi M, Doroud D, Massarrat S, Farahvash MJ. Clarithromycin resistance in Iranian *H.pylori* strains before introduction of clarithromycin. *Helicobacter* 2003;**8**:80.
- Cabrita J, Oleastro M, Matos R, Manherte A, Cabral J, Barros R, et al. Features and trends in *Helicobacter pylori* antibiotic resistance in Lisbon area, Portugal (1990-1999). *J Antimicrob Chemother* 2000;**46**:1029-31.
- Camargo MC, Garcia A, Riquelma A, Otero W, Camargo CA, Hernandez-Garcia T, et al. The problem of *Helicobacter pylori* resistance to antibiotics: A systematic review in Latin America. *Am J Gastroenterol* 2014;**109**:485-95.
- Simsek H, Blaban YH, Gunes DD, Hascelik G, Ozarlan E, Tatar G. Alarming clarithromycin resistance of *Helicobacter pylori* in Turkish population. *Helicobacter* 2005;**10**:360-1.
- Yoon JH, Baik GH, Sohn KM, Kim DY, Kim YS, Suk KT, et al. Trends in the eradication rates of *Helicobacter pylori* infection for eleven years. *World J Gastroenterol* 2012;**18**:6628-34.
- Federico A, Gravina AG, Miranda A, Loguercio C, Roma-

- no M. Eradication of *Helicobacter pylori* infection: Which regimen first? *World J Gastroenterol* 2014;**20**:665-72.
20. Romano M, Cuomo A, Gravina AG, Miranda A, Iovene MR, Tilso A, et al. Empirical levofloxacin-containing versus clarithromycin-containing sequential therapy for *Helicobacter pylori* eradication: a randomized trial. *Gut* 2010;**59**:1465-70.
  21. Berning M, Krasz S, Miehle S. Should quinolones come first in *Helicobacter pylori* therapy? *Therap Adv Gastroenterol* 2011;**4**:103-14.
  22. Talebi Bezmin Abadi A, Ghasemzadeh A, Taghvaei T, Mobaraz AM. Primary resistance of *Helicobacter pylori* to levofloxacin and moxifloxacin in Iran. *Intern. Emerg Med* 2012;**7**:447-52.
  23. Le TP, Xiang YQ. Gemifloxacin. *Drugs Today(Barc)* 2001;**37**:401-10.
  24. Chang WL, Kao CY, Wu CT, Huang AH, Wu JJ, Yang HB, et al. Gemifloxacin can partially overcome Quinolone resistance of *H.pylori* with *gyrA* mutation in Taiwan. *Helicobacter* 2012;**17**:210-5.
  25. Graham DY, Shiotani A. New concepts of resistance in the treatment of *Helicobacter pylori* infections. *Nat Clin Pract Gastroenterol Hepatol* 2008;**5**:321-31.
  26. Megraud F. *H.pylori* antibiotic resistance prevalence, importance, and advances in testing. *Gut* 2004;**53**:1374-84.
  27. Nasa M, Choksey A, Phadke A, Sawant P. Sequential therapy versus standard triple-drug therapy for *Helicobacter pylori* eradication: A randomized study. *Indian J Gastroenterol* 2013;**32**:392-6.