

RESEARCH ARTICLE

A SIX1 Homolog in *Fusarium oxysporum* f. sp. *conglutinans* Is Required for Full Virulence on Cabbage

Erfeng Li¹*, Gang Wang¹*, Jiling Xiao²*, Jian Ling¹, Yuhong Yang¹, Bingyan Xie¹*

1 Institute of Vegetables and Flowers, Chinese Academy of Agricultural Sciences, Beijing 100081, China, **2** Hu'nan Watermelon and Muskmelon Institute, Changsha 410125, China

* These authors contributed equally to this work.

* lfcaas@gmail.com



CrossMark
click for updates

OPEN ACCESS

Citation: Li E, Wang G, Xiao J, Ling J, Yang Y, Xie B (2016) A SIX1 Homolog in *Fusarium oxysporum* f. sp. *conglutinans* Is Required for Full Virulence on Cabbage. PLoS ONE 11(3): e0152273. doi:10.1371/journal.pone.0152273

Editor: Sung-Hwan Yun, Soonchunhyang University, REPUBLIC OF KOREA

Received: April 29, 2015

Accepted: March 13, 2016

Published: March 24, 2016

Copyright: © 2016 Li et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was funded by the National Natural Science Foundation of China (31272003), [\[http://www.nsf.gov.cn/\]](http://www.nsf.gov.cn/), to YHY; National Key Technology R&D Program (2012BAD19B06), [\[http://program.most.gov.cn/\]](http://program.most.gov.cn/), to BYX; National Staple Vegetable Industry Technology System Construction Project (CARS-25-B-01), to BYX; Special Fund for Agro-scientific Research in the Public Interest (200903049-04), to BYX. The funders had roles in study design, data collection and analysis, decision to publish and preparation of the manuscript.

Abstract

Fusarium oxysporum is a soil-born fungus that induces wilt and root rot on a variety of plants. *F. oxysporum* f. sp. *conglutinans* (Foc) can cause wilt disease on cabbage. This study showed that a homolog of SIX1 protein in the *Arabidopsis* infecting isolate Fo5176 (Fo5176-SIX1) had four isoforms in the conidia of Foc by proteomic analysis. Thus, we analyzed the roles of protein Foc-SIX1. Gene expression analysis showed that, compared to the expression in mycelia, dramatically altered expression of *Foc-SIX1* could be detected after infecting cabbages, and *Foc-SIX1* was highly expressed in conidia under axenic culture condition. Furthermore, we knocked out the *Foc-SIX1* gene and found that *Foc-ΔSIX1* mutants had significantly reduced virulence compared with wild type isolate, and full virulence was restored by complementation of *Foc-ΔSIX1* mutants with *Foc-SIX1*. Thus, we concluded that SIX1 in Foc was required for full virulence on cabbage. We also complemented *Foc-ΔSIX1* with *SIX1* gene in *F. oxysporum* f. sp. *lycopersici* (Fol) and found *Foc-ΔSIX1::Fol-SIX1* mutants did not affect the virulence of *Foc-ΔSIX1*. The results confirmed that *Fol-SIX1* was not capable of replacing the role of *Foc-SIX1* in Foc on the disease symptom development of cabbage. The roles of *Fol-SIX1* on virulence might rely on host specificity.

Introduction

The soil-born fungus *Fusarium oxysporum* f. sp. *conglutinans* (Foc) causes vascular wilt disease of cabbage. *Fusarium* wilt of cabbage originated in the United States and was first reported in China in 2001 [1].

The interaction between *F. oxysporum* f. sp. *lycopersici* (Fol) and tomato has become a model system for studying the effectors of the pathogen and R proteins from the host [2]. The secreted-in-xylem (SIX) proteins were first reported in the xylem sap of Fol-infected tomato [3]. In total, eleven SIX proteins have been identified in Fol so far, and the functions of most of these proteins remain unknown [2]. Three of these proteins (SIX1/SIX3/SIX4)

Competing Interests: The authors have declared that no competing interests exist.

could interact with the tomato genes (I/I-1/I-2/I-3) to induce resistance [4–6]. Genomic analysis demonstrated that most of *SIX* genes were primarily located on supernumerary pathogenicity chromosomes that could transfer between strains [7]. The small and cysteine-rich SIX1 effector protein was secreted by Fol during colonization of tomato xylem vessels and was required for virulence in a fully susceptible tomato genotype [8]. SIX1 could interact with I-3, which the tomato plants carry to trigger a resistant recognition reaction [6]. *SIX1* was strongly up-regulated during colonization of the host plant [9], and many effector genes were unexpressed or weakly expressed in axenic cultures and up-regulated during infection [10]. Moreover, it was reported that the SIX4 homolog Fo5176-SIX4, with >99% amino acid identity to Fol-SIX4, was required for full virulence in the *Arabidopsis* infecting isolate Fo5176 [11]. Despite the extensive researches performed in Fol, little research has been conducted on the possible role in pathogenicity of *SIX* genes outside Fol.

In the present study we identified four interesting proteins in the conidia of Foc as Fo5176-SIX1 (gi|342888423) during proteomic analysis. Fo5176-SIX1, identified in the isolate Fo5176, shared 74% identity to Fol-SIX1, which was identified as the avirulence protein of I-3-mediated resistance in tomato [5,6,11]. Compared to *SIX1* expression in mycelia, *SIX1* expression was dramatically altered after Foc infected cabbages. In this work, we also found that *Foc-SIX1* expression was also high in conidia under axenic culture condition. Thus, the objective of this study was to explore the roles played by SIX1 in Foc infection of cabbage.

Methods

Fungal strains, plant materials and culture conditions

Cabbage infecting Foc wild-type isolate (52557TM) was obtained from the Beijing Academy of Agriculture and Forestry Sciences and selected for use in this study. Foc wild type isolate was routinely grown on potato dextrose agar (PDA) at 28°C. Deletion mutants (Foc- Δ SIX1: D1 and D2) and complementation mutants (Foc- Δ SIX1::Foc-SIX1: C1, C2, C3 and C4; Foc- Δ SIX1::Fol-SIX1: T1 and T2) were maintained on PDA containing 100 μ g/mL hygromycin B or 100 μ g/mL neomycin under the same conditions.

A Foc-susceptible cabbage cultivar “Zhong gan 21” was used in this study [12]. Seedlings were grown in plastic pots filled with an autoclaved mixture of vermiculite and turf (1:1, v/v) in a glasshouse at approx. 25°C and a 16-h light photoperiod until the 2–3 true leaf stage.

Protein extraction, quantification and 2-DE

Total proteins were extracted from the conidia of Foc by the borax/PVPP/Phe (BPP) protocol [13]. This method combined ammonium sulfate saturated-methanol and BPP to isolate proteins. Protein concentrations were quantified by the Bradford method with bovine serum albumin (BSA) as the standard [14]. Then, 24 cm, pH 4–7 linear gradient IPG strips were rehydrated, and IEF was carried out using the Hoffer IEF100 at 20°C for focusing. After equilibration, the strips were transferred to 12.5% SDS-PAGE to separate proteins in the second dimension (Ettan DALT six, GE Healthcare) [15]. Three biological replicates were performed to ensure protein reproducibility.

Gel staining and image analysis

The gels were visualized by the GAP method, which used CBB-G250, ammonium sulfate, and phosphoric, and scanned. These images were analyzed with Image Master 2D Platinum Software (Version 5.0, GE Healthcare).

MS identification and database search

Proteins were identified by MALDI TOF/TOF MS/MS (Bruker Daltonics, Bremen, Germany) and analyzed with flexAnalysis software (Version 3.2) [16]. Spectra were then analyzed using a MASCOT search against the taxonomy of fungi in the non-redundant NCBI (NCBIInr) database. The MS/MS ion search parameters were conducted as MS/MS tolerance of ± 0.5 Da with allowances for oxidation (M) modifications and for one missed cleavage as well as with fixed modification of cysteine by carboxymethyl (Carbamidomethylation, C). The optimal match was considered according to a higher Mascot score. Proteins with protein score confidence intervals above 95% were considered as confident identifications. Furthermore, we used a BLASTP search to find SIX1 homologs. The protein alignment was performed and the tree was generated by the neighbor-joining method using MEGA 6 with 1000 bootstraps based on alignments of full amino acid sequences of homologs.

Mycelia, conidia, infected root collection and RNA extraction

The isolate was cultured in autoclaved potato dextrose broth (PDB) at 28°C on a shaker at 150 rpm for 48 h. Mycelia were then filtered through four layers of sterilized lens paper and washed with sterile distilled water. The conidia were separated from the conidial suspensions by centrifugation at 5000 g for 15 min at room temperature. The conidia were obtained for virulence studies and protein and RNA extraction. The mycelia were used for RNA extraction. After 7 days post-inoculation (dpi), the infected cabbage seedling roots were collected. For each sample, three independent biological replicates were conducted. RNA extraction, cDNA synthesis and qRT-PCR reactions were performed as described previously [17].

Quantification of *SIX1* gene expression using qRT-PCR

Gene sequences were obtained from the published Fo5176 reference genome sequence and Foc genome sequence [7]. To compare *SIX1* expression levels in mycelia, conidia and *in planta* conditions, two housekeeping genes, elongation factor 1- α (*EF-1 α*) and β -*tubulin*, were used for normalization and calculation of the relative expression values using the $2^{-\Delta\Delta C_t}$ method [9,18]. The β -*tubulin* gene primers were designed from FOXG_06228 in Foc and had been previously verified for use in Fo5176 [11]. The qRT-PCR reactions for each gene were repeated four times. All primers were tested for expected size and specific for the predicted PCR product and were listed in S1 Table.

Deletion and complementation of *SIX1* in *F. oxysporum* f. sp. *conglutinans*

Deletion of the *SIX1* gene was generated using the overlap PCR and split marker gene strategy. The strategy used a DNA mixture containing overlapping truncations of the selectable marker to enable homologous integration during transformation [19–22]. The target gene was replaced with the 1.4 kb fragment that encoded hygromycin B phosphotransferase as a selectable marker. A 1165 bp region upstream from the start codon and a 1211 bp region downstream from the stop codon of the target gene were amplified from genomic DNA of Foc with primers 1-F/(2+5)-R and (8+3)-F/4-R. Marker tail sites were introduced 3' of the upstream fragment and 5' of the downstream fragment. The split replacement gene was amplified from vector PCH-SGFP carrying the *hph* gene [23]. The two split marker transformation components were amplified by two rounds of PCR. These steps are shown in the S1 Fig. The PCR mixtures of the two split marker transformation components were directly used for protoplast-mediated transformation [24]. The correct deletion mutants (Foc- Δ SIX1) were confirmed by PCR and

Southern hybridization. Genomic DNA was digested with *Sal*I, separated in a 0.8% agarose gel, and hybridized with a 308 bp probe. All primers related to gene deletion and deletion mutant confirmation were listed in [S2](#) and [S3](#) Tables.

Complementation of Foc- Δ SIX1 was achieved by transformation with approximately 3.0 kb of *Foc-SIX1* spanning from -1838 bp (promoter relative to start codon) to 1132 bp (ORF and 3' UTR). The resulting amplicon was cloned into *Bam*HI/*Eco*RI-digested KN to produce Foc-SIX1-comp-KN. This construct was transformed into Foc- Δ SIX1 to produce Foc- Δ SIX1::Foc-SIX1 by protoplast-mediated transformation. The transformants were selected on PDA with neomycin and confirmed by PCR amplification. All primers related to gene complementation and transformants confirmation are listed in [S4 Table](#).

Expression of Fol-SIX1, the Foc-SIX1 homolog in *F. oxysporum* f. sp. *lycopersici*, in the deletion mutants Foc- Δ SIX1

To characterize the role of the homolog Fol-SIX1 in Foc, the 1.9 kb of *Fol-SIX1* spanning from -832 bp to 1091 bp off Fol genomic DNA was amplified and cloned into *Bam*HI/*Eco*RI-digested KN to produce Fol-SIX1-com-KN. This construct was transformed into Foc- Δ SIX1 to produce Foc- Δ SIX1::Fol-SIX1 strain by protoplast-mediated transformation. The transformants were selected on PDA with neomycin and confirmed by PCR amplification. All primers related to gene complementation and transformants confirmation are listed in [S5 Table](#).

Transformation procedures mediated by PEG-CaCl₂

Protoplasts were prepared as described previously, and their concentration was adjusted to 1×10^8 protoplasts/mL [25]. The protoplasts were then transferred into Eppendorf tubes at 100 μ L/tube and kept at -80°C for future use. Transformation components were added to 100 μ L of frozen protoplast suspension and mixed gently with a pipette tip. Next, 160 μ L of PEG solution was dropped into the DNA-protoplasts suspension. The protoplast suspension was then mixed with molten regeneration medium (minimal medium) and poured onto regeneration medium. Deletion mutants and complementation transformants were selected and maintained on PDA medium containing 100 μ g/mL hygromycin B or 100 μ g/mL neomycin.

Inoculum preparation and inoculation procedure

Conidial suspension of Foc and their mutants were used as inocula, which was described above. Inoculum concentration was adjusted to 1×10^6 conidia/mL and the root-dip method was used in this study [26]. Cabbage seedlings dipped in sterilized distilled water served as non-inoculated controls. For each isolate, three independent biological replicates were carried out and a total of more than thirty seedlings were inoculated in every replicate.

Disease assessment and statistical analysis

Disease symptoms were assessed according to the disease severity, divided into five levels according to pre-established rating scales [12,26]. The scoring standard of cabbage susceptibility to Foc was evaluated by statistical analysis: index = 0–5, where “0” meant no evidence of disease symptoms; “1” represented only one leaf showed slight or moderate yellowing; “2” indicated that two leaves showing slight or moderate yellowing; “3” meant all leaves showed moderate yellowing or wilt except for interior leaves; “4” meant all leaves showed severe yellowing or wilt; and “5” indicated that the seedling was dead. The disease index (DI) was calculated according to the rating scales [12]. The statistical analysis of differences in DI was performed

by two-way ANOVA procedures using Duncan's multiple range test in SAS software. The statistical results are shown as the mean \pm SE (standard error).

Results and Discussion

A conserved SIX1 homolog was identified in Foc by proteomic analysis

Two-dimensional electrophoresis (2-DE) and mass spectrometry (MS), combined with careful systematic and statistical analysis, were used to reliably separate and identify pathogenesis-related proteins in Foc. In total, 461 protein spots were detected in the conidia of Foc. Among these, 20 protein spots were identified. From the identification results, it was noteworthy that four protein spots (1, 2, 3 and 4) had higher identities with Fo5176-SIX1 (Fig 1A and 1B). The Fo5176 genomic sequence was previously obtained using Illumina and 454 sequencing [7,11]. These four proteins, which were nicely separated on the gels, were located in horizontal strings with identical functional annotation. We have sequenced the genome of Foc. Only one homolog of Fo5176-SIX1 was found in Foc genome using BLASTN with E-value cutoff of $1e-5$. The gene *Foc-SIX1* had 100% identity with *Fo5176-SIX1*. This result indicated that a single gene product with post-translational modifications changing the *pI* resulted in four isoforms of this protein in Foc (Fig 1B). We named this protein Foc-SIX1. These four identified protein isoforms were listed in Table 1 (See S6 Table, S1 File). BLASTN searches did not reveal the identification of this protein with known functions, but uncovered its homologs in other *F. oxysporum* formae speciales and *Colletotrichum orbiculare*. Phylogenetic analysis showed that Foc-SIX1 was more similar to Fo5176-SIX1 and Fol-SIX1 proteins than to SIX1 of *F. oxysporum* f. sp. *cubeense* (Fig 2). Foc-SIX1 shared 100% and 74% identity with Fo5176-SIX1 and Fol-SIX1, respectively. These two identified proteins were annotated as small, cysteine-rich proteins and contributed to virulence of the pathogens. Therefore, we have focused on Foc-SIX1 in this study.

Comparative analysis of *SIX1* gene expression in mycelia, conidia and infected cabbage roots

To determine whether *Foc-SIX1* is expressed during infection of cabbage, we quantified its mRNA expression using qRT-PCR in mycelia under axenic culture condition and in infected cabbages. The expression of *SIX1* in mycelia was significantly lower than that in infected cabbage. Its expression was detected in cabbage roots at 90-fold higher levels than when measured




Fig 1. Reference maps from 2-DE analysis of conidial proteins from *F. oxysporum* f. sp. *conglutinans*. Proteins were separated by using 24 cm, pH 4–7 IPG strips for IEF and followed by 12.5% SDS-PAGE. The gels were stained by GAP method, which used CBB-G250, ammonium sulfate, and phosphoric acid. A: Gel of conidia proteins from Foc grown in PDB. B: The four proteins (spots 1, 2, 3 and 4), named Foc-SIX1 with high abundance in Foc, were indicated by arrows and numbers and listed in Table 1; Figure B is the partial enlarged details of the rectangular box in Figure A.

doi:10.1371/journal.pone.0152273.g001

Table 1. Identification of the four proteins indicated in Fig 1 from the conidia of Foc.

Spot No. ^a	Acc. ID ^b	Protein identity	Organism	Theor. PI/Mr ^c	Exper. PI/Mr ^d	MP ^e	SC% ^f	Loc ^g	MS ^h
1	342888423	Fo5176-SIX1	<i>F. oxysporum</i>	8.54/31.3	6.38/16.8	1	5	S	86
2	342888423	Fo5176-SIX1	<i>F. oxysporum</i>	8.54/31.3	6.87/16.9	1	5	S	58
3	342888423	Fo5176-SIX1	<i>F. oxysporum</i>	8.54/31.3	7.01/15.9	2	10	S	182
4	342888423	Fo5176-SIX1	<i>F. oxysporum</i>	8.54/31.3	7.18/15.8	1	5	S	65

^aDesignated spot number as indicated and marked on the 2-DE gel in Fig 1;

^bThe accession the accession numbers of NCBI nr database;

^cTheoretical mass (kDa) and *pI* of identified proteins from the database;

^dExperimental mass (kDa) and *pI* of identified proteins. Experimental values were calculated with Image Master 2D Platinum software and standard molecular mass marker;

^eMatched number of peptides (MP) identified from PFF data;

^fThe amino acid sequence coverage (SC) for the identified proteins;

^gThe protein location (Loc) predicted by Target P. (S: secretory pathway);

^hThe Mascot searched score (MS) against the database NCBI nr.

doi:10.1371/journal.pone.0152273.t001

in mycelia (Fig 3). These expression levels indicated that dramatically altered expression of *SIX1* could be detected in Foc after cabbage infection. Previous research reported that many effector genes encoding small secreted proteins were low or undetectable under axenic culture conditions and were up-regulated specifically during infection [10]. Van Der Does indicated that *SIX1* gene expression in Fol was much higher *in planta* compared to expression in axenic cultures [9]. This situation also occurred in Foc: *SIX1* in Foc was weakly expressed in mycelia in axenic culture and up-regulated in *planta*, which were consistent with previous investigations. To verify the results based on proteomic analysis, the expression of *SIX1* in conidia was also quantified. As expected, *SIX1* was highly expressed in conidia under axenic culture condition (Fig 3). That was novel compared to the previous researches. The previous researches mainly focused on the expression of *SIX* gene in mycelia under axenic culture and *in planta* conditions.




Fig 2. Phylogenetic analysis of Foc-SIX1 by using neighbor-joining method. The tree was generated based on alignments of full amino acid sequences of homologs using MEGA 6 with 1000 bootstraps.

doi:10.1371/journal.pone.0152273.g002




Fig 3. The expression of *SIX1* gene in mycelia, conidia and infected cabbage roots at 7 dpi. Mycelia and conidia were obtained from *Foc* grown in PDB for 48 h. Gene expression levels were relative to the internal controls *EF-1α* and *β-tubulin*. Averages of four biological replicates were shown with the standard error.

doi:10.1371/journal.pone.0152273.g003

Deletion of *SIX1* gene in *Foc*

The deletion mutants *Foc-ΔSIX1* were obtained by a targeted gene replacement through homologous recombination. The upstream and downstream regions of the target gene and the selectable marker gene were amplified as shown in [S2 Fig](#) (A, B, C, and D). The two split marker transformation components E and F in [S2 Fig](#) were obtained by fusion PCR with A and C, B and D. Each transformation component was flanked by a target gene and contained two-thirds of a selectable marker gene. Homologous recombination between the overlapping regions of the selectable marker gene and between the flank regions and their genome counterparts resulted in a target gene deletion and replacement with an intact selectable marker gene ([Fig 4A](#)). The mutants *Foc-ΔSIX1* were analyzed by PCR to confirm the correct deletion ([Fig 4B](#)). All mutants showed that the *SIX1* gene was deleted and that the *hph* gene was integrated into the correct position within the genome. Furthermore, the mutants were also analyzed by Southern hybridization ([Fig 4C](#)). *Foc-ΔSIX1* showed a larger band compared with wild type isolate. The deletion mutants lacked the 3006 bp band and instead had a 5295 bp band ([Fig 4A and 4C](#)).

Mutants *Foc-ΔSIX1* reduced virulence in *F. oxysporum* f. sp. *conglutinans* on cabbage

To examine the role of *Foc-ΔSIX1* in cabbage infection, we compared the disease symptoms of cabbage when inoculated with wild type isolate and *Foc-ΔSIX1* mutants. *Foc-ΔSIX1*-inoculated cabbage seedlings showed significantly reduced disease symptom development ([Fig 5](#)). For example, the yellowing leaves appeared on cabbages inoculated with wild type isolate at 8 dpi, while there were no obvious symptoms on the cabbages inoculated with *Foc-ΔSIX1* ([Fig 5A](#)). At 10 dpi, severe chlorotic and necrotic symptoms were visualized on the cabbages inoculated with wild type isolate, but the *Foc-ΔSIX1*-inoculated cabbages still grew well with few visible disease symptoms. Even when the seedlings inoculated with wild type isolate *Foc* died at 11 dpi, the seedlings inoculated with *Foc-ΔSIX1* were still healthy. Note that cabbage seedlings inoculated with wild type isolate *Foc* not only showed yellowing and wilting but also had limited growth. However, the seedlings inoculated with *Foc-ΔSIX1* grew healthily. By comparing the disease indexes between wild type isolate (*Foc*) and mutants (*Foc-ΔSIX1*), we concluded that *Foc-ΔSIX1* had significantly reduced virulence on cabbage ([Fig 5B](#), [S7 Table](#)). The results demonstrated that




Fig 4. Construction and verification of deletion mutants Foc- Δ SIX1. A: *Foc-SIX1* gene deletion and replacement with an intact selectable marker gene (*hph*) by homologous recombination. B: Confirmation of the correct deletion by PCR. Numbers 1, 2, 3 and 4 represent the four primer pairs that specify the *SIX1* gene, *hph* gene, and the correct upstream and downstream homologous recombination, respectively. C: Southern hybridization analysis of wild type isolate and mutants Foc- Δ SIX1. The genomic DNAs from the wild type Foc isolate and deletion mutants Foc- Δ SIX1 were digested with *Sal* I. A fragment amplified from upstream of the target gene was used as probe.

doi:10.1371/journal.pone.0152273.g004




Fig 5. Functional analysis of the SIX1 gene in Foc. A: Disease symptoms of cabbage seedlings inoculated with wild type isolate Foc and their mutants Foc-ΔSIX1 were shown at 8 dpi, 10 dpi and 11 dpi. B: Disease index of cabbage seedlings inoculated with wild type isolate Foc and mutants Foc-ΔSIX1 from 8 to 13 days after inoculation. The average of three replicates was shown with standard error of the mean. Each value in the figure was an average of three independent biological replicates. Each replicate included more than 30 seedlings for each isolate.

doi:10.1371/journal.pone.0152273.g005

SIX1 played crucial roles in Foc, which could influence its virulence on cabbage. The growth rates did not differ between wild type isolate (Foc) and mutants (Foc-ΔSIX1) (S3 Fig), suggesting that Foc-SIX1 did not affect fungal growth or development. It was reported that the resistance of cabbage to *Fusarium* wilt was controlled by a single dominant gene *FOCI* [27,28]. SIX1 in *Fol* acted as an Avr protein where it triggered resistance in I-3-containing tomato [6]. In our work, we also analyzed if SIX1 in Foc works as an avirulence gene by the inoculation test using yellow resistance cabbage varieties (“Zhenqi”). The results showed the inoculated seedlings were no disease symptoms even at 15 dpi, whether in Foc-cabbage or Foc-ΔSIX1-cabbage interactions. It indicated that SIX1 was not an avirulence gene, but a virulence gene in Foc.




Fig 6. Construction, verification, and virulence analysis of complementation mutants *Foc-ΔSIX1::Foc-SIX1*. A: *Foc-ΔSIX1* mutants were complemented by transformation with approximately 3.0 kb of *Foc-SIX1* spanning from -1838 bp (promoter relative to start codon) to 1132 bp (ORF and 3' UTR). B: The *SIX1* gene, *hph* gene and *Neo* gene fragments were amplified by PCR to confirm the correct complementation mutants. C: Disease symptoms of cabbage seedlings inoculated with wild type isolate *Foc*, deletion mutants *Foc-ΔSIX1*, and complementation mutants *Foc-ΔSIX1::Foc-SIX1* were shown at 14 dpi. Similar results were obtained in three independent biological replicates.

doi:10.1371/journal.pone.0152273.g006

Restoration of full virulence by complementation of *Foc-SIX1* in the deletion mutants *Foc-ΔSIX1*

To confirm that the reduced virulence of mutants *Foc-ΔSIX1* was due to deletion of *SIX1*, deletion mutants *Foc-ΔSIX1* were complemented with an intact copy of *Foc-SIX1* fragment (Fig 6A). Complementation of *Foc-ΔSIX1* was confirmed by PCR amplification (Fig 6B). *Foc-SIX1*-complemented isolates (C1-C4: *Foc-ΔSIX1::Foc-SIX1*) showed that *SIX1* gene was complemented, and *hph* gene and *Neo* gene were integrated into the genome. After inoculation, all of *Foc-SIX1*-complemented isolates showed full virulence symptom (Fig 6C). The disease indexes for the complemented isolates were not significantly different from that of wild type isolate (*Foc*) and were significantly higher than that of the two deletion mutants (D1 and D2) (S8 Table). After complementation, *Foc-SIX1*-complemented isolates restored full *Foc* virulence on cabbage. It demonstrated that *SIX1* gene was required for full virulence of *Foc* on cabbage.

No complementation of Foc- Δ SIX1 by a Fol-SIX1, the Foc-SIX1 homolog in *F. oxysporum* f. sp. *lycopersici*

To test whether SIX1 might have a nonspecific effect on virulence, we also introduced Fol-SIX1, which was a Foc-SIX1 homolog in Fol, into Foc- Δ SIX1 mutants (Fig 7A). Foc- Δ SIX1::Fol-SIX1 was confirmed by PCR amplification (Fig 7B). The results showed no difference in disease symptom development between Foc- Δ SIX1 and Foc- Δ SIX1::Fol-SIX1 strains when infected cabbage (Fig 7C). Disease development caused by both wild type strain Foc and complemented mutants Foc- Δ SIX1::Foc-SIX1 was significantly higher than that caused by the two




Fig 7. Construction, verification, and virulence analysis of transformants Foc- Δ SIX1::Fol-SIX1. A: Foc- Δ SIX1 mutants were complemented by transformation with approximately 2.0 kb of *Fol-SIX1* spanning from -832 bp (promoter relative to start codon) to 1091 bp (ORF and 3' UTR). B: The *Fol-SIX1* gene and *Neo* gene fragments were amplified by PCR to confirm the correct complementation mutants. C: Disease symptoms of cabbage seedlings inoculated with wild type isolate Foc, deletion mutants Foc- Δ SIX1 (D1 and D2), complementation mutants (C2 and C3) and transformants Foc- Δ SIX1::Fol-SIX1 (T1 and T2) were shown at 18 dpi. Similar results were obtained in three independent biological replicates.

doi:10.1371/journal.pone.0152273.g007

Foc- Δ SIX1 mutants and the two Foc- Δ SIX1::Fol-SIX1 mutants. The disease indexes for complemented isolates (T1 and T2: transformants Foc- Δ SIX1::Fol-SIX1) were not significantly different from those of Foc- Δ SIX1 mutants and were significantly lower than that of wild type isolate (Foc) and the two complemented mutants (C2 and C3: Foc- Δ SIX1::Fol-SIX1) (S9 Table). These experiments confirmed that Fol-SIX1 was not capable of replacing the role of Foc-SIX1 in Foc on the disease symptom development of cabbage. It had been reported that Fo5176-SIX4 and Fol-SIX4 could restore the virulence of Fo5176- Δ SIX4. The expression of Fo5176-SIX4 in *A. thaliana* increased susceptibility to Fo5176 [11]. While Fol-SIX6 did not promote growth of *A. tumefaciens* in *N. benthamiana* [29]. The Fol-SIX6 expression in *A. thaliana* could not enhance susceptibility to Fo5176 or *V. dahliae* [29]. Our research confirmed that Fol-SIX1 did not alter the virulence of Foc- Δ SIX1 isolate on cabbage. The roles of Fol-SIX1 on virulence might rely on the host specificity. The roles of Foc-SIX1 has led us to study its molecular mechanism in further detail. Further research is required to determine the roles played by SIX1 in the conidia of Foc and to notably decrease Foc virulence on cabbage.

Supporting Information

S1 Fig. Overlap PCR and split marker gene disruption strategy. A. Amplification of 5'- and 3'- flanking regions of the target gene and split replacement of selectable marker gene. B. Fusion the flanking regions and split marker sequences. C. Homologous recombination between the flank regions of the target gene and their genome counterparts and between the overlapping regions of the selectable marker gene.
(DOCX)

S2 Fig. The construction of transformation component. A and B. The upstream and downstream homologous recombination fragments of target gene. C and D. The split replacements of *hph* gene. E and F. The fusion fragments between A and C, B and D, which were directly used for protoplast-mediated transformation. M. Marker.
(DOCX)

S3 Fig. The analysis of fungal growth rates between wild type isolate Foc and deletion mutants Foc- Δ SIX1.
(DOCX)

S1 File. Spectrum and MALDI TOF/TOF MS/MS identification information for the identified protein Foc-SIX1.
(DOCX)

S1 Table. The primer pairs used in qRT-PCR analysis.
(DOCX)

S2 Table. The primer pairs for deletion construction of *SIX1* gene.
(DOCX)

S3 Table. The primer pairs used to ensure the correct deletion mutants.
(DOCX)

S4 Table. The primer pairs used for complementation *Foc-SIX1* in Foc- Δ SIX1.
(DOCX)

S5 Table. The primer pairs used for complementation homolog *Fol-SIX1* in Foc- Δ SIX1.
(DOCX)

S6 Table. Detailed information on MS identification, functional classification and GO analysis of the four protein spots.

(XLSX)

S7 Table. Disease index on cabbage seedlings inoculated with wild type isolate Foc and mutants Foc-ΔSIX1 at 8, 10 and 12 dpi.

(DOCX)

S8 Table. Disease index on cabbage seedlings inoculated with wild type Foc, deletion mutants Foc-ΔSIX1 (D1 and D2) and complementation mutants Foc-ΔSIX1::Foc-SIX1 (C1, C2, C3 and C4) at 14 dpi.

(DOCX)

S9 Table. Disease index on cabbage seedlings inoculated with wild type Foc, Foc-ΔSIX1 (D1 and D2), Foc-ΔSIX1::Foc-SIX1 (C2 and C3) and Foc-ΔSIX1::Fol-SIX1 (T1 and T2) at 18 dpi.

(DOCX)

Acknowledgments

The work in the ministry of agriculture key laboratory of genetic improvement of horticultural crops was and is supported in part by grants from the National Natural Science Foundation of China (No.31272003), the National Key Technology R&D Program (No.2012BAD19B06), the Special Fund for Agro-scientific Research in the Public Interest (No.200903049-04) and the National Staple Vegetable Industry Technology System Construction Project (No.CARS-25-B-01). The authors declared no conflicts of interest.

Author Contributions

Conceived and designed the experiments: BYX. Performed the experiments: EFL J LX GW. Analyzed the data: EFL GW JL. Contributed reagents/materials/analysis tools: J LX YHY. Wrote the paper: EFL.

References

1. Smith E (1899) The fungous infestation of agricultural soils in the United States. *Sci Am Suppl* 48: 1981–1982.
2. Takken F, Rep M (2010) The arms race between tomato and *Fusarium oxysporum*. *Mol Plant Pathol* 11: 309–314. doi: [10.1111/j.1364-3703.2009.00605.x](https://doi.org/10.1111/j.1364-3703.2009.00605.x) PMID: [20447279](https://pubmed.ncbi.nlm.nih.gov/20447279/)
3. Rep M, Dekker HL, Vossen JH, De Boer AD, Houterman PM, Speijer D, et al. (2002) Mass spectrometric identification of isoforms of PR proteins in xylem sap of fungus-infected tomato. *Plant Physiol* 130: 904–917. doi: [10.1104/pp.007427](https://doi.org/10.1104/pp.007427) PMID: [12376655](https://pubmed.ncbi.nlm.nih.gov/12376655/)
4. Houterman PM, Cornelissen BJ, Rep M (2008) Suppression of plant resistance gene-based immunity by a fungal effector. *PLoS Pathog* 4: e1000061. doi: [10.1371/journal.ppat.1000061](https://doi.org/10.1371/journal.ppat.1000061) PMID: [18464895](https://pubmed.ncbi.nlm.nih.gov/18464895/)
5. Houterman PM, Ma L, Van Ooijen G, De Vroomen MJ, Cornelissen BJ, Takken FL, et al. (2009) The effector protein Avr2 of the xylem-colonizing fungus *Fusarium oxysporum* activates the tomato resistance protein I-2 intracellularly. *Plant J* 58: 970–978. doi: [10.1111/j.1365-3113.2009.03838.x](https://doi.org/10.1111/j.1365-3113.2009.03838.x) PMID: [19228334](https://pubmed.ncbi.nlm.nih.gov/19228334/)
6. Rep M, Van Der Does HC, Meijer M, Van Wijk R, Houterman PM, Dekker HL, et al. (2004) A small, cysteine-rich protein secreted by *Fusarium oxysporum* during colonization of xylem vessels is required for I-3-mediated resistance in tomato. *Mol Microbiol* 53: 1373–1383. doi: [10.1111/j.1365-2958.2004.04177.x](https://doi.org/10.1111/j.1365-2958.2004.04177.x) PMID: [15387816](https://pubmed.ncbi.nlm.nih.gov/15387816/)
7. Ma LJ, Van Der Does HC, Borkovich KA, Coleman JJ, Daboussi MJ, Di Pietro A, et al. (2010) Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* 464: 367–373. doi: [10.1038/nature08850](https://doi.org/10.1038/nature08850) PMID: [20237561](https://pubmed.ncbi.nlm.nih.gov/20237561/)

8. Rep M, Meijer M, Houterman P, Van Der Does H, Cornelissen B (2005) *Fusarium oxysporum* evades I-3-mediated resistance without altering the matching avirulence gene. *Mol Plant Microbe In* 18: 15–23. doi: [10.1094/MPMI-18-0015](https://doi.org/10.1094/MPMI-18-0015)
9. Van Der Does HC, Duyvesteijn RG, Goltstein PM, Van Schie CC, Manders EM, Cornelissen BJ, et al. (2008) Expression of effector gene *SIX1* of *Fusarium oxysporum* requires living plant cells. *Fungal Genet Biol* 45: 1257–1264. doi: [10.1016/j.fgb.2008.06.002](https://doi.org/10.1016/j.fgb.2008.06.002) PMID: [18606236](https://pubmed.ncbi.nlm.nih.gov/18606236/)
10. Rep M (2005) Small proteins of plant-pathogenic fungi secreted during host colonization. *FEMS Microbiol Lett* 253: 19–27. doi: [10.1016/j.femsle.2005.09.014](https://doi.org/10.1016/j.femsle.2005.09.014) PMID: [16216445](https://pubmed.ncbi.nlm.nih.gov/16216445/)
11. Thatcher LF, Gardiner DM, Kazan K, Manners JM (2012) A highly conserved effector in *Fusarium oxysporum* is required for full virulence on *Arabidopsis*. *Mol Plant Microbe In* 25: 180–190. doi: [10.1094/MPMI-08-11-0212](https://doi.org/10.1094/MPMI-08-11-0212)
12. Li E, Wang G, Yang Y, Xiao J, Mao Z, Xie B (2015) Microscopic analysis of the compatible and incompatible interactions between *Fusarium oxysporum* f. sp. *conglutinans* and cabbage. *Eur J Plant Pathol* 141: 597–609. doi: [10.1007/s10658-014-0567-6](https://doi.org/10.1007/s10658-014-0567-6)
13. Wang X, Li X, Deng X, Han H, Shi W, Li Y (2007) A protein extraction method compatible with proteomic analysis for the eukaryote *Salicornia europaea*. *Electrophoresis* 28: 3976–3987. doi: [10.1002/elps.200600805](https://doi.org/10.1002/elps.200600805) PMID: [17960840](https://pubmed.ncbi.nlm.nih.gov/17960840/)
14. Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248–254. doi: [10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3) PMID: [942051](https://pubmed.ncbi.nlm.nih.gov/942051/)
15. Li E, Ling J, Wang G, Xiao J, Yang Y, Mao Z, et al. (2015) Comparative proteomics analyses of two races of *Fusarium oxysporum* f. sp. *conglutinans* that differ in pathogenicity. *Sci Rep* 5. doi: [10.1038/srep13663](https://doi.org/10.1038/srep13663)
16. Shen S, Jing Y, Kuang T (2003) Proteomics approach to identify wound-response related proteins from rice leaf sheath. *Proteomics* 3: 527–535. doi: [10.1002/pmic.200390066](https://doi.org/10.1002/pmic.200390066) PMID: [12687619](https://pubmed.ncbi.nlm.nih.gov/12687619/)
17. Pu X, Xie B, Li P, Mao Z, Ling J, Shen H, et al. (2014) Analysis of the defence-related mechanism in cucumber seedlings in relation to root colonization by nonpathogenic *Fusarium oxysporum* CS-20. *FEMS Microbiol Lett* 355: 142–151. doi: [10.1111/1574-6968.12461](https://doi.org/10.1111/1574-6968.12461) PMID: [24810367](https://pubmed.ncbi.nlm.nih.gov/24810367/)
18. Beaubois E, Girard S, Lallechere S, Davies E, Paladian F, Bonnet P, et al. (2007) Intercellular communication in plants: evidence for two rapidly transmitted systemic signals generated in response to electromagnetic field stimulation in tomato. *Plant Cell Environ* 30: 834–844. doi: [10.1111/j.1365-3040.2007.01669.x](https://doi.org/10.1111/j.1365-3040.2007.01669.x) PMID: [17547655](https://pubmed.ncbi.nlm.nih.gov/17547655/)
19. Fu J, Hettler E, Wickes B (2006) Split marker transformation increases homologous integration frequency in *Cryptococcus neoformans*. *Fungal Genet Biol* 43: 200–212. doi: [10.1016/j.fgb.2005.09.007](https://doi.org/10.1016/j.fgb.2005.09.007) PMID: [16497523](https://pubmed.ncbi.nlm.nih.gov/16497523/)
20. Son H, Seo YS, Min K, Park AR, Lee J, Jin J, et al. (2011) A phenome-based functional analysis of transcription factors in the cereal head blight fungus, *Fusarium graminearum*. *PLoS Pathog* 7: e1002310. doi: [10.1371/journal.ppat.1002310](https://doi.org/10.1371/journal.ppat.1002310) PMID: [22028654](https://pubmed.ncbi.nlm.nih.gov/22028654/)
21. Yu JH, Hamari Z, Han KH, Seo JA, Reyes Domínguez Y, Scazzocchio C (2004) Double-joint PCR: a PCR-based molecular tool for gene manipulations in filamentous fungi. *Fungal Genet Biol* 41: 973–981. doi: [10.1016/j.fgb.2004.08.001](https://doi.org/10.1016/j.fgb.2004.08.001) PMID: [15465386](https://pubmed.ncbi.nlm.nih.gov/15465386/)
22. Catlett N, Lee BN, Yoder O, Turgeon BG (2003) Split-marker recombination for efficient targeted deletion of fungal genes. *Fungal Genet Newsl*: 9–11.
23. Li E, Wang D, Yang Y, Xie B (2011) *Agrobacterium tumefaciens* mediated transformation of cabbage *Fusarium* wilt pathogen. *China Vegetables* 14: 008.
24. Turgeon BG, Garber RC, Yoder O (1987) Development of a fungal transformation system based on selection of sequences with promoter activity. *Mol Cell Biol* 7: 3297–3305. doi: [10.1128/MCB.7.9.3297](https://doi.org/10.1128/MCB.7.9.3297) PMID: [2823126](https://pubmed.ncbi.nlm.nih.gov/2823126/)
25. Talbot NJ, Ebole DJ, Hamer JE (1993) Identification and characterization of *MPG1*, a gene involved in pathogenicity from the rice blast fungus *Magnaporthe grisea*. *Plant Cell* 5: 1575–1590. doi: [10.1105/tpc.5.11.1575](https://doi.org/10.1105/tpc.5.11.1575) PMID: [8312740](https://pubmed.ncbi.nlm.nih.gov/8312740/)
26. Lv H, Fang Z, Yang L, Xie B, Liu Y, Zhuang M, et al. (2011) Research on screening of resistant resources to *Fusarium* wilt and inheritance of the resistant gene in cabbage. *Acta Horti Sin* 5: 011.
27. Shimizu M, Pu ZJ, Kawanabe T, Kitashiba H, Matsumoto S, Ebe Y, et al. (2014) Map-based cloning of a candidate gene conferring *Fusarium* yellows resistance in *Brassica oleracea*. *Theor Appl Genet* 128: 119–130. doi: [10.1007/s00122-014-2416-6](https://doi.org/10.1007/s00122-014-2416-6) PMID: [25351523](https://pubmed.ncbi.nlm.nih.gov/25351523/)

28. Lv H, Fang Z, Yang L, Zhang Y, Wang Q, Liu Y, et al. (2014) Mapping and analysis of a novel candidate *Fusarium* wilt resistance gene *FOC1* in *Brassica oleracea*. *BMC genomics* 15: 1094. doi: [10.1186/1471-2164-15-1094](https://doi.org/10.1186/1471-2164-15-1094) PMID: [25495687](https://pubmed.ncbi.nlm.nih.gov/25495687/)
29. Gawehns F, Houterman PM, Ichou FA, Michielse CB, Hijdra M, Cornelissen BJ, et al. (2014) The *Fusarium oxysporum* effector Six6 contributes to virulence and suppresses I-2-mediated cell death. *Mol Plant Microbe In* 27: 336–348. doi: [10.1094/MPMI-11-13-0330-R](https://doi.org/10.1094/MPMI-11-13-0330-R)