

SCIENTIFIC REPORTS



OPEN

Rainfall reduction amplifies the stimulatory effect of nitrogen addition on N₂O emissions from a temperate forest soil

Shicong Geng^{1,2}, Zhijie Chen^{1,2}, Shijie Han¹, Fang Wang^{1,2} & Junhui Zhang¹

Received: 29 June 2016

Accepted: 25 January 2017

Published: 24 February 2017

Soil is a significant source of atmospheric N₂O, and soil N₂O emissions at a global scale are greatly affected by environment changes that include continuous deposition of atmospheric nitrogen and changing precipitation distribution. However, to date, field simulations of multiple factors that control the interaction between nitrogen deposition and precipitation on forest soil N₂O emissions are scarce. In this study, we conducted a 2-year continuous assessment of N₂O emissions from November 2012 to October 2014 at a nitrogen addition and rainfall reduction manipulation platform in an old broad-leaved Korean pine mixed forest at Changbai Mountain in northeastern China. We found that N₂O emissions from control plots were 1.25 ± 0.22 kg N₂O-N ha⁻¹ a⁻¹. Nitrogen addition significantly increased N₂O emissions, with the emission factor of 1.59%. A 30% reduction in rainfall decreased N₂O emissions by 17–45%. However, in combination, nitrogen addition and rainfall reduction increased N₂O emissions by 58–140%, with the emission factor of 3.19%, and had a larger promotional effect than the addition of nitrogen alone. Our results indicated that drought slightly decreases forest soil N₂O emission; however, with increasing deposition of atmospheric N in temperate forest soils, the effect of drought might become altered to increase N₂O emission.

As a component of the nitrogen cycle, N₂O has attracted significant attention because N₂O (1) is a long-lived greenhouse gas with an atmospheric lifetime of 114 years and a 100-year global warming potential of 298 relative to CO₂¹, (2) acts as a catalyst to deplete stratospheric ozone², and (3) represents a form of nitrogen loss³. The atmospheric concentration of N₂O has increased linearly by approximately 0.26% yr⁻¹ over the past several decades and reached 319 ppbv in 2005¹. Soils are the primary source of N₂O emissions and contribute approximately 65% of the total N₂O emissions to the atmosphere⁴. Soil microbial processes, including nitrification, denitrification and nitrifier denitrification, lead to the release of N₂O from soils^{5,6}, and these processes are all affected by soil temperature, soil moisture, substrate N availability, dissolved organic carbon (DOC) and soil pH^{7,8}.

Changes in the global environment are anticipated and include continuous atmospheric nitrogen deposition⁹ and redistribution of precipitation¹⁰. These changes will likely strongly affect soil factors and therefore, alter N₂O emissions from soil. Forest soils are an important source of atmospheric N₂O^{11,12}, and nitrogen addition experiments have been conducted to evaluate the effect of enhanced nitrogen deposition on N₂O emissions in many forests^{13,14}. Reductions in rainfall also reportedly affect forest soil N₂O emissions^{15–17}, but these results are not consistent. Some studies find that exclusion of throughfall, by decreasing soil moisture, reduces N₂O emissions^{17,18}, whereas other studies show that reductions in throughfall increase N₂O emissions by increasing the soil content of DOC, which acts as an energy source for heterotrophic nitrifiers and denitrifiers¹⁵. Currently, field simulation experiments are rare that investigate the effects of the multifactor interaction between nitrogen addition and rainfall reduction on forest soil N₂O emissions. Smolander *et al.*¹⁹ found that N mineralization is apparently inhibited more during drought in soil containing a supplemental nitrogen addition than in the control soil, indicating an interaction between drought and nitrogen addition. However, the combined effect of nitrogen addition and rainfall reduction on forest soil N₂O emissions remains unknown.

¹Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese Academy of Sciences, 72 Wenhua Road, Shenyang 110016, China. ²University of Chinese Academy of Sciences, Beijing, 100039, China. Correspondence and requests for materials should be addressed to J.Z. (email: jhzhang@iae.ac.cn)




Figure 1. Daily mean air temperature, soil temperature (5 cm soil depth), and cumulative precipitation (A). N_2O emission rates from three replicates in NF (B), FF (C), RR (D) and RRF (E) treatments measured from November 2012 to October 2014 are shown by the black circles. Soil moisture (WFPS, %) at a 5 cm soil depth was only determined during the growing seasons.

Broad-leaved Korean pine (*Pinus koraiensis*) mixed forest covers an area of 42 334 ha in the Changbai Mountain region and is one of the dominant forest types in northeastern China. This area is exposed to high nitrogen deposition of at least $23 \text{ kg N ha}^{-1} \text{ a}^{-1}$ ²⁰ and receives less precipitation in the summer and fall²¹. To examine the influence of an increase in nitrogen deposition, a decrease in precipitation and the effect of annual variability in temperature and precipitation on soil N_2O emissions, both nitrogen addition and rainfall reduction were experimentally manipulated at this site. We hypothesized that rainfall reduction would amplify the stimulatory effect of nitrogen addition on N_2O emissions, for less leaching losses of nitrogen²².

Results

Environmental conditions. Compared with the long-term annual mean air temperature (3.2°C) and precipitation (700 mm), the annual precipitation in 2013 was higher, and extremely heavy daily precipitation occurred. The cumulative precipitation in two days exceeded 150 mm (8/15 and 8/16; Fig. 1A). By contrast, 2014 was warmer and drier than long-term averages, with a mean air temperature of 4.3°C and 605.6 mm of precipitation.

Significant annual and seasonal fluctuations in soil moisture were observed (Table 1). The soil moisture in all plots in 2014 was lower than that in 2013, which was consistent with the precipitation. The lowest soil moisture occurred in June 2013 and from August to October in 2014, ranging from 5.9% to 48.6% (WFPS; Fig. 1B–E). Although the magnitude of throughfall reaching the soil was reduced in the RR treatment plots, the differences observed in soil moisture between NF and RR treatments were not significant ($P > 0.05$, Table 2). In the nitrogen

	All treatments			Nitrogen addition			Rainfall reduction			Nitrogen addition and rainfall reduction		
	df	F	P	df	F	P	df	F	P	df	F	P
N₂O												
Treatment	3	16.34	****	1	45.18	****	1	0.03		1	3.82	*
Month	11	16.96	****	11	5.63	****	11	3.92	****	11	3.20	***
Year	1	3.79	.	1	3.36	.	1	2.69		1	0.04	
Soil moisture												
Treatment	3	4.28	**	1	11.48	***	1	0.49		1	0.90	
Month	7	80.44	****	7	3.08	**	7	2.19	*	7	0.43	
Year	1	267.29	****	1	0.19		1	0.45		1	0.01	
Soil temperature												
Treatment	3	0.00		1	0.00		1	0.00		1	0.00	
Month	7	1716.93	****	7	0.09		7	0.08		7	0.09	
Year	1	18.17	****	1	1.37		1	1.41		1	1.92	
NH₄⁺												
Treatment	3	0.19		1	0.12		1	0.13		1	0.97	
Month	4	16.13	****	5	2.71	*	5	1.55		5	0.66	
NO₃⁻												
Treatment	3	4.10	*	1	10.00	*	1	7.94	*	1	0.36	
Month	4	14.95	****	5	2.07	.	5	1.41		5	0.34	
DOC												
Treatment	3	5.03	*	1	13.70	**	1	1.30		1	0.07	
Month	4	22.56	****	5	0.73		5	1.96		5	0.73	
pH												
Treatment	3	0.15		1	0.11		1	0.15		1	0.18	
Month	5	7.29	***	5	0.78		5	0.30		5	0.59	

Table 1. ANOVA results of the linear mixed-effects model testing the treatment, seasonal and yearly effects of experimental manipulations on soil variables. $P < 0.1$. * $P < 0.05$. ** $P < 0.01$. *** $P < 0.001$. **** $P < 0.0001$.

		NF	FF	RR	RRF
Soil moisture [†] (WFPS, %)	2013	30.17 ± 2.47 aA	41.16 ± 1.37 bA	31.06 ± 4.65 aA	37.25 ± 1.79 abA
	2014	24.15 ± 1.25 aA	34.63 ± 4.44 bA	24.40 ± 3.16 aA	30.21 ± 0.69 abB
Soil temperature [†] °C	2013	11.69 ± 0.20 aA	11.90 ± 0.00 aA	11.73 ± 0.22 aA	11.68 ± 0.20 aA
	2014	11.28 ± 0.10 aA	11.06 ± 0.01 aB	11.28 ± 0.10 aA	11.28 ± 0.10 aA
NH ₄ ⁺ (mg N kg ⁻¹) [‡]		19.99 ± 1.87 a	17.25 ± 1.78 a	17.24 ± 1.99 a	18.53 ± 1.01 a
NO ₃ ⁻ (mg N kg ⁻¹) [‡]		22.86 ± 2.71 a	31.51 ± 4.04 a	30.35 ± 1.58 a	43.08 ± 4.40 b
DOC (mg C kg ⁻¹) [‡]		300.08 ± 13.53 ab	236.69 ± 11.71 a	347.66 ± 36.96 b	259.32 ± 28.73 a
pH [‡]		5.27 ± 0.51 a	5.52 ± 0.26 a	5.54 ± 0.24 a	5.51 ± 0.24 a

Table 2. Annual means (±SE) for soil variables in different treatment plots in a broad-leaved Korean pine (*Pinus koraiensis*) mixed forest. [†]Measured across the 2013 and 2014 growing seasons. [‡]Measured only for the growing season in 2014. Different capital letters indicate a significant difference at the level of 0.05 between years for the same treatment; different lowercase letters indicate a significant difference at the level of 0.05 among treatments in the same year.

fertilizer treatments (FF and RRF), soil moisture was relatively higher than that in nonfertilizer treatments (NF and RR; Table 2).

Soil temperature followed a pattern similar to that of air temperature during the experiment, with significant annual and seasonal differences ($P < 0.0001$, Table 1). However, no significant differences in temperature were observed among treatments ($P > 0.05$, Table 2).

Significant differences in the seasonal dynamics of NH₄⁺, NO₃⁻, DOC and pH were observed ($P < 0.0001$, Table 1). However, we did not observe significant differences in soil NH₄⁺ among treatments (Table 2), whereas nitrogen addition increased concentrations of soil NO₃⁻ and DOC compared with the control plots. Soil pH was not significantly affected by the treatments ($P > 0.05$, Table 1).

Seasonal and annual dynamics of N₂O emissions. For the 2 years of this study, the mean annual N₂O emission was 1.25 ± 0.22 kg N₂O-N ha⁻¹ a⁻¹ in the NF treatment; however, the emissions during 2014 (1.41 ± 0.32 kg N₂O-N ha⁻¹ a⁻¹) were approximately 30% higher than those during 2013 (1.09 ± 0.13 kg N₂O-N

		NF	FF	RR	RRF
2013	Winter	0.24 ± 0.02 (22.22) a	0.44 ± 0.11 (21.47) a	0.36 ± 0.08 (40.30) a	0.55 ± 0.11 (21.01) a
	Freeze-thaw period	0.12 ± 0.02 (10.83) a	0.23 ± 0.05 (11.33) b	0.21 ± 0.03 (23.27) ab	0.49 ± 0.01 (18.76) c
	Growing season	0.77 ± 0.09 (70.45) a	1.37 ± 0.04 (67.19) c	0.33 ± 0.05 (36.43) b	1.57 ± 0.17 (60.23) c
2014	Winter	0.49 ± 0.14 (34.42) ab	0.35 ± 0.08 (16.97) ab	0.24 ± 0.04 (31.38) a	0.63 ± 0.11 (28.11) b
	Freeze-thaw period	0.47 ± 0.09 (33.28) bc	0.30 ± 0.06 (14.36) ab	0.17 ± 0.06 (22.84) a	0.60 ± 0.06 (26.94) c
	Growing season	0.46 ± 0.10 (32.30) a	1.41 ± 0.44 (68.67) b	0.35 ± 0.06 (45.78) a	1.00 ± 0.19 (44.94) ab

Table 3. Seasonal N₂O-N emissions (Mean ± SE, kg N₂O-N ha⁻¹) in different treatments, with the percentage contribution to annual total emissions in brackets (Mean ± SE, n = 3). Different letters indicate a significant difference at the level of 0.05 among different treatments during the same season.




Figure 2. Annual cumulative N₂O emissions in the different treatments (NF, FF, RR and RRF). Different lowercase letters indicate a significant difference at the level of 0.05 among treatments in the same year.

ha⁻¹ a⁻¹). The seasonal dynamics of the N₂O emission rates in the four treatments during the two-year observation period are shown in Fig. 1B–E. The rates were generally consistent with the level of soil moisture during the growing season. In the control plots, the highest N₂O emission rates generally occurred in July and August in 2013, when the soil moisture was high due to abundant precipitation. However, in 2014, the highest N₂O emission rates occurred in late April and early May, when the soil experienced freeze-thaw events.

The seasonal, cumulative N₂O emissions in the control plots were different between the two years (Table 3). In 2013, the N₂O emissions during the growing season accounted for more than 70% of the annual emissions. However, the emissions in the three seasonal periods were approximately equal in 2014, with each period accounting for approximately 1/3 of the annual emissions.

Effects of nitrogen addition and rainfall reduction on N₂O emissions. Nitrogen addition significantly increased the N₂O emission rates (Table 1), and the annual emissions increased by 87% in 2013 and 45% in 2014, averaging an increase of 63% across the two years (Fig. 2). Nitrogen addition stimulated N₂O emissions by 78% during the growing season in 2013 and by 209% in 2014. However, the effect of nitrogen addition on N₂O emissions was not always positive. In 2014, compared with the NF treatment, the cumulative N₂O emissions in the FF treatment decreased by 28% and 37% during the winter and freeze-thaw period, respectively (Table 3). Rainfall reduction decreased the annual N₂O emissions by 17% in 2013 and 45% in 2014, averaging a decrease of 33% across the two years (Fig. 2), with the decrease significant ($P < 0.05$) during the growing season in 2013 and the following freeze-thaw period in 2014 (Table 3). A significant interaction was detected between nitrogen addition and rainfall reduction on N₂O emission rates ($P < 0.05$, Table 1). Although the annual N₂O emissions decreased in the RR treatment, the annual N₂O emissions in RRF plots were 19% (29% in 2013 and 9% in 2014) higher than those in FF plots, with the increase significant ($P = 0.036$) in 2013 (Fig. 2). Seasonally, the cumulative N₂O emissions in RRF plots were higher than those in the NF plots (Table 3).

The EF_N and EF_{NP} were 1.59% (1.89% in 2013 and 1.28% in 2014) and 3.19% (3.43% in 2013 and 2.94% in 2014), respectively, in the broad-leaved Korean pine mixed forest at Changbai Mountain, China.

Effects of environmental factors on N₂O emissions. Stepwise multiple linear regressions with backward elimination were used to examine the relationships between the measured soil variables and N₂O emissions across all treatments during the entire 2014 growing season. In this model, the best predictors of observed N₂O emissions following log-transformation were soil moisture ($t = 7.68$, $P < 0.001$), soil temperature ($t = 4.85$,




Figure 3. Variable importance (VIMP) for predicting N₂O emissions.

$P < 0.001$) and DOC ($t = -1.92$, $P = 0.059$). The parameter estimations obtained from this model were determined in the following equation (residual standard error 0.67, $df = 68$; F-statistic 29.23, $P < 0.001$; multiple $R^2 = 0.56$, Adjusted $R^2 = 0.54$):

$$\ln[\text{N}_2\text{O}] = 0.05 \times [\text{soil moisture}] + 0.13 \times [\text{soil temperature}] - 0.001[\text{DOC}] - 0.29$$

In general, soil inorganic N (NH_4^+ and NO_3^-) concentration is the critical factor influencing N_2O emissions. However, based on the stepwise multiple linear regressions, the effects of NH_4^+ and NO_3^- on N_2O emissions were not significant and could be eliminated. The regression model with the random ForestSRC package in the R statistical software package was also used to determine the influences of environmental factors and soil nutrients on N_2O emissions. The values of variable importance (VIMP) in order were soil moisture > soil temperature > NO_3^- > NH_4^+ > DOC > pH (Fig. 3). The response of N_2O emissions for each soil variable is shown in Fig. 4. Rates of N_2O emission increased with soil moisture and soil temperature when soil moisture ranged from 5% to 45% and soil temperature ranged from 7 °C to 17 °C. N_2O emission rates increased rapidly when NH_4^+ and NO_3^- concentrations were below 20 mgN kg⁻¹, and then increased slowly. The N_2O emission rates exhibited a negative relationship with DOC when the concentration was below 300 mg of C kg soil⁻¹. The effect of pH on N_2O emission rates was segmented, and the demarcation point was 6.

Discussion

The mean annual N_2O emission from NF plots was 1.25 ± 0.22 kg $\text{N}_2\text{O-N ha}^{-1} \text{ a}^{-1}$, which is consistent with data reported in previous studies on N_2O emissions for temperate forest soils^{12,23–25}. However, low emissions of less than 0.1 kg $\text{N}_2\text{O-N ha}^{-1} \text{ a}^{-1}$ ^{26,27} and high emissions that exceed 4 kg $\text{N}_2\text{O-N ha}^{-1} \text{ a}^{-1}$ ^{28,29} have both been reported. The marked differences in emissions might be due to spatial variation, and the annual N_2O emissions could make a large difference in the same area³⁰. These large temporal and spatial variations in emissions can be attributed to two causes. First, soil N_2O emissions are influenced by numerous factors, including precipitation, temperature, forest type, and soil texture and characteristics⁷, and small changes in these variables may result in large differences in N_2O emissions³. Second, N_2O emissions are characterized by short pulse emissions related to nitrogen deposition³¹, precipitation^{32,33}, freeze-thaw cycles³⁴ and drying-wetting events³⁵. And these pulse events of N_2O emission may be extremely important contributions to the total annual budget³⁰. Field studies on N_2O emissions often use discontinuous measurements performed with chambers over durations of weeks to months^{36,37}, and even automated measuring systems can have observation intervals. Thus, missing pulse emission events or measuring during pulse emission events could result in underestimating or overestimating annual N_2O emission, respectively³⁸.

During the growing season, N_2O emissions were higher in 2013 than in 2014 ($P = 0.053$), which was likely because of higher soil moisture due to abundant rain in summer 2013. However, during the non-growing seasons, particularly the freeze-thaw period, the annual N_2O emission in 2014 was larger than that in 2013. With less snow cover and lower soil temperatures, the creation of an adverse soil environment can result in increased mortality of plant roots and soil biota³⁹, which leads to the accumulation of labile soil organic carbon and nitrogen. Therefore, when this sufficient supply of soil nutrients was coupled with suitable soil moisture, a burst in N_2O emissions occurred during the freeze-thaw period in 2014. Additionally, the result of the high N_2O emission rates during the freeze-thaw period was likely greatly reduced soil concentrations of nitrate or carbon³³, which then led to reduced N_2O emissions during the growing season in 2014.

The contribution to N_2O emissions from the nitrogen addition was significant in the broad-leaved Korean pine mixed forest, and the emission factor was 1.59%. Increases in forest soil N_2O emissions following nitrogen




Figure 4. Partial correlations between N_2O emission rates and soil variables during the growing season in 2014 in the broad-leaved Korean pine (*Pinus koraiensis*) mixed forest.

addition are widely reported^{13,14}, but the N_2O emission factors differ by region and forest type. In this study, the N_2O emission factor was higher than the 0.1% for a spruce forest in Germany⁴⁰, the 0.7–0.8% for a subtropical forest in southern China¹³, and the default factor of 1% used by the IPCC¹. In natural ecosystems, particularly those N-limited and without a history of N addition, emission levels of N_2O are generally low and the responses to added N are weak⁴¹. By contrast, our study area was exposed to high levels of nitrogen deposition estimated at more than $23 \text{ kg N ha}^{-1} \text{ a}^{-1}$ ²⁰; and the FF treatment plots received supplemental nitrogen for four years before beginning the N_2O observations. Therefore, additional N_2O would be emitted in this area compared with natural forest following nitrogen addition, resulting in a higher N_2O emission factor. Compared with the NF plots, the cumulative N_2O emission of the FF plots was significantly higher during the growing season in both years (Table 3). Pulse N_2O emissions 2- to 4-fold greater were observed after nitrogen addition (Fig. 1C), which was consistent with previous studies conducted in the identical area³¹; the increase in emission rates persisted for approximately two weeks. These pulse emissions were the primary reason for the differences between FF and NF plots during the growing season. Additionally, soil moisture and NO_3^- concentrations were higher in FF plots than those in NF plots (Table 2), which provided a more favourable environment for denitrification, leading to the increase in N_2O emissions. In winter and the freeze-thaw period, the differences in cumulative N_2O emissions between NF and FF plots decreased compared with that in growing season (Table 3). There are two possible explanations for this decrease in differences. First, N was only applied from May to October, and therefore, the effect of N addition likely decreased during the non-growing season. Second, the limiting factors might be soil temperature and/or soil moisture rather than soil N content during winter and the freeze-thaw period⁴².

As expected, rainfall reduction lowered the annual N_2O emissions by 17–45%, similar to reports in previous studies^{16,17,43}. However, no significant difference in soil moisture was observed between RR and NF plots, which was similar to the results for a tropical forest in which total rainfall was reduced by 25% and 50%¹⁵. This lack of response in soil moisture to rainfall reduction might be because precipitation is mainly concentrated in summer period, and subsurface lateral water flow reduced the effect of rainfall reduction. Although rainfall reduction had no significant effect on soil moisture; it did decrease the precipitation pulses which could cause short peak emissions of N_2O emission^{32,33}.

The combination treatment of nitrogen addition and rainfall reduction increased N_2O emissions by 58–140% and had a larger stimulatory effect on emissions than the addition of nitrogen alone. A reduction in rainfall could prevent the added nitrogen from leaching out of the soil²². Thus, more nitrogen would remain in the soil, resulting in increased N_2O emissions under RRF conditions compared with those under FF, as evidenced by the significantly higher soil NO_3^- concentration on RRF compared to FF plots (Table 2). Because the level of precipitation in 2014 was far less than that in 2013, the protective effect of rain reduction on nitrogen would be expected to weaken, which helped to explain the difference in N_2O emissions between RRF and FF plots between 2014 and

2013 (Fig. 2). Therefore, we concluded that a reduction in rainfall might increase the stimulatory effect of N addition on soil N₂O emissions.

Stepwise multiple linear regressions with backward elimination indicated that soil moisture, soil temperature and DOC were the best predictors of the observed N₂O emission patterns during the growing season in 2014. Soil moisture and soil temperature are widely reported as the best predictors of N₂O emissions in other temperate forests^{12,23,44}. Previous studies have demonstrated that a WFPS value of 60% is an important threshold because nitrification activity is the highest and denitrification activity begins to increase when the WFPS is approximately 60%^{45,46}. In our study, soil WFPS values were slightly less than 60%, indicating that nitrification might be the most important process regulating N₂O emissions in the broad-leaved Korean pine mixed forest. Significant positive correlations between soil temperature and N₂O emissions were observed in this study that were consistent with those reported in other studies^{23,44}. In our study, N₂O emissions and DOC concentration were slightly negatively correlated. In general, as a labile nutrient, a higher concentration of DOC promotes the activities of soil biota (i.e., nitrifiers and denitrifiers), resulting in increases in N₂O emissions. However, with the addition of external N, soil N₂O emissions and DOC concentration might become negatively correlated⁴⁷. On one hand, N addition could stimulate soil N₂O emissions and consume more DOC; on the other hand, N addition could inhibit soil C mineralization and decrease soil DOC concentration^{48,49}, as evidenced by the higher N₂O emissions and lower DOC concentrations in FF than in NF plots.

In addition, soil inorganic N (NH₄⁺ and NO₃⁻) content was the essential factor influencing N₂O emission. However, based on stepwise multiple linear regressions, the effects of NH₄⁺ and NO₃⁻ were not significant and could be eliminated from the model. To further investigate the influence of soil variables on N₂O emissions, we used the regression model provided with the randomForestSRC package in the R statistical software package. Similar to the results of the stepwise multiple linear regressions, soil moisture and soil temperature were also the two most important factors influencing N₂O emissions. However, NO₃⁻ and NH₄⁺ were more important factors than DOC in this model. From the partial correlation plots (Fig. 4), we observed that the N₂O emission rates were only influenced by NO₃⁻ or NH₄⁺ at concentrations less than 20 mgN kg⁻¹. This small range of influence might explain why inorganic nitrogen factors were eliminated by the stepwise multiple linear regressions.

Conclusions

The response of soil N₂O emissions to nitrogen addition and rainfall reduction was studied for two continuous years in an old broad-leaved Korean pine mixed forest at Changbai Mountain in northeastern China. We found that nitrogen addition significantly increased N₂O emissions and rainfall reduction slightly decreased N₂O emissions. However, the combination of nitrogen addition and rainfall reduction stimulated N₂O emissions more than the nitrogen addition alone. This result could be explained by the reduction in rainfall preventing the added nitrogen from leaching out of the soil, and therefore, with more nitrogen remaining in the soil, N₂O emissions increased. Our findings indicated that drought would slightly decrease forest soil N₂O emission; however, with increasing atmospheric N deposition, drought might alter to increase the N₂O emission in temperate forests.

Methods

Study site. The experiment was conducted in an old broad-leaved Korean pine (*Pinus koraiensis*) mixed forest (~200 years old) on Changbai Mountain (42°24' N, 128°05' E; 766 m a.s.l.) in northeastern China. The detailed description of this site can be found in Wang *et al.*⁵⁰. The site is characterized as having a temperate, continental climate. The mean annual temperature is 3.2 °C, and the mean annual precipitation is 700 mm, primarily occurring from June to August. The soil in this region was developed from volcanic ash and is classified as a Eutric cambisol (FAO classification) containing a high content of organic matter in the organic horizon (O-horizon). The bulk density was 0.35 g cm⁻³ in the surface soil (0–10 cm), and the contents of total carbon, nitrogen and phosphorus were 11.8%, 0.9% and 0.1%, respectively.

Experimental design. The experiment was initiated in September 2009 with the following four treatments: rainfall-reduction (RR), N-fertilization (FF), rainfall-reduction and N-fertilization (RRF) and control plots (NF). The study design included three replicates of each treatment. Thus, twelve 50 × 50 m plots with a buffer zone of at least 20 m between any two plots were established randomly within the study area. For the RR treatment, V-shaped polycarbonate (PC) of high transparency (approximately 95%) was used to prevent approximately 30% of natural throughfall. The PC was fixed on an aluminium frame ~1 m above the soil surface. For the FF treatment, ammonium nitrate (NH₄NO₃) solution was applied monthly (six times from May to October each year) with a sprayer. The application rate of 50 kg N ha⁻¹a⁻¹ was approximately double the annual total N deposition (~23 kg N ha⁻¹a⁻¹) in this area²⁰. In the RRF treatment, throughfall was reduced by 30% and 50 kg N ha⁻¹a⁻¹ as NH₄NO₃ fertilizer was applied. The control plots received natural rainfall and atmospheric N deposition.

N₂O emissions. N₂O emissions were measured using the static opaque chamber method³⁷. Trapezoid-shaped chambers with the body constructed of polyvinyl chloride plastic (top: 55 × 40 cm; bottom: 47 × 32 cm; height: 30 cm) were permanently inserted into the forest soil to a depth of 5 cm, which resulted in a volume of 0.055 m³. Two parallel chambers were installed in each plot five months before gas sampling.

Emissions of N₂O were measured from November 2012 to October 2014. Measurements were conducted between 8:00 and 11:00 a.m. every three to four days (8 times per month) during the growing season (April to October) and every seven to eight days (4 times per month) during the non-growing season. On each sampling, plastic tops were fitted to the chambers to ensure a gas-tight environment, and a 60-mL plastic syringe with a three-way valve was used to obtain air samples. During a 1 h incubation at 15 min intervals, five gas samples were collected from each chamber. Air temperature inside the chambers was recorded using a temperature sensor probe (JM624; Jinming Instrument CO., LTD, Tianjin, China).

All air samples were analysed within 24 h of sampling using a gas chromatograph (Agilent-7890A; Agilent Technologies, USA) equipped with an electron capture detector. The initial rates of change of the gas concentrations within the chambers were determined using significant nonlinear (exponential) or linear fitting of the five concentration observations versus enclosure time. An observation was rejected as null datum when the correlation between the gas concentrations and the enclosure time was not significant ($P < 0.05$)⁵¹.

Meteorological data and soil physicochemical analyses. Meteorological data (i.e., air temperature, soil temperature and precipitation) were obtained from the Changbai Forest Ecosystem Research Station. During the non-frozen period (April through October), the soil temperature (5 cm) and soil moisture (WFPS, 0–6 cm) of each field plot were manually measured simultaneously during gas sampling using a temperature sensor probe (JM624; Jinming Instrument CO., LTD, Tianjin, China) and time-domain reflectometry (Hydro Sense II; USA), respectively. To determine the factors that influenced N₂O emissions, soil samples were collected monthly from a depth of 10 cm in each plot using a 3 cm diameter soil auger from May to October in 2014. The soil samples were analysed for ammonium (NH₄⁺), nitrate (NO₃⁻), and dissolved organic carbon (DOC) concentrations and pH. Inorganic nitrogen (NH₄⁺ and NO₃⁻) was extracted from twenty grams of soil for 1 h in 100 mL of a 2 M potassium chloride (KCl) solution. The concentrations of NH₄⁺ and NO₃⁻ were determined with an automatic nitrogen analyser (AA3; BRAN & Lubbe, Nordstedt, Germany). The DOC was extracted by shaking 20 g of soil sample for 1 h in 100 mL of deionized water. The extracts were centrifuged at 6,000 rpm for 15 min. The supernatant was filtered through a 0.45 μm polyethersulfone membrane filter (Membran, Germany) before analysis with a Multi N/C 3000 analyser (Analytic Jena AG, Germany). Air-dried soils were sieved through a 2 mm mesh, and the pH was determined by electrode at a soil: water ratio of 1:1.

Statistical analyses. All statistical analyses were conducted using the R statistical software package (version 3.2.0, Core Team 2009). Linear mixed-effects models with plot as the random factor were used to test the effects of treatment and time (i.e., month and year) on N₂O emission rates and soil variables. Annual means for all variables were calculated for each plot and were analysed using linear mixed-effects models, with treatment and year as fixed factors and plot as the random factor. Seasonal cumulative N₂O emissions were calculated for each plot and were analysed using linear mixed-effects models, with treatment, season and year as fixed factors and plot as the random factor. Similarly, annual cumulative N₂O emissions were calculated for each plot and were analysed using linear mixed-effects models, with treatment and year as fixed factors and plot as the random factor.

The Shapiro test revealed that N₂O emissions were log-distributed. After log transformation, stepwise multiple linear regressions with backward elimination using a linear model were conducted to examine the relationships between monthly mean N₂O emissions and soil variables. We also used the randomForestSRC package (version 1.6.1) to assess the influences of soil variables on N₂O emissions.

The N₂O emission factors under nitrogen addition (EF_N) and nitrogen addition with rainfall reduction (EF_{NP}) for broad-leaved Korean pine mixed forest soil were calculated as follow:

$$EF_N = ([E_N - E_0]/N_{\text{added}}) \times 100\% \quad (1)$$

$$EF_{NP} = ([E_{NP} - E_0]/N_{\text{added}}) \times 100\% \quad (2)$$

where E_N, E_{NP} and E₀ are the N₂O emissions from the FF, RRF and NF plots during the same period, respectively, and N_{added} is the amount of nitrogen fertilizer applied.

References

- IPCC. *Climate change 2007-the physical science basis: Working group I contribution to the fourth assessment report of the IPCC*. Vol. 4 (Cambridge University Press, 2007).
- Ravishankara, A., Daniel, J. S. & Portmann, R. W. Nitrous oxide (N₂O): the dominant ozone-depleting substance emitted in the 21st century. *Science* **326**, 123–125 (2009).
- Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R. & Zechmeister-Boltenstern, S. Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philosophical Transactions of the Royal Society B: Biological Sciences* **368** (2013).
- Seitzinger, S. P., Kroeze, C. & Styles, R. V. Global distribution of N₂O emissions from aquatic systems: natural emissions and anthropogenic effects. *Chemosphere - Global Change Science* **2**, 267–279 (2000).
- Thomson, A. J., Giannopoulos, G., Pretty, J., Baggs, E. M. & Richardson, D. J. Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. *Philos. Trans. R. Soc. B-Biol. Sci.* **367**, 1157–1168 (2012).
- Stange, F. & Döhling, F. ¹⁵N tracing model SimKIM to analyse the NO and N₂O production during autotrophic, heterotrophic nitrification, and denitrification in soils. *Isotopes in environmental and health studies* **41**, 261–274 (2005).
- Rowlings, D. W., Grace, P. R., Kiese, R. & Weier, K. L. Environmental factors controlling temporal and spatial variability in the soil-atmosphere exchange of CO₂, CH₄ and N₂O from an Australian subtropical rainforest. *Global Change Biology* **18**, 726–738 (2012).
- Weslien, P., Klemetsson, A. K., Borjesson, G. & Klemetsson, L. Strong pH influence on N₂O and CH₄ fluxes from forested organic soils. *European Journal of Soil Science* **60**, 311–320 (2009).
- Galloway, J. N. *et al.* Transformation of the nitrogen cycle: Recent trends, questions, and potential solutions. *Science* **320**, 889–892 (2008).
- Dore, M. H. I. Climate change and changes in global precipitation patterns: What do we know? *Environ Int* **31**, 1167–1181 (2005).
- Ambus, P., Zechmeister-Boltenstern, S. & Butterbach-Bahl, K. Sources of nitrous oxide emitted from European forest soils. *Biogeosciences* **3**, 135–145 (2006).
- Papen, H. & Butterbach-Bahl, K. A 3-year continuous record of nitrogen trace gas fluxes from untreated and limed soil of a N-saturated spruce and beech forest ecosystem in Germany - 1. N₂O emissions. *Journal of Geophysical Research-Atmospheres* **104**, 18487–18503 (1999).
- Zhang, W. *et al.* Responses of nitrous oxide emissions to nitrogen and phosphorus additions in two tropical plantations with N-fixing vs. non-N-fixing tree species. *Biogeosciences* **11**, 4941–4951 (2014).
- Hall, S. J. & Matson, P. A. Nitrogen oxide emissions after nitrogen additions in tropical forests. *Nature* **400**, 152–155 (1999).

15. Wieder, W. R., Cleveland, C. C. & Townsend, A. R. Throughfall exclusion and leaf litter addition drive higher rates of soil nitrous oxide emissions from a lowland wet tropical forest. *Global Change Biology* **17**, 3195–3207 (2011).
16. Goldberg, S. D. & Gebauer, G. N₂O and NO fluxes between a Norway spruce forest soil and atmosphere as affected by prolonged summer drought. *Soil Biol. Biochem.* **41**, 1986–1995 (2009).
17. Davidson, E. A., Ishida, F. Y. & Nepstad, D. C. Effects of an experimental drought on soil emissions of carbon dioxide, methane, nitrous oxide, and nitric oxide in a moist tropical forest. *Global Change Biology* **10**, 718–730 (2004).
18. Vasconcelos, S. S. *et al.* Moisture and substrate availability constrain soil trace gas fluxes in an eastern Amazonian regrowth forest. *Global Biogeochemical Cycles* **18** (2004).
19. Smolander, A., Barnette, L., Kitunen, V. & Lumme, I. N and C transformations in long-term N-fertilized forest soils in response to seasonal drought. *Applied Soil Ecology* **29**, 225–235 (2005).
20. Lu, C. Q. & Tian, H. Q. Spatial and temporal patterns of nitrogen deposition in China: Synthesis of observational data. *Journal of Geophysical Research-Atmospheres* **112** (2007).
21. Piao, S. L. *et al.* The impacts of climate change on water resources and agriculture in China. *Nature* **467**, 43–51 (2010).
22. Thomas, D., Johannes, K., David, K., Rüdiger, G. & Ralf, K. Impacts of management and climate change on nitrate leaching in a forested karst area. *Journal of environmental management* **165**, 243–252 (2016).
23. Wu, X. *et al.* Environmental controls over soil-atmosphere exchange of N₂O, NO, and CO₂ in a temperate Norway spruce forest. *Global Biogeochemical Cycles* **24** (2010).
24. Horvath, L., Fuhrer, E. & Lajtha, K. Nitric oxide and nitrous oxide emission from Hungarian forest soils; linked with atmospheric N-deposition. *Atmospheric Environment* **40**, 7786–7795 (2006).
25. Chen, G. X. *et al.* Nitrous oxide emissions from terrestrial ecosystems in China. *Chemosphere - Global Change Science* **2**, 373–378 (2000).
26. Jungkunst, H. F., Fiedler, S. & Stahr, K. N₂O emissions of a mature Norway spruce (*Picea abies*) stand in the Black Forest (southwest Germany) as differentiated by the soil pattern. *Journal of Geophysical Research-Atmospheres* **109** (2004).
27. Bowden, R. D., Melillo, J. M., Steudler, P. A. & Aber, J. D. Effects of nitrogen additions on annual nitrous oxide fluxes from temperate forest soils in the northeastern United States. *Journal of Geophysical Research: Atmospheres (1984–2012)* **96**, 9321–9328 (1991).
28. Klemmedtsson, L. *et al.* Reduction of greenhouse gas emissions by wood ash application to a *Picea abies* (L.) Karst. forest on a drained organic soil. *European Journal of Soil Science* **61**, 734–744 (2010).
29. Dong, Y. S. *et al.* Experimental study on N₂O and CH₄ fluxes from the dark coniferous forest zone soil of the Gongga Mountain, China. *Sci China Ser D* **46**, 285–295 (2003).
30. Butterbach-Bahl, K., Rothe, A. & Papen, H. Effect of tree distance on N₂O and CH₄-fluxes from soils in temperate forest ecosystems. *Plant Soil* **240**, 91–103 (2002).
31. Bai, E. *et al.* Pulse Increase of soil N₂O emission in response to N addition in a temperate forest on Mt Changbai, Northeast China. *Plos One* **9** (2014).
32. Liu, X. C. *et al.* Response of soil N₂O emissions to precipitation pulses under different nitrogen availabilities in a semiarid temperate steppe of Inner Mongolia, China. *Journal of Arid Land* **6**, 410–422 (2014).
33. Zona, D. *et al.* N₂O fluxes of a bio-energy poplar plantation during a two years rotation period. *Global Change Biology Bioenergy* **5**, 536–547 (2013).
34. Wolf, B. *et al.* Grazing-induced reduction of natural nitrous oxide release from continental steppe. *Nature* **464**, 881–884 (2010).
35. Beare, M. H., Gregorich, E. G. & St-Georges, P. Compaction effects on CO₂ and N₂O production during drying and rewetting of soil. *Soil Biol. Biochem.* **41**, 611–621 (2009).
36. Morishita, T., Aizawa, S., Yoshinaga, S. & Kaneko, S. Seasonal change in N₂O flux from forest soils in a forest catchment in Japan. *Journal of Forest Research* **16**, 386–393 (2011).
37. Bowden, R. D., Steudler, P. A., Melillo, J. M. & Aber, J. D. Annual nitrous-oxide fluxes from temperate forest soils in the northeastern United-States. *Journal of Geophysical Research-Atmospheres* **95**, 13997–14005 (1990).
38. Barton, L. *et al.* Sampling frequency affects estimates of annual nitrous oxide fluxes. *Scientific Reports* **5** (2015).
39. Hardy, J. P. *et al.* Snow depth manipulation and its influence on soil frost and water dynamics in a northern hardwood forest. *Biogeochemistry* **56**, 151–174 (2001).
40. Eickenscheidt, N., Brumme, R. & Veldkamp, E. Direct contribution of nitrogen deposition to nitrous oxide emissions in a temperate beech and spruce forest - a N-15 tracer study. *Biogeosciences* **8**, 621–635 (2011).
41. Bouwman, A. F., Boumans, L. J. M. & Batjes, N. H. Emissions of N₂O and NO from fertilized fields: Summary of available measurement data. *Global Biogeochemical Cycles* **16** (2002).
42. Shrestha, R. K., Strahm, B. D. & Sucre, E. B. Greenhouse gas emissions in response to nitrogen fertilization in managed forest ecosystems. *New Forests* **46(2)**, 167–193 (2015).
43. Nepstad, D. C. *et al.* The effects of partial throughfall exclusion on canopy processes, aboveground production, and biogeochemistry of an Amazon forest. *Journal of Geophysical Research-Atmospheres* **107(D20)** (2002).
44. Borken, W., Xu, Y. J., Davidson, E. A. & Beese, A. Site and temporal variation of soil respiration in European beech, Norway spruce, and Scots pine forests. *Global Change Biology* **8** (2002).
45. Mu, Z. J., Kimura, S. D., Toma, Y. & Hatano, R. Nitrous oxide fluxes from upland soils in central Hokkaido, Japan. *J Environ Sci-China* **20**, 1312–1322 (2008).
46. Davidson, E. A. Fluxes of nitrous oxide and nitric oxide from terrestrial ecosystems. In *Microbial production and consumption of greenhouse gases: methane, nitrogen oxides, and halomethanes*. (eds. J. E. Rogers & W. B. Whitman) 219–235 (1991).
47. Huang, Y. N₂O emission of farmland in The exchange of carbon and nitrogen in the system of earth and atmosphere: from experiment to model. (ed. Huang, Y.) Ch. 5, 130–162 (2003).
48. Filep, T. & Rekas, M. Factors controlling dissolved organic carbon (DOC), dissolved organic nitrogen (DON) and DOC/DON ratio in arable soils based on a dataset from Hungary. *Geoderma* **162(3–4)**, 312–318 (2011).
49. Guo, R. H., Zheng, J. Q., Han, S. J., Zhang, J. H. & Li, M. H. Carbon and nitrogen turnover in response to warming and nitrogen addition during early stages of forest litter decomposition-an incubation experiment. *Journal of Soils and Sediments* **13(2)**, 312–324 (2013).
50. Wang, C. *et al.* Responses of Fine roots and soil N availability to short-term nitrogen fertilization in a broad-leaved Korean Pine mixed forest in northeastern China. *Plos One* **7** (2012).
51. Mei, B. L. *et al.* Characteristics of multiple-year nitrous oxide emissions from conventional vegetable fields in southeastern China. *Journal of Geophysical Research-Atmospheres* **116** (2011).

Acknowledgements

This work was funded by the strategic priority research program of the Chinese Academy of Sciences (XDA05020300) and the National Natural Science Foundation of China (41430639, 41675150, 41330530, and 31470522).

Author Contributions

S.G., J.Z. and S.H. designed the experiment. S.G., Z.C. and F.W. conducted the experiments. S.G. wrote the manuscript text and executed statistical analysis. J.Z. revised the manuscript text and analysis methods. All authors reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Geng, S. *et al.* Rainfall reduction amplifies the stimulatory effect of nitrogen addition on N₂O emissions from a temperate forest soil. *Sci. Rep.* 7, 43329; doi: 10.1038/srep43329 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017