

Cite this: *Chem. Sci.*, 2022, 13, 11427

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 19th May 2022  
Accepted 24th August 2022

DOI: 10.1039/d2sc02827a

rsc.li/chemical-science

# A base-catalyzed approach for the anti-Markovnikov hydration of styrene derivatives†

Spencer P. Pajk,‡ Zisong Qi,§ Stephen J. Sujansky and Jeffrey S. Bandar ID\*

The base-catalyzed addition of 1-cyclopropylethanol to styrene derivatives with an acidic reaction workup enables anti-Markovnikov hydration. The use of either catalytic organic superbase or crown ether-ligated inorganic base permits hydration of a wide variety of styrene derivatives, including electron-deficient, *ortho*-substituted and heteroaryl variants. This protocol complements alternative routes to terminal alcohols that rely on stoichiometric reduction and oxidation processes. The utility of this method is demonstrated through multigram scale reactions and its use in a two-step hydration/cyclization process of *ortho*-halogenated styrenes to prepare 2,3-dihydrobenzofuran derivatives.


## Introduction

$\beta$ -Aryl alcohols are an important alcohol subclass that find wide utility in organic chemistry. This substructure is prominently featured in bioactive compounds (Fig. 1a)<sup>1</sup> and is a common synthetic intermediate for phenethyl-functionalized compounds, including several pharmaceuticals (Fig. 1b).<sup>2</sup> Medicinal chemistry discovery efforts also frequently employ this building block to perform structure activity relationship (SAR) studies on phenethyl units, with three such examples shown in Fig. 1c.<sup>3</sup> Two common approaches to  $\beta$ -aryl alcohols are reduction of arylacetic acids<sup>4</sup> and hydroboration/oxidation of styrene derivatives.<sup>5,6</sup> While reliable for simple substrates, these methods have limitations that can prevent rapid access to alcohols with diverse aryl groups. For instance, non-commercial arylacetic acids require separate multistep syntheses.<sup>7</sup> Meanwhile, traditional stoichiometric hydroboration/oxidation protocols<sup>8</sup> are often unselective and low yielding for electron-deficient or *ortho*-substituted styrenes and heteroaryl variants, thus requiring alternative hydroboration methods and reagents.<sup>9</sup> A new approach to  $\beta$ -aryl alcohols from styrene derivatives that does not rely on reduction or oxidation events could therefore improve access to this valuable class of alcohol.<sup>10</sup>

The catalytic anti-Markovnikov hydration of aryl-substituted alkenes has long been desired as a sustainable and complementary alternative to stoichiometric hydroboration/oxidation processes.<sup>11</sup> Several strategies toward this goal have recently been reported for styrenes, including Grubbs' triple relay

catalytic process,<sup>12</sup> Arnold's and Li's biocatalytic approaches,<sup>13</sup> and Lei's photocatalytic alkene oxidation method.<sup>14</sup> While these methods are impressive in their strategy for achieving anti-Markovnikov selectivity, they have yet to be applied toward more complex styrene substrates, including vinyl *N*-heteroarenes. Thus, there remains a current challenge to develop catalytic hydration methods that can access densely functionalized  $\beta$ -aryl alcohols to better reflect the structural diversity represented in Fig. 1.

We proposed to use base catalysis for the anti-Markovnikov hydration of styrene derivatives as a new approach to  $\beta$ -aryl alcohols. We recently disclosed the use of the phosphazene superbase  $P_4$ -*t*-Bu or 18-crown-6-ligated KO-*t*-Bu as basic

Fig. 1 Examples of the importance of  $\beta$ -aryl substituted alcohols.

Department of Chemistry, Colorado State University Fort Collins, Colorado, 80523 USA. E-mail: jeff.bandar@colostate.edu

† Electronic supplementary information (ESI) available. See <https://doi.org/10.1039/d2sc02827a>

‡ These authors contributed equally.






Fig. 2 Overview of a base-catalyzed approach to anti-Markovnikov styrene hydration.


catalysts for the nucleophilic addition of alcohols to aryl-substituted alkenes.<sup>15,16</sup> Water does not participate in this reaction, so we instead sought a nucleophilic “protected water” source that could be used to achieve formal hydration (Fig. 2).<sup>17</sup> Thus, following hydroetherification, facile deprotection could provide straightforward access to  $\beta$ -aryl alcohols with complete regiocontrol. We herein report the discovery of an unconventional nucleophilic water surrogate that enables anti-Markovnikov hydration of styrene derivatives.

## Results and discussion


Base-catalyzed alcohol addition reactions to styrene derivatives are reversible such that the substrate identities and reaction conditions contribute to the observed equilibrium yields. These considerations and the identification of effective catalysts are extensively documented in a recent report from our group that served as the basis for this work.<sup>15b,d</sup> We began by examining  $\text{P}_4\text{-}t\text{-Bu}$ -catalyzed addition reactions using oxygen pronucleophiles that could potentially serve as water surrogates.<sup>18</sup> These reactions were conducted under our previously optimized reaction conditions with 2-chloro-3-vinylpyridine (**1**) as a model alkene (Scheme 1a).<sup>15d</sup> Alcohols comprised of common *O*-protecting groups, such as trimethylsilanol, phenol, *p*-methoxybenzyl alcohol and allyl alcohol do not undergo addition. These results are consistent with our previous studies that revealed alcohol addition to styrenes is both kinetically and thermodynamically challenging, indicating a more nucleophilic water surrogate is necessary for addition.<sup>15,19</sup> However, alcohols comprised of common aliphatic protecting groups, such as *tert*-butanol and 2-trimethylsilylethanol, provide low addition yields. Examination of cyclopropyl-substituted alcohols led to the discovery that 1-cyclopropylethanol undergoes addition in 82% isolated yield. The resulting 1-methyl 1'-cyclopropylmethyl (MCPM) ether is an uncommon protecting group developed for oligosaccharide synthesis.<sup>20</sup> For comparison, cyclopropylmethanol, dicyclopropylmethanol and *iso*-propanol all provide lower addition yields. The isolated MCPM ether **2** undergoes high-yielding deprotection in 3 min using methanesulfonic acid, presumably through a cyclopropyl-stabilized carbocation hydrolysis mechanism (Scheme 1b).<sup>21</sup>

We next investigated the substrate scope for anti-Markovnikov hydration using 1-cyclopropylethanol (**4**) and an

### (a) Identification of a nucleophilic water surrogate reagent<sup>a</sup>



### (b) Rapid acidic hydrolysis of 1-cyclopropylethyl ether adduct



Scheme 1 Optimization studies for (a) superbase-catalyzed alcohol addition and (b) ether hydrolysis. <sup>a</sup> Yields determined by <sup>1</sup>H NMR spectroscopy. <sup>b</sup> Isolated yield of purified product.

*in situ* acidic workup procedure (Table 1). First, in Table 1a, a comparison of superbase ( $\text{P}_4\text{-}t\text{-Bu}$  in PhMe) and inorganic ( $\text{KO-}t\text{-Bu}$ /18-crown-6 in DME) catalysts is shown for two styrene derivatives (**5** and **6**) and two chlorinated vinyl pyridines (**3** and **7**). The inorganic conditions were selected based on our prior extensive screening of bases for hydroetherification reactions,<sup>15b</sup> with additional information provided in the ESI.† As seen in Table 1a, the inorganic conditions enable synthetically useful alcohol yields (49–61%) while the use of  $\text{P}_4\text{-}t\text{-Bu}$  generally provides higher yields (60–83%). The yield differences between the superbase and inorganic conditions are likely a solvent equilibrium effect, as we previously measured alcohol addition to be more favorable in PhMe than in DME.<sup>15b</sup> While  $\text{P}_4\text{-}t\text{-Bu}$  is an efficient catalyst in both solvents, the  $\text{KO-}t\text{-Bu}$ /18-crown-6 system is most active in DME and typically does not catalyze the reaction in PhMe to its equilibrium position.<sup>15d</sup>

Table 1b shows hydration products of other vinyl arenes using the superbase conditions, while a full comparison to inorganic conditions is provided in the ESI.† Styrenes (**5**, **6**, **8–15**) featuring diverse substitution patterns with nitro, fluoro, chloro, bromo, sulfonyl, trifluoromethyl and benzoyl substituents undergo hydration in good yields. A notable feature of this scope is that *ortho*-substituents are well tolerated. Vinyl heteroarenes are also effective substrates, including those with halogens prone to nucleophilic aromatic substitution. Thus, halogenated 2-, 3-, and 4-vinyl pyridine derivatives (**3**, **7**, **17** and **18**) undergo hydration in high yields. Beyond pyridines, quinoline (**16**) and thiazole (**19**) derivatives are also hydrated in good yields. Although 1-cyclopropylethanol (**4**) will add to electron-neutral styrenes and 1,1-

Table 1 Product scope for anti-Markovnikov hydration<sup>a</sup>

**(a) Yields obtained using superbases and inorganic conditions**

*superbase* = P<sub>4</sub>-t-Bu in PhMe    *inorganic* = KO-t-Bu + 18-crown-6 in DME

 <b>5</b> <i>superbase</i> 40 °C, 60% yield <i>inorganic</i> 60 °C, 49% yield	 <b>3</b> <i>superbase</i> 50 °C, 83% yield <i>inorganic</i> 40 °C, 61% yield <sup>b</sup>
 <b>6</b> <i>superbase</i> rt, 64% yield <sup>b</sup> <i>inorganic</i> rt, 61% yield <sup>b</sup>	 <b>7</b> <i>superbase</i> 40 °C, 65% yield <i>inorganic</i> 40 °C, 51% yield

**(b) Additional substrate examples using superbase conditions**

 <b>8</b> , 30 °C 64% yield	 <b>9</b> , 60 °C 48% yield	 <b>10</b> , 100 °C 81% yield
 <b>11</b> , 40 °C 77% yield	 <b>12</b> , 65 °C 83% yield	 <b>13</b> , 40 °C 49% yield <sup>b</sup>
 <b>14</b> , 50 °C 75% yield	 <b>15</b> , 60 °C 44% yield	 <b>16</b> , 40 °C 62% yield
 <b>17</b> , rt 75% yield <sup>c</sup>	 <b>18</b> , 35 °C 60% yield	 <b>19</b> , rt 74% yield <sup>d</sup>

<sup>a</sup> Yields are of purified alcohol product; reactions performed on 0.5 to 1.0 mmol scale. <sup>b</sup> 5 equiv. of **4** used. <sup>c</sup> 1 h reaction time. <sup>d</sup> 4 equiv. of **4** used.


disubstituted variants, only low equilibrium yields are observed under these conditions.<sup>15b</sup> Representative substrates documenting the electronic limitations of this method are shown in the ESI.†

We next compared the hydration yields of several substrates from Table 1 to traditional hydroboration/oxidation protocols using BH<sub>3</sub>·THF and 9-BBN (Scheme 2a). Although use of these boranes results in high yields of β-phenethyl alcohols from electron-neutral styrenes, decreased yields and selectivities are common for electron-deficient and heteroaryl variants.<sup>8</sup> Hydroboration/oxidation of 2,6-dichlorostyrene (**20**) leads to mixtures of α- and β-phenethyl alcohols while 2-nitro-4-methylstyrene (**21**) results in low yields of β-phenethyl alcohol. An even greater limitation is observed for 3-bromo-4-vinylpyridine (**22**), in which only net hydrogenation occurs with no alcohol formation. In contrast, base-catalyzed hydration gives high yields with exclusive β-regioselectivity for these substrates.

**(a) Substrate scope comparison to traditional hydroboration/oxidation<sup>a</sup>**


Ar-CH=CH <sub>2</sub> (1 equiv)	+ 4 (3 equiv)	$\xrightarrow[\text{solvent, 12-24 h}]{\text{catalyst (10 mol\%)}}$ (HCl or MsOH workup)	Ar-CH <sub>2</sub> -CH <sub>2</sub> -OH	
			β-alcohol	α-alcohol
			$\xrightarrow[\text{THF, 0 °C to rt or 80 °C, 14 h}]{\text{i) BH}_3\cdot\text{THF or 9-BBN}}$	Ar-CH <sub>2</sub> -CH <sub>2</sub> -OH + Ar-CH(OH)-Me
			$\xrightarrow[\text{MeOH, 0 °C to rt, 4 h}]{\text{ii) H}_2\text{O}_2, \text{NaOH}}$	β-alcohol + α-alcohol

	<b>20</b>	<b>21</b>	<b>22</b>
BH <sub>3</sub> ·THF	15% β + 34% α	39% β + 0% α	0% β + 0% α
9-BBN	63% β + 11% α	18% β + 0% α	0% β + 0% α
from Table 1	81% β + 0% α	64% β + 0% α	75% β + 0% α

**(b) Examples of hydroboration/oxidation-sensitive functional groups<sup>b</sup>**


Scheme 2 Comparison of base-catalyzed hydration to hydroboration/oxidation methods. <sup>a</sup> Yields and regioselectivities determined by <sup>1</sup>H NMR spectroscopy of the crude reaction mixture; yields from Table 1 represent isolated yields. <sup>b</sup> Yields are of purified alcohol product. <sup>c</sup> 5 equiv. of **4** used.

Base-catalyzed hydration is also likely to have complementary functional group tolerance to hydroboration/oxidation protocols. This is illustrated in Scheme 2b, where substrates with an aryl thioether (**23**) and a thiomorpholine (**24**) that are sensitive to oxidizing conditions undergo P<sub>4</sub>-t-Bu-catalyzed hydration in moderate yields.<sup>22</sup> Substrate **25**, featuring a terminal alkene that is reactive toward traditional hydroboration, illustrates the chemoselectivity of this method for styrene hydration.<sup>23</sup>

**10 to 30 mmol scale anti-Markovnikov hydration reactions**

Scheme 3 Examples of preparative scale anti-Markovnikov hydration reactions; yields are of isolated alcohol product.





**Scheme 4** Utility of Table 1 alcohol products for dihydrobenzofuran derivative synthesis; yields are of isolated product from the cyclization step using given conditions. <sup>a</sup> CuI (10 mol%), 8-quinolinol (15 mol%), Cs<sub>2</sub>CO<sub>3</sub> (1.5 equiv.), PhMe, 110 °C, 16 h. <sup>b</sup> NaH (1.5 equiv.), THF, 70 °C, 16 h. <sup>c</sup> Pd(OAc)<sub>2</sub> (3 mol%), JohnPhos (4 mol%), Cs<sub>2</sub>CO<sub>3</sub> (1.5 equiv.), PhMe, 80 °C, 22 h.

Scheme 3 demonstrates the scalability of this protocol through three hydration reactions conducted on 10 mmol scale or greater. Hydration of 3-bromo-4-vinylpyridine (**22**) is best accomplished using P<sub>4</sub>-t-Bu and thus 5 mol% catalyst loading provided access to 1.4 grams of alcohol **17**.<sup>24</sup> The hydration of 2,6-dichlorostyrene (**20**, 30 mmol) was achieved in 73% yield using catalytic KO-t-Bu, exemplifying the practicality of the inorganic conditions. Although the highest styrene hydration yields are obtained using 3 equiv. of 1-cyclopropylethanol (**4**), we also identified conditions that allow for only moderately decreased addition yields using 1.2 equiv. of **4**. This finding, shown in Table S3† of the ESI,† exploits our previous discovery of a negative alcohol rate order for base-catalyzed hydro-etherification reactions.<sup>15b</sup> Thus, reactions can be run at lower temperatures when less alcohol is used, thereby decreasing the entropic penalty of addition to counteract Le Chatelier's principle. This effect was extended to a challenging oxa-Michael addition process for acrylamide **26**, where just 1.5 equiv. of **4** and 30 min are required for a high-yielding KO-t-Bu-catalyzed hydration process at room temperature.<sup>25</sup>

Given that this method is suitable for *ortho*-halogenated vinyl (hetero)arenes, we reasoned hydration could be sequenced with a cyclization reaction to prepare 2,3-dihydrobenzofuran derivatives.<sup>26</sup> Thus, employing substrates from Table 1, base-promoted or metal-catalyzed cyclization reactions produce **28–32** in high yields (Scheme 4). This approach should improve access to this important heterocycle class,<sup>27</sup> especially functionalized and aza variants where there is limited availability of the corresponding benzofuran compounds.<sup>28</sup>

## Conclusions

In conclusion, base-catalyzed addition of 1-cyclopropylethanol to styrene derivatives provides a convenient method for preparing β-aryl substituted alcohols. The use of catalytic P<sub>4</sub>-t-Bu superbase allows for superior yields to inorganic bases, although KO-t-Bu/18-crown-6 may be a more practical catalyst system for hydration on larger scale. In the broader context,

analogous application of the nucleophilic yet easily deprotected 1-cyclopropylethanol water surrogate can likely enable other challenging hydration processes.

## Data availability

Compound characterization data and experimental procedures are openly available in the ESI.†

## Author contributions

S. P. P., Z. Q. and S. J. S. performed the experimental work. The manuscript was written by J. S. B. with input from all authors.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This material is based upon work supported by the National Science Foundation under Grant No. 1944478. We also thank Colorado State University (CSU) for funding and the Analytical Resources Core (RRID: SCR\_021758) at CSU for instrument access, training and assistance with sample analysis.

## Notes and references

- For references that discuss compounds shown in Fig. 1a, see: (a) S. Manzetti, J. Zhang and D. van der Spoel, *Biochem.*, 2014, **53**, 821–835; (b) R.-H. Guo, Q. Zhang, Y.-B. Ma, J. Luo, C.-A. Geng, L.-J. Wang, X.-M. Zhang, J. Zhou, Z.-Y. Jiang and J.-J. Chen, *Eur. J. Med. Chem.*, 2011, **46**, 307–319; (c) H. R. Dexter, E. Allen and D. M. Williams, *Tetrahedron Lett.*, 2018, **59**, 4323–4325.
- For references that discuss compounds shown in Fig. 1b, see: (a) H. Chen, Y. Chen, Q. Zou and L. Yuan, *Org. Process Res. Dev.*, 2013, **17**, 714–717; (b) L. R. Madivada, R. R. Anumala, G. Gilla, S. Alla, K. Charagondla, M. Kagga, A. Bhattacharya and R. Bandichhor, *Org. Process Res. Dev.*, 2009, **13**, 1190–1194.
- For references that discuss compounds shown in Fig. 1c, see: (a) J. Xie, F. Yang, M. Zhang, C. Lam, Y. Qiao, J. Xiao, D. Zhang, Y. Ge, L. Fu and D. Xie, *Bioorg. Med. Chem. Lett.*, 2017, **27**, 131–134; (b) J. T. Sanderson, H. Clabault, C. Patton, G. Lassalle-Claux, J. Jean-François, A. F. Paré, M. J. G. Hébert, M. E. Surette and M. Touaibia, *Bioorg. Med. Chem.*, 2013, **21**, 7182–7193; (c) L. A. T. Cleghorn, P. C. Ray, J. Odingo, A. Kumar, H. Wescott, A. Korkegian, T. Masquelin, A. Lopez Moure, C. Wilson, S. Davis, M. Huggett, P. Turner, A. Smith, O. Epemolu, F. Zuccotto, J. Riley, P. Scullion, Y. Shishikura, L. Ferguson, J. Rullas, L. Guijarro, K. D. Read, S. R. Green, P. Hipskind, T. Parish and P. G. Wyatt, *J. Med. Chem.*, 2018, **61**, 6592–6608; (d) D. E. Jeffries, J. O. Witt, A. L. McCollum, K. J. Temple, M. A. Hurtado, J. M. Harp, A. L. Blobaum, C. W. Lindsley

- and C. R. Hopkins, *Bioorg. Med. Chem. Lett.*, 2016, **26**, 5757–5764.
- 4 (a) M. B. Smith, *March's Advanced Organic Chemistry*, 7th edn, Wiley, New York, 2013, pp. 1497–1519; (b) S. Werkmeister, K. Junge and M. Beller, *Org. Process Res. Dev.*, 2014, **18**, 289–302; (c) J. Pritchard, G. A. Filonenko, R. van Putten, E. J. M. Hensen and E. A. Pidko, *Chem. Soc. Rev.*, 2015, **44**, 3808–3833.
- 5 For reviews on uncatalyzed hydroboration, see: (a) H. C. Brown, *Hydroboration*, W. A. Benjamin, Inc., New York, 1962; (b) R. S. Dhilon, *Hydroboration and Organic Synthesis*, Springer, Berlin, Germany, 2007; (c) H. C. Brown, *Tetrahedron*, 1961, **12**, 117–138.
- 6 A variety of alternative routes have been employed toward  $\beta$ -aryl alcohols; for representative examples, see: (a) S. Mohanty, S. Talasila, A. K. Roy and A. C. Karmakar, *Org. Process Res. Dev.*, 2014, **18**, 168–173; (b) C. Huynh, F. Derguini-Boumechal and G. Lindstrumelle, *Tetrahedron Lett.*, 1979, **20**, 1503–1506; (c) J. V. Obligation and P. J. Chirik, *Nat. Rev. Chem.*, 2018, **2**, 15–34; (d) M. D. Greenhalgh, D. J. Frank and S. P. Thomas, *Adv. Synth. Catal.*, 2014, **356**, 584–590; (e) M. Zeng, L. Li and S. B. Herzon, *J. Am. Chem. Soc.*, 2014, **136**, 7058–7067; (f) T. Zhang, Z. Zhang, Y. Nishiyama and H. Maekawa, *Tetrahedron*, 2016, **72**, 2293–2299.
- 7 For example routes to arylacetic acid derivatives, see the following references and references therein: (a) M. R. Biscoe and S. L. Buchwald, *Org. Lett.*, 2009, **11**, 1773–1775; (b) T. Hama, S. Ge and J. F. Hartwig, *J. Org. Chem.*, 2013, **78**, 8250–8266; (c) P. J. Moon, S. Yin and R. J. Lundgren, *J. Am. Chem. Soc.*, 2016, **138**, 13826–13829; (d) P. Xie, Y. Xie, B. Qian, H. Zhou, C. Xia and H. Huang, *J. Am. Chem. Soc.*, 2012, **134**, 9902–9905; (e) Q.-Y. Meng, T. E. Schirmer, A. L. Berger, K. Donabauer and B. König, *J. Am. Chem. Soc.*, 2019, **141**, 11393–11397.
- 8 For discussions of stoichiometric hydroboration on challenging styrene substrates, see: (a) H. C. Brown, J. V. N. Vara Prasad and S. H. Zee, *J. Org. Chem.*, 1986, **51**, 439–445; (b) H. C. Brown and R. L. Sharp, *J. Am. Chem. Soc.*, 1966, **88**, 5851–5854; (c) L. C. Vishwakarma and A. Fry, *J. Org. Chem.*, 1980, **45**, 5306–5308; (d) P. V. Ramachandran, S. Madhi and M. J. O'Donnell, *J. Fluorine Chem.*, 2006, **127**, 1252–1255.
- 9 For reviews and examples of metal-catalyzed and other approaches to hydroboration, see: (a) K. Semba, T. Fujihara, J. Terao and Y. Tsuji, *Tetrahedron*, 2015, **71**, 2183–2197; (b) K. Burgess and M. J. Ohlmeyer, *Chem. Rev.*, 1991, **91**, 1179–1191; (c) I. Beletskaya and A. Pelter, *Tetrahedron*, 1997, **53**, 4957–5026; (d) J. H. Docherty, J. Peng, A. P. Dominey and S. P. Thomas, *Nat. Chem.*, 2017, **9**, 595–600; (e) R. Corberán, N. W. Mszar and A. H. Hoveyda, *Angew. Chem., Int. Ed.*, 2011, **50**, 7079–7082; (f) J. V. Obligation and P. J. Chirik, *J. Am. Chem. Soc.*, 2013, **135**, 19107–19110; (g) Z. Kuang, K. Yang, Y. Zhou and Q. Song, *Chem. Commun.*, 2020, **56**, 6469–6479; (h) T. Taniguchi, *Eur. J. Org. Chem.*, 2019, 6308–6319.
- 10 An alternative oxidation/reduction route to terminal alcohols is the regioselective reductive opening of styrene oxide compounds. For selected recent examples, see: (a) C. Yao, T. Dahmen, A. Gansäuer and J. Norton, *Science*, 2019, **364**, 764–767; (b) X. Liu, L. Longwitz, B. Spiegelberg, J. Tönjes, T. Beweries and T. Werner, *ACS Catal.*, 2020, **10**, 13659–13667; (c) M. Magre, E. Paffenholz, B. Maity, L. Cavallo and M. Rueping, *J. Am. Chem. Soc.*, 2020, **142**, 14286–14294; (d) W. Liu, W. Li, A. Spannenberg, K. Junge and M. Beller, *Nat. Catal.*, 2019, **2**, 523–528.
- 11 For reviews on anti-Markovnikov hydration and oxidation of alkenes, see: (a) J. Guo and P. Teo, *Dalton Trans.*, 2014, **43**, 6952–6964; (b) M. Beller, J. Seayad, A. Tillack and H. Jiao, *Angew. Chem., Int. Ed.*, 2004, **43**, 3368–3398; (c) J. J. Dong, W. R. Browne and B. L. Feringa, *Angew. Chem., Int. Ed.*, 2015, **54**, 734–744. For two recent examples of anti-Markovnikov addition of oxygen reagents to terminal alkenes, see: (d) S.-Q. Lai, B.-Y. Wei, J.-W. Wang, W. Yu and B. Han, *Angew. Chem., Int. Ed.*, 2021, **60**, 21997–22003; (e) L. Quach, S. Dutta, P. M. Pflüger, F. Sandfort, P. Bellotti and F. Glorius, *ACS Catal.*, 2022, **12**, 2499–2504.
- 12 G. Dong, P. Teo, Z. K. Wickens and R. H. Grubbs, *Science*, 2011, **333**, 1609–1612.
- 13 (a) S. C. Hammer, G. Kubik, E. Watkins, S. Huang, H. Minges and F. H. Arnold, *Science*, 2017, **358**, 215–218; (b) S. Wu, J. Liu and Z. Li, *ACS Catal.*, 2017, **7**, 5225–5233; (c) For a review, see: S. Wu, Y. Zhou and Z. Li, *Chem. Commun.*, 2019, **55**, 883–896.
- 14 (a) X. Hu, G. Zhang, F. Bu and A. Lei, *ACS Catal.*, 2017, **7**, 1432–1437; (b) Y. Chen, J. Zhang, Z. Tang and Y. Sun, *J. Photochem. Photobiol. A*, 2020, **392**, 112340.
- 15 (a) C. Luo and J. S. Bandar, *J. Am. Chem. Soc.*, 2018, **140**, 3547–3550; (b) C. Luo, J. V. Alegra-Requena, S. J. Sujansky, S. P. Pajk, L. C. Gallegos, R. S. Paton and J. S. Bandar, *J. Am. Chem. Soc.*, 2022, **144**, 9586–9596; (c) C. Luo and J. S. Bandar, *Synlett*, 2018, **29**, 2218–2224; (d) Additional information regarding catalyst and condition optimization for both the superbases and inorganic catalyst systems can be found in the ESI.†
- 16 For reviews and references on  $P_4$ -*t*-Bu superbases catalysis, see: (a) R. Schwesinger and H. Schlemper, *Angew. Chem., Int. Ed.*, 1987, **26**, 1167–1169; (b) *Superbases for Organic Synthesis: Guanidines, Amidines, Phosphazenes and Related Organocatalysts*, ed. T. Ishikawa, John Wiley & Sons, Ltd: Chichester, UK, 2009; (c) T. R. Puleo, S. J. Sujansky, S. E. Wright and J. S. Bandar, *Chem. - Eur. J.*, 2021, **27**, 4216–4229.
- 17 For an example of addition of hydroxide to a highly activated vinyl *N*-heteroarene, see: J. M. Bartolomé-Nebreda, S. A. Alonso de Diego, M. Artola, F. Delgado, O. Delgado, M. L. Martín-Martín, C. M. Martínez-Viturro, M. A. Pena, H. M. Tong, M. Van Gool, J. M. Alonso, A. Fontana, G. J. Macdonald, A. Megens, X. Langlois, M. Somers, G. Vanhoof and S. Conde-Ceide, *J. Med. Chem.*, 2015, **58**, 978–993.
- 18 P. G. M. Wuts, *Greene's Protective Groups in Organic Synthesis*, 5th edn, Wiley, Hoboken, NJ, 2014.

- 19 T. R. Puleo, A. J. Strong and J. S. Bandar, *J. Am. Chem. Soc.*, 2019, **141**, 1467–1472.
- 20 (a) E. Eichler, F. Yan, J. Sealy and D. M. Whitfield, *Tetrahedron*, 2001, **57**, 6679–6693; (b) S. H. Yu and D. M. Whitfield, *Tetrahedron*, 2011, **67**, 5750–5761.
- 21 (a) G. A. Olah, V. P. Reddy and G. K. S. Prakash, *Chem. Rev.*, 1992, **92**, 69–95; (b) C. F. Wilcox, L. M. Loew and R. Hoffman, *J. Am. Chem. Soc.*, 1973, **95**, 8192–8193.
- 22 M. Madesclaire, *Tetrahedron*, 1986, **42**, 5459–5495.
- 23 For an example of selective hydroboration of an unactivated alkene in the presence of a styrene, see: J. Y. Dong, E. Manias and T. C. Chung, *Macromolecules*, 2002, **35**, 3439–3447.
- 24 We found that  $P_4-t\text{-Bu}$  catalyzes the addition of alcohol **4** to alkene **22** in 95% yield whereas the inorganic catalyst system provides a maximum 68% addition yield; see ESI† for details.
- 25 For a discussion of the challenges of oxa-Michael addition and hydration reactions, see: (a) C. F. Nising and S. Bräse, *Chem. Soc. Rev.*, 2012, **41**, 988–999; (b) E. Hartmann, D. J. Vyas and M. Oestreich, *Chem. Commun.*, 2011, **47**, 7917–7932; (c) K. Zheng, X. Liu and X. Feng, *Chem. Rev.*, 2018, **118**, 7586–7656.
- 26 For example other routes to compounds similar to those in Scheme 4, see: (a) S. El Sayed, A. Bordet, C. Weidenthaler, W. Hetaba, K. L. Luska and W. Leitner, *ACS Catal.*, 2020, **10**, 2124–2130; and references therein.; (b) S. Kuwabe, K. E. Torraca and S. L. Buchwald, *J. Am. Chem. Soc.*, 2001, **123**, 12202–12206; (c) J. Alvarado, J. Fournier and A. Zakarian, *Angew. Chem., Int. Ed.*, 2016, **55**, 11625–11628; (d) A. E. Frissen, A. T. M. Marcelis and H. C. Van Der Plas, *Tetrahedron*, 1989, **45**, 803–812; (e) R. Willand-Charnley, B. W. Puffer and P. H. Dussault, *J. Am. Chem. Soc.*, 2014, **136**, 5821–5823.
- 27 For examples of bioactive dihydrobenzofurans, see: (a) R. J. Nevagi, S. N. Dighe and S. N. Dighe, *Eur. J. Med. Chem.*, 2015, **97**, 561–581; (b) D. P. Zlotos, R. Jockers, E. Cecon, S. Rivara and P. A. Witt-Enderby, *J. Med. Chem.*, 2014, **57**, 3161–3185; (c) Z. Chen, M. Pitchakuntla and Y. Jia, *Nat. Prod. Rep.*, 2019, **36**, 666–690.
- 28 The corresponding benzofurans of products in Scheme 4 are expensive (>\$500/gram) or not commercially available according to a search of the online chemical catalog, <https://www.emolecules.com>, accessed May 17, 2022.