

The Membrane-Associated Transient Receptor Potential Vanilloid Channel Is the Central Heat Shock Receptor Controlling the Cellular Heat Shock Response in Epithelial Cells

Zohar Bromberg¹, Pierre Goloubinoff², Younousse Saidi², Yoram George Weiss^{1,3*}

1 Dept. of Anesthesiology and Critical Care Medicine and the Goldyne Savad Institute of Gene Therapy, Hadassah-Hebrew University School of Medicine, Jerusalem, Israel, **2** Dept. of Plant Molecular Biology, University of Lausanne, Lausanne, Switzerland, **3** Dept. of Anesthesiology and Critical Care Medicine, University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania, United States of America

Abstract

The heat shock response (HSR) is a highly conserved molecular response to various types of stresses, including heat shock, during which heat-shock proteins (Hsps) are produced to prevent and repair damages in labile proteins and membranes. In cells, protein unfolding in the cytoplasm is thought to directly enable the activation of the heat shock factor 1 (HSF-1), however, recent work supports the activation of the HSR via an increase in the fluidity of specific membrane domains, leading to activation of heat-shock genes. Our findings support the existence of a plasma membrane-dependent mechanism of HSF-1 activation in animal cells, which is initiated by a membrane-associated transient receptor potential vanilloid receptor (TRPV). We found in various non-cancerous and cancerous mammalian epithelial cells that the TRPV1 agonists, capsaicin and resiniferatoxin (RTX), upregulated the accumulation of Hsp70, Hsp90 and Hsp27 and Hsp70 and Hsp90 respectively, while the TRPV1 antagonists, capsazepine and AMG-9810, attenuated the accumulation of Hsp70, Hsp90 and Hsp27 and Hsp70, Hsp90, respectively. Capsaicin was also shown to activate HSF-1. These findings suggest that heat-sensing and signaling in mammalian cells is dependent on TRPV channels in the plasma membrane. Thus, TRPV channels may be important drug targets to inhibit or restore the cellular stress response in diseases with defective cellular proteins, such as cancer, inflammation and aging.

Citation: Bromberg Z, Goloubinoff P, Saidi Y, Weiss YG (2013) The Membrane-Associated Transient Receptor Potential Vanilloid Channel Is the Central Heat Shock Receptor Controlling the Cellular Heat Shock Response in Epithelial Cells. PLoS ONE 8(2): e57149. doi:10.1371/journal.pone.0057149

Editor: Sue Cotterill, St. Georges University of London, United Kingdom

Received: December 8, 2011; **Accepted:** January 22, 2013; **Published:** February 27, 2013

Copyright: © 2013 Bromberg et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported in part by grants from the Israel Science Foundation (#529/08 - YGW), the ESICM (European Society of Intensive Care Medicine) Basic Science Award, the Hadassah Medical Organization internal research grant (YGW), the Hadassah Medical Organization internal research grant (YGW) and the Zwahlen foundation from Lausanne University (PG). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: weiss@hadassah.org.il

Introduction

The heat shock response (HSR), which is induced upon exposure of living cells to acute or subacute stressors, is characterized by the expression of a group of phylogenetically-conserved intracellular heat shock proteins (HSPs), which have a protective effect against injury, especially of proteins [1]. Among the most massively expressed HSPs are molecular chaperones that mediate conformational changes in proteins and prevent protein misfolding. Some heat-induced chaperones, such as Hsp70, can hydrolyze ATP to unfold toxic protein aggregates into natively refoldable polypeptides and promote the degradation of toxic misfolded proteins by the proteasome [2–5]. Further, HSP chaperones modulate apoptosis, NF- κ B signaling, cell division, and other pathways associated with the cellular response to stress, resulting in cytoprotection from various environmental stresses, heat shock, and infection-induced pro-inflammatory pathways [1,2,6–9]. Hsp70 is a core element of the chaperone network in the cell that controls stress signaling, protein trafficking and the onset of damage-preventing and repairing mechanisms [3,10]. Hsp70

functions in collaboration with chaperones and co-chaperones such as Hsp90, Hsp110, Hsp27, Hsp40, etc. [1–3]. Altered levels of HSP chaperones are often observed in inflammation, as in severe sepsis [11]. The HSR also provides cytoprotection from chemical aggressors, such as chemotherapy [12]. Hence, cancer cells that constitutively up-regulate the expression of HSPs, Hsp70 and Hsp90 in particular, are often resistant to heat as well as to chemotherapeutic treatments [12]. Reciprocally, inhibition of Hsp90 using specific inhibitors such as geldanamycin or radicicol, or reducing Hsp70 expression using siRNA, leads to growth arrest and cell death [13–16].

Heat shock factors (HSFs), in particular HSF-1, control the transcription of HSPs [4]. In the absence of stress, inactive HSF-1 monomers are hypo-phosphorylated and are apparently associated in the cytoplasm with Hsp70 and Hsp90 [17]. One mechanism by which diverse stress stimuli, heat-shock or toxic chemicals in particular, are proposed to induce the HSR, is the presumed stress-induced formation of misfolded proteins in the cytoplasm, which would in turn titrate Hsp70 and Hsp90 chaperones away from their stable inhibitory association with HSF-1, thereby,




Figure 1. Capsaicin and capsazepine effects on heat shock protein expression in HEK293e, MLE-12, MCF-7 and HT-29 cells. Representative immunoblot analysis of Hsp70, Hsp90 and Hsp27 abundances. Hsp70, Hsp90 and Hsp27 were detected in HEK293e (Panel A), MLE-12 cells (Panel B), HT-29 (Panel C) and MCF-7 (Panel D). Cells in culture were incubated with 0.01% DMSO, 32 μ M of capsaicin (Caps) for 1 hr, 32 μ M of capsaicin (Caps) for 1 hr with the addition of 100 μ M capsazepine (CPZ) or 100 μ M capsazepine alone at 37°C. Cells were treated with either heat shock at 42°C for 3 hrs, heat shock at 42°C for 3 hrs with the addition of 100 μ M capsazepine. Cells were treated with 32 μ M of capsaicin with the addition of 5 mM EGTA, 5 mM EGTA alone at 37°C or heat shock at 42°C for 3 hrs with the addition of 5 mM EGTA. 15 μ g of total protein/lane were separated on 9% SDS-PAGE gel.

doi:10.1371/journal.pone.0057149.g001

presumably, resulting in HSF-1 activation by way of hyperphosphorylation, trimerization and translocation to the nucleus [4,18–20]. However, this mechanism fails to explain situations where HSP levels are increased in the presence of non-denaturing chemicals and without heat under physiological conditions, where protein aggregates are unlikely to form in the cell. Thus, a strong HSR may already develop in response to a very mild temperature increase or in the presence of membrane-interfering compounds such as arimoclomol, benzyl alcohol, NG-094, BGP-15 [21–26]. A different mechanism for heat-sensing and for activation by heat or chemicals of the HSR has been demonstrated whereby activation of the HSR occurs via increase in the fluidity of specific membrane microdomains, leading to activation of heat-shock genes [22]. This

activation of the HSR occurs, not only by heat, but may also happen in response to chemicals, such as bimocmolol sterol glycoside [27], acetyl salicylic acid or arachidonic acid [25]. The activation occurs through subsequent secondary messengers, such as calmodulin-binding domains, which are likely to associate with entering Ca^{2+} -ions and specific calmodulins in the cytoplasm [27], sterol glycoside [27] or arachidonic acid [25], thus providing cells with a means to mediate a highly specific Heat Shock-like signal without temperature changes.

Here, we provide new supportive evidence to the seminal findings that binding of HSF-1 to HSE can be activated by the entry of extracellular Ca^{2+} also in mammalian cells [28]. This work identifies specific Ca^{2+} -channels as components of a central

cellular thermosensory apparatus in mammalian cells. We show that mammalian epithelial cells contain specific Ca^{2+} -channels, the membrane-dependent transient receptor potential vanilloid (TRPV) that can respond primarily to a temperature rise and also to chemicals such as capsaicin, by sending a Ca^{2+} -entry dependent signal to activate HSF-1 and develop a cellular HSR. We found in various non-cancerous and cancerous mammalian epithelial cells that the TRPV1 agonists, capsaicin and resiniferatoxin (RTX) [29], upregulated the accumulation of Hsp70, Hsp90 and Hsp27 and Hsp70 and Hsp90, respectively, while the TRPV1 antagonists, capsazepine, and AMG-9810 [30] attenuated the accumulation of Hsp70, Hsp90 and Hsp27 and Hsp70, Hsp90, respectively. Capsaicin was also shown to activate HSF-1. TRPV1 siRNA, capsazepine, or AMG-9810, or extracellular EGTA all prevented the heat-induced accumulation of Hsp70 and Hsp90, suggesting that in mammalian cells, heat-sensing and cellular stress signaling in general depend on specific membrane-dependent transient receptor potential vanilloid (TRPV) calcium channel-like receptors. Together, our findings suggest that heat or specific chemicals can change the plasma membrane state [22], thus triggering the transient opening of calcium channel-like receptors (TRPVs), which in turn send a specific signal to generate a cellular stress response, under conditions that are not necessarily dependent on stress damages in proteins, in particular the formation of intra-cellular protein aggregates.

Results

Capsaicin induces, and capsazepine inhibits HSP expression in non-neuronal epithelial cells

Western blot analysis showed that when kidney cells (HEK-293e found to express TRPV1), alveolar epithelial cells (MLE-12), carcinoma breast cancer cell line (MCF-7) or colon cancer cells (HT-29) are treated with the TRPV1 agonist capsaicin, they accumulate Hsp70, Hsp90 and Hsp27 proteins (**Fig. 1**). Capsazepine, a specific antagonist of capsaicin [28,29], abolished the heat-induced HSR (**Fig. 1**). Capsaicin and capsazepine were found to respectively activate, or inhibit the HSR. Similarly to capsazepine, treatment with the strong calcium chelator EGTA abolished HSP overexpression by capsaicin or by heat (**Fig. 1**). Furthermore, exposure of HEK-293e cells to 32 μM capsaicin, at different times (0, 1 hr, 4 hrs, and 6 hrs) demonstrated significantly elevated HSP levels after 1 hr (**Fig. 2C**).

A 3-hour heat-shock at 42°C or an isothermal treatment with capsaicin, both induced the accumulation of HSPs, which was directly associated with an increase of HSF-1 and Phospho-HSF-1 abundance and, expectedly, with HSF-1 activation, as determined also by DNA-shift binding activity in HEK-293e cells (**Fig. 3**). Confirming the isothermal induction we show that in cells harboring TRPV1, either heat-shock or treatment with capsaicin induces an increase of HSF-1 and Phospho-HSF-1 abundances, a finding confirmed by semi quantitative PCR depicting an increase of HSF-1 mRNA (**Fig. 3**). The addition of capsazepine prior to HS impaired the heat-dependent induction of HSF-1 (**Fig. 3B**). Capsazepine-suppression of heat-induced HSP accumulation was directly associated with the inhibition of HSF-1 abundance and with its reduced DNA binding activity (**Fig. 3C**). Treatment of cells with both capsaicin and its competitive inhibitor capsazepine, resulted in a partial inhibition of the capsaicin-induced HSR (**Fig. 1 & 3**).

Other TRPV1 agonists and antagonists, RTX and AMG-9810, respectively, were also tested. RTX dose response was clearly associated with increased Hsp90, and Hsp70 expressions, while AMG-9810 blocked these effects (**Fig. 4B**). The RTX findings

correlate with the dose response findings of capsaicin (**Fig. 4A**). Doses in the range of 4-32 μM of capsaicin showed elevated levels of Hsp70, Hsp90 and Hsp27 in a dose-dependent manner in epithelial cell cultures (**Fig. 4A**).

The role of TRPV in the HSR

In our work, the acute cellular and physiologic effects of heat or capsaicin are principally mediated by the transient receptor potential vanilloid type-1 (TRPV1), a non-selective membrane calcium channel, responsive to noxious temperatures or to vanilloid compounds providing an explanation for the previously unnoted inhibitory effect of EGTA on the heat shock response. The presence of TRPV1 has been previously detected in a wide range of neuronal and non-neuronal cells [31,32]. In contrast to previous suggestions [33,34], we found by Western blots the presence of TRPV1 receptors also in HEK-293e cells (**Fig. 2B**). A capsaicin dose-response experiment in HEK-293e cells depicted an increased TRPV1 abundance (**Fig. 2B**). This was also depicted by immuno-histochemistry which showed TRPV1 abundance in capsaicin-treated HEK-293e cells (**Fig. 5**). To confirm these findings, we performed RT-PCR for TRPV1 (**Fig. 2A**). We found TRPV1 expression in both control and heat-treated HEK-293e cells. Interestingly, the mRNA signal was significantly abolished in capsaicin-treated HEK-293e cells in 1 hr. To test these findings, we used prostate cancer cells (PC3), as positive controls expressing TRPV1 [35]. Untreated PC3 cells had lower TRPV1 mRNA levels compared to high levels found following treatment with 32 μM capsaicin for 1 hr, or heat treatment (**Fig. 2A**). This data clearly suggests that TRPV1 is expressed in HEK-293e cells and that presumably, it is over expressed following HS. TRPV1 and Hsp90 protein levels 1 hr after treatment with capsaicin depicted an increased expression compared to control (0 hr) and at 4 and 6 hrs (**Fig. 2C**). Substantiating these findings, a second line of HEK cells devoid of TRPV1 demonstrated that transfection with the TRPV1 gene was associated with an increased expression of Hsp70 in response to treatment with capsaicin (**Fig. 6A**). Furthermore, *Drosophila* cells, known to be devoid of TRPV1, showed no expression of TRPV1 and no response of Hsp70 to treatment with capsaicin (**Fig. 6A**).

To address the specificity of our finding showing a feedback mechanism causing increased TRPV1 expression following treatment with either capsaicin or heat shock, we performed RT-PCR for TRPV4 depicting a constitutive expression of TRPV4 in HEK293e and PC3 cells following treatment with either DMSO, capsaicin, heat shock or heat shock with capsaicin. There was no effect of these treatments on the level of TRPV4 expression. Furthermore, HEK cells which do not express TRPV1 also do not express TRPV4 (**Fig. 6B**).

These results suggest that the cellular stress response and HSP expression in response to heat treatment is controlled, at least in part, by the membrane-associated TRPV1 receptor, and that capsaicin and RTX can induce an isothermal HSR response in different cell types. In support of this finding, we show a decrease in TRPV1 expression in an *in-vivo* model of sepsis-induced lung injury (**Fig. 7C**). Moreover, we found that treatment with an adenoviral vector overexpressing Hsp70 maintained high levels of TRPV1 expression in septic lung cells (**Fig. 7C**), suggesting that the protective effects of the HSR is primarily due to the overexpression of Hsp70.

Confirming the inhibitory effect by capsazepine, a TRPV1 antagonist, we demonstrate a strong TRPV1-dependence of the HSR using two different sequences of TRPV1 siRNAs. TRPV1 siRNA reduced HSP abundance (**Fig. 8**). Blast analysis showed low homology between TRPV1 siRNA and other TRPVs




Figure 2. TRPV1 mRNA and protein abundances in HEK-293e cells. A. Semi-quantitative RT-PCR for TRPV1. Upper left panel: mRNA levels from PC3 prostate cancer cells: non treated cells, cells treated with 32 μ M capsaicin for 1 hr and cells treated with heat shock at 42°C for 3 hrs. Lower right panel: mRNA levels of GAPDH as well as densitometric. Upper right panel: mRNA levels from HEK-293e cells: non treated cells, cells treated with 32 μ M capsaicin for 1 hr (Caps), cells treated with heat shock at 42°C for 3 hrs, and cells treated with heat shock at 42°C for 3 hrs with the addition of 32 μ M capsaicin. Lower right Panel: mRNA levels of GAPDH as well as densitometry. B. Representative immunoblot analysis of TRPV1 abundance. TRPV1 was detected in HEK-293e cells. Cells in culture were incubated with 0.01% DMSO or treated with 4 μ M, 16 μ M or 32 μ M of capsaicin for 1 hr. C. Representative immunoblot analysis of TRPV1 and Hsp90 abundances in HEK-293e cells. Cells were treated with 32 μ M capsaicin for 0 hr, 1 hr, 4 hrs and 6 hrs.

doi:10.1371/journal.pone.0057149.g002

sequences [33,36,37], implying that the observed siRNA effects were specific to TRPV1. Transfection with 200 nM of TRPV1 siRNA attenuated Hsp70 and Hsp90 abundances in a dose-dependent manner, depending on cell type: HEK-293e or HT-29 cells (Fig. 8), compared to scrambled siRNA.

Evidence of Interaction between HSP70 and TRPV1

Co-immunoprecipitation studies in lungs treated, *in-vivo* with an adenoviral vector overexpressing Hsp70 showed an unexpected interaction between TRPV1 and Hsp70 (Fig. 7D). Further, *in-vitro* immuno-histochemistry in HEK-293e cells using co-localized staining revealed an association between TRPV1 and Hsp70, preferentially close to the plasma membrane (Fig. 5A). Hsp70 levels increased in capsaicin and heat shock treated HEK-293e cells compared to non-treated HEK-293e, and decrease in 200 nM TRPV1 siRNA HEK-293e transfected cells (Fig. 5B). In cancerous MCF-7 cells, cytoplasmic expression as well as co-localization of Hsp70 to the membrane can be seen (Fig. 7A).

Discussion

Here we find that TRPV1 acts as a central component of the cellular thermosensory pathway, with direct effects on HSP expression, and that TRPV1 antagonists can block the cellular HSP expression. It has been previously documented that TRPV1

channels are mostly expressed in primary sensory neurons and that they are naturally activated by heat or by capsaicin, protons, and anandamide [38]. We show evidence that TRPV1 acts as a general cellular stress sensor, which in addition of being a thermosensor, elicits a heat-shock-like cellular stress response following heat or other chemical treatments in epithelial cells, such as in the lungs, as well as in various cancerous cells. We demonstrate that treatment of human embryonic kidney cells, alveolar epithelial cells, human breast cancer cells or colon cancer cells with heat, or TRPV1 agonists, capsaicin or RTX, induced the accumulation of Hsp70, Hsp90 and Hsp27, and Hsp70 and Hsp90, respectively, which are hallmarks of the cellular heat-shock response. Thus, in addition to the well-established ability of capsaicin to induce isothermally the activation of an inter-cellular signal resulting in the deceptive sensation of heat [38], TRPV1 agonists can also induce an intracellular signal resulting in the synthesis of HSPs. Further, we show by immunoblots, RT-PCR and Gel-shift assays that capsaicin treatment resulted in the isothermal activation of HSF-1, suggesting that TRPV1, the only known capsaicin target, is directly responsible for the heat or the chemical signal that up-regulates the heat shock genes, and the cellular heat shock response (HSR) in general. In support of these findings, we found that cells devoid of the TRPV1 gene, could not over express Hsp70 in response to heat-shock or capsaicin treatments. Further, we show that in cells harboring TRPV1,




Figure 3. Capsazepine inhibits HSF-1 expression and nuclear translocation. A. Representative immunoblot analysis of nuclear HSF-1 and phospho-HSF-1 abundances. Intra-nuclear HSF-1 and Phospho-HSF-1 were detected in HEK-293e cells. Cells were incubated with 4 μ M, 16 μ M or 32 μ M of capsaicin for 1 hr. HEK-293e cells were treated as indicated with heat shock at 42°C for 3 hrs, heat shock at 42°C for 3 hrs with the addition of 5 mM EGTA, heat shock at 42°C for 3 hrs with the addition of 100 μ M capsazepine or 100 μ M capsazepine with the addition of capsaicin. B. Semi-quantitative RT-PCR for HSF-1. Upper panel: mRNA levels from non treated HEK293e cells (control), cells treated with 32 μ M capsaicin for 1 hr (Caps), cells treated with heat shock at 42°C for 3 hrs, cells treated with heat shock at 42°C for 3 hrs with the addition of 100 μ M capsazepine. Lower Panels: GAPDH mRNA levels from HEK293e cells as indicated above as well as densitometric analysis (Right panel). C. Electrophoretic mobility shift assay for HSF-1 DNA binding activity. Nuclear protein isolated from HEK-293e cells treated with 4 μ M, 16 μ M or 32 μ M of capsaicin for 1 hr. Cells treated as indicated with heat shock at 42°C for 3 hrs, heat shock at 42°C for 3 hrs with the addition of 5 mM EGTA, heat shock at 42°C for 3 hrs with the addition of 100 μ M capsazepine or 100 μ M capsazepine with the addition of capsaicin. 10 μ g of nuclear protein incubated for 20 min at room temperature with a 32 P-labeled double-stranded DNA oligonucleotide containing a consensus –HSF-1 binding site. Right panel indicates densitometric analysis.

doi:10.1371/journal.pone.0057149.g003

either heat-shock or treatment with capsaicin induces an increase of HSF-1 and Phospho-HSF-1 abundances, a finding confirmed by semi quantitative PCR depicting an increase of HSF-1 mRNA, in agreement with previous reports that riluzole benzothiazole can upregulate mRNA levels of HSF-1 and result in HSP expression [39].

TRPV1 mediates heat- and pain sensations by allowing the transient entry mostly of Ca^{2+} , but possibly also Na^{+} and H^{+} , into the cytoplasm of neuronal sensory cells [40]. It has been shown that application of capsaicin to HEK-293 cells stably expressing TRPV1, initiates a marked elevation of Ca^{2+} influx [41]. Here, we show that the specific TRPV1 antagonists (capsazepine and AMG-

9810) and the non-specific calcium chelator EGTA, blocked heat-induced accumulation of HSPs. Inhibition of Ca^{2+} entry in A-431 cells by removing external Ca^{2+} with the addition of EGTA in the medium greatly attenuated HSF-1 translocation from the cytosol to the nucleus, HSF binding to HSE, Hsp70 gene expression, and protein synthesis [42]. The current model of heat-sensing and signaling in animal cells asserts that the activation of HSF-1 by heat results in the presumed unfolding of some unknown heat-labile proteins in the cell, which would recruit Hsp90, Hsp70 and Hsp40 chaperones away from their normal inhibitory association with HSF-1 [18]. Yet, a different mechanism for heat-sensing and activation by heat or chemicals of the HSR has been initially




Figure 4. Effect of various TRPV1 agonists and antagonists on heat shock protein expression. A. Capsaicin & Capsazepine: Representative immunoblots analysis of Hsp70, Hsp90 and Hsp27 abundances. Hsp70, Hsp90 and Hsp27 were detected in MLE-12 (Panel a), HEK-293e (Panel b) cells. Hsp70 was detected in Cells in L3 - primary cells isolated from human lung tumors (Panel d). Cells culture were incubated with 0.01% DMSO, 4 μ M, 16 μ M or 32 μ M of capsaicin (Caps) for 1 hr. 15 μ g of total protein/lane were separated on 9% SDS-PAGE gel. (Panel c) represents Hsp70 and Hsp90 levels in control HEK-293e cells, HEK-293e cells treated with 32 μ M of capsaicin for 1 hr, heat shock treatment for 3 hrs at 42°C and cells treated with both 32 μ M of capsaicin for 1 hr and heat shock for 3 hrs at 42°C. B. RTX & AMG-9810: Representative immunoblot analysis of Hsp70, Hsp90. Hsp70, Hsp90 were detected in MLE-12 and HT-29 cells. Cell cultures were incubated with 0.01% DMSO, 10 nM, 40 nM, 50 nM or 80 nM of RTX for 1 hr, 80 nM RTX with the addition of 5 μ M of AMG-9810 for 1 hr, 5 μ M AMG-9810 at 37°C for 1 hr or 10–50 μ M AMG-9810 at 42°C for 3 hrs. 15 μ g of total protein/lane were separated on 9% SDS-PAGE gel.
doi:10.1371/journal.pone.0057149.g004

suggested [22] and was demonstrated in plants [23], whereby activation occurs via an increase in the fluidity of specific membrane domains, leading to the activation of heat-shock genes [22]. Our results support the existence of such a plasma membrane-dependent thermosensory pathway also in mammalian cells, whereby HSF-1 would become activated primarily through subtle changes in the fluidity state of the plasma membrane, which would in turn activate the transient opening of the specific TRPV1 channels, allowing a specific calcium-dependent signal to propagate and activate HSF-1. Furthermore, we and others have shown that membrane fluidizers, such as benzyl alcohol, cause a HSR under non-inducing temperatures in plants and bacteria [43].

The direct heat-sensing or the perception of a non-thermal chemical stimulus by non-neuronal cells as a HSR stimulus, resulting in the accumulation of HSPs may be associated with changes in the membrane state, especially the organization of the

membrane micro-domains and the activation of a signal protein such as calmodulin [44]. It has been demonstrated that the C-terminus of TRPV1 contains modulatory domains able to bind calmodulin in the presence of bound Ca^{2+} ions [45]. However, the exact second messengers remain elusive.

Our work extends previous evidence that the primary heat-sensing mechanism activating the HSR is located in the plasma membrane of plants, bacteria, yeast and mammalian cells [46,47,48]. Our findings are especially supported by the biochemical, biophysical and pharmacokinetic studies in plants that demonstrated a tight dependence of the heat-induced HSR or isothermal drug treatments, on the transient entry of external Ca^{2+} through specific heat-sensitive Ca^{2+} -channels in the plant plasma membrane [49,50].

Our data point to the plasma membrane-dependent transient potential vanilloid 1 (TRPV) calcium channel-like receptors as




Figure 5. TRPV1 expression in HEK293e cells. A. Left panel: Double immunofluorescence staining of TRPV1 and Hsp70 in HEK-293e cells treated with 32 μM capsaicin for 1 hr. TRPV1 visualized in green, Hsp70 in red and DAPI (Nuclei) in blue. White arrows indicate Hsp70 abundance (left picture) and co-localization (right picture). B. Right panel: Immunofluorescence staining of Hsp70 in HEK-293e cells in non-treated cells, treated cells with 32 μM of capsaicin for 1 hr, heat shock treated cells for 3 hrs at 42°C and cells transfected with 200 nM TRPV1 siRNA and treated with heat shock for 3 hrs at 42°C.

doi:10.1371/journal.pone.0057149.g005

being a major heat shock response sensor in different epithelial non-cancerous and cancerous cells, capable of triggering the cellular HSR in response to thermal or isothermal pharmaceutical treatments. Using two different sequences of TRPV1 siRNA, we demonstrated in animal cells dependence between the TRPV1-receptor and the cellular HSR that recapitulated the inhibitory effect of capsazepine. A scrambled siRNA sequence lacked these effects. Cancer cells devoid of TRPV1 were unable to augment a HSR while introducing the TRPV1 into these cells restored this HSR capacity. Furthermore, S2 *Drosophila* cells, devoid of the TRPV1 gene, treated with capsaicin did not express Hsp70.

TRPV1 is a channel activated by a wide range of agonists (capsaicin, RTX, ethanol, anandamide, NADA, 12-HPETE, camphor, allicin, 2-APB, lidocaine, gingerol, shogaol, piperine monoacylglycerols, w-3 fatty acids, membrane stretch), by low pH and by physical factors such as heat and membrane depolarization [51,52]. Noxious heat directly gates the channel with a threshold for activation at >43°C. However, as others have noted, the concept of 'threshold' is not strictly applicable to a channel such as TRPV1 wherein increases in temperature activate the channel by increasing its probability of activation and wherein the channel, therefore, has some degree of activity at any temperature [53]. Hence, the term 'threshold' should be used as an operational definition to indicate the temperature at which the inward current through a thermo-TRP channel becomes large enough to trigger action potentials in afferent nerve fibers. However, here we look at the effect of the TRPV1 channel on the cell itself and conclude that much lower levels of TRPV1 activation may be necessary for the activation of this intra-cellular response.

In all experiments, HSP expression was found to be repeatedly decreased by the exposure to CZP and AMG-9810 at 42°C. However, although capsaicin and RTX are well-documented specific agonists of TRPV1 [54,28], it cannot be ruled out that some of the depicted effects are also modulated through other non-specific membrane proteins.

Moreover, our data clearly indicate that TRPV1 is not only a sensor of pain and temperature variations that initiate the propagation of a signal among neuronal cells, but it may also act on neuronal as well as non-neuronal cells as a molecular sensor of temperature variation and of other non-thermal cellular stresses, such as inflammatory stimuli (sepsis), activating HSP expression. We demonstrate a significant loss of the TRPV1 receptor in the presence of a septic insult to the lungs and that this loss can be corrected by treatment with an adenovirus expressing Hsp70; this is in agreement with earlier findings indicating that TRPV1 KO mice are highly susceptible to septic insults, compared to the wild type [54]. Further, confocal microscopy in cells treated with capsaicin showed that TRPV1 co-localizes with Hsp70, suggesting that the induction of the HSR promotes the co-localization of Hsp70 to the membrane; this is in accordance with the previously observed case of human tumors, wherein Hsp70 was found to be associated with cholesterol-rich microdomains in the plasma membrane of mouse tumors [55]. Further, our co-immunoprecipitation data suggest a possible interaction between TRPV1 and Hsp70 within lungs that overexpress Hsp70. These changes may be associated with modulation by Hsp70 of TRPV1 trafficking, through PI3K and PKC as shown in sensory nerve endings [55,56]. Others have shown that Hsp70, together with Hsp90, regulate the function, trafficking and turnover of a wide variety of




Figure 6. HEK293e cells versus HEK cells and S2 KO cells – TRPV1 & TRPV4 expressions. A. Representative immunoblot analysis of Hsp70 and TRPV1 in HEK with or without the transfection of the TRPV1 gene (HEK, HEK+V1) and S2 drosophila cells lacking the TRPV1 gene (S2). B. Semi-quantitative RT-PCR for TRPV4. Upper panel: mRNA levels from HEK and PC3 prostate cancer cells: non treated cells (DM), cells treated with 32 μ M capsaicin for 1 hr (Cps) and cells treated with heat shock at 42°C for 3 hrs (HS) or with heat shock at 42°C for 3 hrs with the addition of capsaicin (Cps+HS). Lower panels: mRNA levels from HEK-293e cells: non treated cells (DM), cells treated with 32 μ M capsaicin for 1 hr (Cps), cells treated with heat shock at 42°C for 3 hrs (HS), and cells treated with heat shock at 42°C for 3 hrs with the addition of 32 μ M (Cps+HS) and mRNA levels of GAPDH. doi:10.1371/journal.pone.0057149.g006

signaling proteins [57]. Thus, we speculate that overexpression of Hsp70 or induction of the HSR may initiate TRPV1 stabilization or its trafficking to the membrane.


Several tumor cell lines which express high levels of HSPs have been associated with an increased resistance to chemotherapy [58]. Further, the down-regulation of HSP expression or activity resulted in suppressed tumor growth [14,16]. Here, we demonstrated that capsaicin and capsazepine have similar contradictory effects on cancer cells and on epithelial non-cancerous cells. Others have already described the pro-apoptotic effect of capsazepine [59]. Given that TRPV receptors sense heat and respond to chemicals and to cellular stresses in general, their therapeutic inhibition with specific inhibitors, such as capsazepine, could potentiate various chemotherapies otherwise hindered by an over effective cellular HSR. Furthermore, thermotherapy of particular cancers has been shown to be efficient at first, but a loss of efficacy and resistance was observed, in correlation with increased HSP levels [60]. It is tempting to speculate that by blocking the cellular stress response, capsazepine could rehabilitate cancer chemotherapy [61]. In addition to the possible therapy of resistant cancers associated with HSP over-expression, the pharmacological control of stress-induced HSPs, molecular chaperones in particular, holds promise in the modulation of inflammation, protein misfolding diseases and aging in general.

Materials and Methods


Isolation and preparation of cytosolic, nuclear and whole extracts from cells

HEK293e (Human Embryonic Kidney cells found to express TRPV1), HEK cells, a gift from Prof. Baruch Minke's lab, (not expressing TRPV1), S2 (Drosophila) cells a gift from Prof. Baruch Minke's lab, (lack of TRPV1 gene), MLE-12 (Murine Lung Epithelial) cells (obtained from ATCC, Manassas, VA, USA), HT-29 (human colon cancer) cells, PC3 (human prostate cancer) cells and MCF-7 (human breast carcinoma cells) were grown in DMEM or RPMI containing 10% FCS, 100 units/ml penicillin and 100 μ g/ml streptomycin (Gibco BRL, Grand Island, NY, USA). Cells were incubated with 0.01% DMSO, 4 μ M, 16 μ M and 32 μ M of capsaicin for 1 hr with or without the addition of 100 μ M capsazepine, or treated with either heat shock of 42°C for 3 hrs, with or without 5 mM EGTA (a non-specific calcium chelator) or 100 μ M capsazepine. Others have demonstrated that capsaicin concentrations in the order of 10^{-4} M [62] affected neuronal cells in culture. Here, we have used higher capsaicin concentrations in non-neuronal cells, based on dose response curves; these demonstrated that the doses in the range of 4–32 μ M did not inadvertently affect these epithelial cell cultures (Data not shown). Additional TRPV1 agonist (RTX) and antagonist (AMG-9810) were added to MLE-12 and HT-29. RTX was added in a dose-dependent manner of 10, 40, 50, 80 nM and AMG-9810 in concentrations of 5–50 μ M.

For cell fractionation preparations, media were collected and centrifuged. Detached floating cells (a consequence of addition of capsazepine or EGTA), were added to the attached cell pellets.

A. Hsp70 co-localizes in MCF-7 cell membrane

Immunostaining to Hsp70
in MCF-7 cells treated with 32 μ M Capsaicin – Confocal microscopy
Hsp70 - red Nuclei - blue

B. Hsp70 expression in MCF-7 cells

Non-treated cells

cells treated with 32 μ M capsaicin



C. TRPV1 expression in lung homogenate *in vivo***D. TRPV1 IP in lungs treated with AdHSP *in vivo***

Figure 7. TRPV1 membrane receptor activates the HSR. A. Hsp70 co-localized in MCF-7 cell membrane. Immunofluorescence staining of Hsp70 in MCF-7 cells treated with 32 μ M capsaicin for 1 hr. Hsp70 visualized in red and DAPI (Nuclei) in blue. White arrows indicate co-localization of Hsp70 to the membrane (right picture). B. Hsp70 expression in MCF-7 cells. Immunofluorescence staining of Hsp70 in MCF-7 cells in non treated cells compared to treated cells with 32 μ M capsaicin for 1 hr. C. TRPV1 and Hsp70 were detected in lungs of untreated rats (T0), septic rats treated with PBS (2CLPPBS) or septic rats treated with adenovirus vector over-expressing Hsp70 (2CLPAdHSP). The numbers (1, 2) represent repeated experiments. D. Co-IPs studies in lung homogenates treated with adenoviral vector over-expressing Hsp70 *in-vivo*. 100 μ g of lung homogenates were Co-IPed with TRPV1. Representative immunoblot analysis of TRPV1 and Hsp70 abundances.
doi:10.1371/journal.pone.0057149.g007

The cells were washed with PBS x 1 and lysed with lysis buffer containing 10 mM Tris, (pH 7.9), 60 mM KCl, 1 mM EDTA, 1 mM DTT, 0.4% Nonidet P-40, 0.5 mM PMSF, 1 mM Na₃-Vo₄ and a protease inhibitor cocktail (Roche Diagnostics, GmbH, Mannheim, Germany). The lysates were incubated in ice for 10 min and centrifuged at 2,500 rpm for 5 min at 4°C. The supernatants (cytoplasmic extracts) were collected and pellets were washed three times, suspended in lysis buffer consisting of 20 mM Tris (pH 7.9), 0.4 NaCl, 1.5 mM MgCl₂, 1.5 mM EDTA, 1 mM DTT, 25% glycerol and the protease inhibitor cocktails listed above. After suspension and vortex mixing, the extracts were incubated 10 min on ice and centrifuged 5 min at 10,000 rpm. The supernatant (nuclear extract) was collected for nuclear protein detection. The commercial kit (cell lysis buffer by cell signaling, Technology, Inc) was used for whole cell extractions. Protein concentration was determined using the Bradford method.

Western Blot Analysis

Lysates containing 15 μ g of cytosolic, nuclear or whole cell proteins were separated on 9% SDS-PAGE gels. Immunoblot

signals were detected using enhanced chemiluminescence and quantified using TINA software with scanning densitometry. Hsp70 was identified with a primary mouse monoclonal Hsp70 antibody (Stressgen, Biotechnologies Corp., Canada), Hsp90 with a primary rat Hsp90 monoclonal antibody (Stressgen, Biotechnologies Corp., Canada), Hsp27 with a primary rabbit polyclonal Hsp27 antibody (Stressgen, Biotechnologies Corp., Canada), HSF-1 with a primary rabbit polyclonal HSF-1 antibody (Santa Cruz Biotech Inc., CA USA), Phospho-HSF-1 with a primary rabbit monoclonal phospho-HSF-1 antibody (Abcam Inc, Cambridge, UK), TRPV-1 with rabbit polyclonal TRPV-1 antibodies (Alomone Labs Inc, Israel and Abcam Inc., Cambridge, UK), β -actin with primary mouse monoclonal β -actin antibody (Abcam Inc, Cambridge, UK). Secondary antibodies were goat anti-rabbit IgG, or goat anti-rat IgG or goat anti mouse IgG (Jackson, Immunoresearch Lab., Inc., West Grove, PA., USA).

The TRPV1 receptor's role in the HSR

TRPV1 siRNA assay




Figure 8. TRPV1 siRNA assay. Upper panels: A. TRPV1 siRNA – sequence No. 1: Knockdown of TRPV1, using siRNA, in HEK-293 and HT-29 cells resulted in inhibition of Hsp90, and Hsp70 expression. HEK293 & HT-29 cells were treated with either oligofectamine alone, heat shock at 42°C for 3 hrs, heat shock at 42°C for 3 hrs with the transfection of 100 nM TRPV1 siRNA or with the transfection of 200 nM TRPV1 siRNA. Cells were transfected with TRPV1 siRNA for 72 hrs, prior to heat shock. Lanes grouped by curly brackets represent repeated RNAi experiments. B. TRPV1 siRNA – sequence No. 2 & scrambled siRNA: Knockdown of TRPV1, using a second sequence of siRNA and 200 nM of scrambled siRNA in HEK-293 and HT-29 cells, resulted in inhibition of Hsp90, and Hsp70 expression. Lower panel: Representative immunoblot of TRPV1 abundance. TRPV1 was detected in HEK-293 cells treated with heat shock at 42°C for 3 hrs with the transfection of either 100 nM TRPV1 siRNA or 200 nM TRPV1 siRNA. Lanes grouped by curly brackets represent repeated RNAi experiments.
doi:10.1371/journal.pone.0057149.g008

Electrophoretic Mobility Shift Analysis of HSF-1 DNA Binding Activity

A 32 P-labeled double-stranded DNA oligonucleotide containing a consensus HSE binding site (5'-CTAGAAGCTTCTA-GAAGCTTCAG-3') was used. The labeled oligonucleotide was purified on a G-25 Sephadex column. Nuclear extracts containing 10 μ g of protein were incubated with binding buffer (20 mM Hepes, (pH 7.9), 60 mM KCl, 2 mM EDTA, 5 mM MgCl₂, 10% glycerol, 1 mM PMSF, 1 mM DTT, 0.1% NP-40), dIdC (1 μ g/ μ l) for 20 min at room temperature. The labeled oligonucleotide was added to the reaction mixture for 20 min. Specificity for the binding site was determined by cold competition using a ten-fold excess of unlabeled oligonucleotide. Complexes were visualized by autoradiography.

RT-PCR

Total RNAs from the cell tissues were extracted using Trizol (Sigma, Saint Louis, MO) according to the manufacturer's instructions. 2.5 μ g of extracted RNA (for each sample) was reverse transcribed using 0.5 μ g/ μ l oligo dT, 200 u/ μ l M-MLV

RT and 2.5 mM dNTPs. 2 μ l of the reacting cDNA products were used as template for PCR. PCR was performed using PCR primers specific for human HSF-1: 5' -GTGCAGTCAAACCG-GATCCT -3' (sense) and 5' -GAGATGATGGGTCCA-GAGCTG- 3' (anti sense), and carried out for 40 cycles.

Human TRPV1: 5' - TGTGCCGTTTCATTTT - 3' (sense) and 5' -TGCAGCTTCCAGATGTT - 3' (anti sense), Human TRPV4: 5' -CGTGAGAACACCAAGTTTGT - 3' (sense) and 5 - CTTGCTGTTGTACACCAGGAT - 3 (anti sense), GAPDH was used as a house keeping control gene, using PCR primers specific for GAPDH: 5'- ACCACAGTCCATGCCATCAC-3' and 5'- TCCACCACCCTGTTGCTGTA-3'.

Double immunofluorescence – Confocal microscopy

Double immunofluorescence was performed as follows: cells were fixed in 4% PFA for twenty minutes. Cells were permeabilized using PBS \times 1 buffer containing 5% donkey serum and 0.5% Triton for ten minutes. TRPV1 rabbit polyclonal antibody 1:200 (Alomone Labs Inc., Israel) and Hsp70 mouse monoclonal 1:50 (Stressgen, Biotechnologies Corp., Canada) were added for an hour. The secondary antibodies, goat anti rabbit Alexa 488 and

goat anti mouse RhoDamin X (Jackson, Immunoresearch Lab., Inc., West Grove, PA., USA) were added for thirty minutes. The cells were washed three times and the nuclei were counterstained with DAPI (Vector Laboratories Inc., Burlingame, CA, USA). Confocal imaging was visualized using Nikon Confocal microscope in magnification x 60.

Induction of Sepsis

All studies were approved by the Institutional Animal Care and Use Committees (IACUC) of both collaborating institutions and conform to both University Laboratory Animal Resources (ULAR), The Hebrew University School of Medicine Ethics Committee – research number MD 109.14-4 and the National Institutes of Health standards. Severe sepsis was induced by cecal ligation and double puncture (2CLP) in male adolescent rats under isoflurane anesthesia as previously described [7–9,63,64]. Animals were fluid resuscitated (50 cc/kg of 0.9% saline injected subcutaneously) at the time of surgery and at 24 hours. 48 hours after 2CLP, animals were euthanized with an overdose (150 mg/kg) of pentobarbital. Lungs were homogenized for extraction as previously described [7–9,63,64].

Adenoviral Vector Administration

As previously described, 10^{11} viral plaque-forming units (PFUs) of AdHSP recombinant E1,E3-deleted adenoviral vector dissolved in PBS (total volume, 300 μ l) were administered via tracheal puncture at the time of 2CLP [7–9,63,64]. Control animals were subjected to 2CLP and treated with intratracheal PBS alone.

Immunoprecipitation

Immunoprecipitation was performed as previously described [7–9]. Briefly, 100 μ g of extracts were immunoprecipitated using a rabbit polyclonal anti-TRPV1, (Abcam Inc, Cambridge, UK), that was diluted 1:100. Samples were agitated overnight at 4°C. Protein A/G beads (Pierce Inc., USA) were added and the samples were agitated for two more hours at 4°C and centrifuged at 14,000 rpm for 5 min at 4°C. The resulting pellet was washed three times, suspended in sample buffer and boiled for 5 min. The samples were centrifuged at 14,000 rpm for 5 min at room temperature and the resulting supernatants were subjected to immunoblot as described above.

References

- Westerheide SD, Morimoto RI (2005) Heat shock response modulators as therapeutic tools for diseases of protein conformation. *J Biol Chem* 280(39):33097–33100.
- Kampinga HH (2006) Chaperones in preventing protein denaturation in living cells and protecting against cellular stress. *Handb Exp Pharmacol* 72, 1–42.
- Pratt WB, Toft DO (2003) Regulation of signaling protein function and trafficking by the hsp90/hsp70-based chaperone machinery. *Exp. Biol. Med* 228, 111–133.
- Prahlad V, Cornelius T, Morimoto RI (2008) Regulation of the Cellular Heat Shock Response in *Caenorhabditis elegans* by Thermosensory. *Neurons Science* 320, 811–814.
- Hinault MP, Farina-Henriquez-Cuendet A, Goloubinoff P (2011) Molecular Chaperones and Associated Cellular Clearance Mechanisms against Toxic Protein Conformers in Parkinson's Disease. *Neurodegener Dis* 8(6):397–412.
- Bromberg Z, Deutschman CS, Weiss YG (2005) Heat shock protein 70 and the acute respiratory distress syndrome. *J Anesth* 19(3) 236–242. Review.
- Weiss YG, Bromberg Z, Raj N, Raphael J, Goloubinoff P, et al. (2007) Enhanced heat shock protein 70 expression alters proteasomal degradation of I κ B kinase in experimental acute respiratory distress syndrome. *Crit Care Med* 35(9) 2128–2138.
- Bromberg Z, Raj N, Goloubinoff P, Deutschman CS, Weiss YG (2008) Enhanced Expression of Hsp70 Limits Cell Division in a Sepsis- Induced Model of ARDS. *Crit Care Med* 36, 246–255.
- Aschkenasy G, Bromberg Z, Raj N, Deutschman CS, Weiss YG (2011) Enhanced Hsp70 expression protects the lung during acute lung injury modulating apoptotic cell signaling pathways. *PLoS One* Vol. 6 (11) e26956.

TRPV1 siRNA experiments

RNA interference (siRNA) of human TRPV1 was performed using sequence siRNA reagents (IDT Inc, Coralville, IA, USA) using human TRPV1 siRNA sequence: 5'-GGA TTGCCCTCACGAGGAA-3' [33]. The second TRPV1 siRNA sequence was: 5' – AACUGGAGACUAUUUCCGA- 3' (sense), 5' – UCGGAAAUAGUCUCCAGUU - 3' (anti sense). Furthermore, a scrambled siRNA was used as a transfection negative control. HEK293 and HT-29 cells were transfected with TRPV1 siRNA for 72 hrs using Oligofectamine (Invitrogen Inc., California, USA) as recommended by the manufacturer.

Statistical analysis

Each experiment was repeated at least four times. Where appropriate, a t-test was performed. The significance level was set at $P < 0.05$. Bars in the graphs represent the standard error of the mean.

Acknowledgments

We wish to thank Prof. Eithan Galun, The Goldyne Savad Institute of Gene Therapy, The Hebrew University, Hadassah Medical Center, Jerusalem, for the ongoing support of this research, Hila Giladi, Ph.D., The Goldyne Savad Institute of Gene Therapy, The Hebrew University, Hadassah Medical Center, Jerusalem for her guidance in performing the siRNA studies, Temima Schnitzer Perlman, The Goldyne Savad Institute of Gene Therapy, The Hebrew University, Hadassah Medical Center, Jerusalem for TRPV1 sequences alignments, Sharon Even-Ram, Ph.D., The Goldyne Savad Institute of Gene Therapy, The Hebrew University, Hadassah Medical Center, Jerusalem, for Confocal microscopy and Jonathan Axelrod, Ph.D., The Goldyne Savad Institute of Gene Therapy, The Hebrew University, Hadassah Medical Center, Jerusalem, for his suggestions and comments. We also wish to thank Prof. Baruch Minke, Department of Physiology, the Hebrew University- Hadassah School of Medicine, Jerusalem, Israel, and his Ph.D. student, Maximilian Peters, for giving us the HEK and the S2 Drosophila cells, and for their advice. Finally, we wish to thank Mery Clausen, The Goldyne Savad Institute of Gene Therapy, The Hebrew University, Hadassah Medical Center, Jerusalem, for her editing.

Author Contributions

Conceived and designed the experiments: ZB PG YGW. Performed the experiments: ZB. Analyzed the data: ZB. Contributed reagents/materials/analysis tools: ZB YGW. Wrote the paper: ZB PG YS YGW.

- De Los Rios P, Ben-Zvi A, Slutsky O, Azem A, Goloubinoff P (2006) Hsp70 chaperones accelerate protein translocation and the unfolding of stable protein aggregates by entropic pulling. *Proc Natl Acad Sci U S A* Apr 18;103(16):6166–71.
- Weiss YG, Bouwman A, Gehan B, Scheers G, Raj N, et al. (2000) Cecal ligation and double puncture impairs heat shock protein 70 (HSP-70) expression in the lungs of rats. *Shock* 13(1) 19–23.
- Tiligada E (2006) Chemotherapy: induction of stress responses. *Endocrin Related Cancer* 13 S115–S124.
- Hohfeld J (1998) Regulation of the heat shock conjugate Hsc70 in the mammalian cell: the characterization of the anti-apoptotic protein BAG-1 provides novel insights. *Biol Chem* 379(3) 269–274. Review.
- Nylandsted J, Rohde M, Brand K, Bastholm L, Elling F, et al. (2000) Selective depletion of heat shock protein 70 (Hsp70) activates a tumor-specific death program that is independent of caspases and bypasses Bcl-2. *Proc Natl Acad Sci* 97(14) 871–876.
- Schulte TW, Akinaga S, Soga S, Sullivan W, Stensgard B, et al. (1998) Antibiotic radicicol binds to the N-terminal domain of Hsp90 and shares important biologic activities with geldanamycin. *Cell Stress Chaperones* 3(2) 100–108.
- Banerjee, Mustafi S, Chakraborty PK, Raha S (2010) Modulation of Akt and ERK1/2 pathways by resveratrol in chronic myelogenous leukemia (CML) cells results in the downregulation of Hsp70. *PLoS One* 14;5(1) e8719.
- Akerfelt M, Morimoto RI, Sistonen L (2010) Heat shock factors: integrators of cell stress, development and lifespan. *Nat Rev Mol Cell Biol* Aug;11(8):545–55.
- Voellmy R, Boellmann F (2007) Chaperones regulation of the heat shock protein response. *Adv Exp Med Biol* 594, 89–99.

19. Akerfelt M, Trouillet D, Mezger V, Sistonen L (2007) Heat shock factors at crossroad between stress and development. *Ann N Y Acad Sci* 13, 15–27, review.
20. Trinklein ND, Chen WC, Kingston RE, Myers RM (2004) Transcriptional regulation and binding of heat shock factor 1 and heat shock factor 2 to 32 human heat shock genes during thermal stress and differentiation. *Cell Stress Chaperones* 9(1), 21–8.
21. Török Z, Tsvetkova NM, Balogh G, Horváth I, Nagy E, et al. (2003) Heat shock protein coinducers with no effect on protein denaturation specifically modulate the membrane lipid phase. *Proc Natl Acad Sci U S A* 18;100(6):3131–6.
22. Balogh G, Horváth I, Nagy E, Hoyk Z, Benkő S, et al. (2005) The hyperfluidization of mammalian cell membranes acts as a signal to initiate the heat shock protein response. *FEBS J* 272(23):6077–86.
23. Saidi Y, Finka A, Muriset M, Bromberg Z, Weiss YG, et al. (2009) The heat shock response in moss plants is regulated by specific calcium-permeable channels in the plasma membrane. *Plant Cell* 21(9):2829–43.
24. Nagy E, Balogi Z, Gombos I, Akerfelt M, Björkholm A, et al. (2007) Hyperfluidization-coupled membrane microdomain reorganization is linked to activation of the heat shock response in a murine melanoma cell line. *Proc Natl Acad Sci U S A* 8;104(19):7945–50.
25. Balogh G, Péter M, Liebisch G, Horváth I, Török Z, et al. (2010) Lipidomics reveals membrane lipid remodelling and release of potential lipid mediators during early stress responses in a murine melanoma cell line. *Biochim Biophys Acta* 1801(9):1036–47.
26. Gombos I, Crul T, Poggio S, Güngör B, Török Z, et al. (2011) Membrane-lipid therapy in operation: the HSP co-inducer BGP-15 activates stress signal transduction pathways by remodeling plasma membrane rafts. *PLoS One* 6(12):e28818.
27. Akiyama H, Sasaki N, Hanazawa S, Gotoh M, Kobayashi S, et al. (2011) Novel sterol glucosyltransferase in the animal tissue and cultured cells: evidence that glucosylceramide as glucose donor. *Biochim Biophys Acta* 1811(5):314–22.
28. Mosser DD, Kotzbauer PT, Sarge KD, Morimoto RI (1990) In vitro activation of heat shock transcription factor DNA-binding by calcium and biochemical conditions that affect protein conformation. *Proc Natl Acad Sci U S A* May; 87(10):3748–52.
29. Roy A, Mandadi S, Fiamma MN, Rodikova E, Ferguson EV, et al. (2012) Anandamide modulates carotid sinus nerve afferent activity via TRPV1 receptors increasing responses to heat. *J Appl Physiol* 112(1):212–24.
30. Klafke JZ, da Silva MA, Trevisan G, Rossato MF, da Silva CR, et al. (2012) Involvement of the glutamatergic system in the nociception induced intrathecally for a TRPA1 agonist in rats. *Neuroscience* 222:136–46.
31. Marincák R, Tóth BI, Czifra G, Márton I, Rédl P, et al. (2009) Increased expression of TRPV1 in squamous cell carcinoma of the human tongue. *Oral Dis* 15:328–35.
32. Szallasi A (2011) New developments in the medicinal chemistry of vanilloid TRPV1 and related receptors. *Curr Top Med Chem* 11(17):2116–7.
33. Li WH, Lee YM, Kim JY, Kang S, Kim S, et al. (2007) Transient receptor potential vanilloid-1 mediates heat-shock-induced matrix metalloproteinase-1 expression in human epidermal keratinocytes. *J Invest Dermatol* 127:2328–2335.
34. Vos MH, Neelands TR, McDonald HA, Choi W, Kroeger PE, et al. (2006) TRPV1b overexpression negatively regulates TRPV1 responsiveness to capsaicin, heat and low pH in HEK293 cells. *J Neurochem* Nov; 99(4):1088–102.
35. Czifra G, Varga A, Nyeste K, Marincák R, Tóth BI, et al. (2009) Increased expressions of cannabinoid receptor-1 and transient receptor potential vanilloid-1 in human prostate carcinoma. *J Cancer Res Clin Oncol* Apr;135(4):507–14.
36. Wang Q, Carmichael GG (2004) Effects of length and location on the cellular response to double-stranded RNA. *Microbiol Mol Biol Rev* 68(3), 432–52.
37. Walton SP, Wu M, Gredell JA, Chan C (2010) Designing highly active siRNAs for therapeutic applications. *FEBS J* 277(23), 4806–13.
38. Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, et al. (1997) The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* Oct 23;389(6653):816–24.
39. Liu AY, Mathur R, Mei N, Langhammer CG, Babiarz B, et al. (2011) Neuroprotective drug riluzole amplifies the heat shock factor 1 (HSF1)- and glutamate transporter 1 (GLT1)-dependent cytoprotective mechanisms for neuronal survival. *J Biol Chem* 28, 2785–2794
40. Caterina MJ, Rosen TA, Tominaga M, Brake AJ, Julius D (1999) A capsaicin-receptor homologue with a high threshold for noxious heat. *Nature* 398, 436–441.
41. Han P, McDonald HA, Bianchi BR, Kouhen RE, Vos MH, et al. (2007) Capsaicin causes protein synthesis inhibition and microtubule disassembly through TRPV1 activities both on the plasma membrane and intracellular membranes. *Biochem Pharmacol* May 15;73(10):1635–45.
42. Kiang JG (1994) Corticotropin-releasing factor increases $[Ca^{2+}]_i$ via receptor-mediated Ca^{2+} channels in human epidermoid A-431 cells. *Eur J Pharmacol* Apr 15;267(2):135–42.
43. Saidi Y, Peter M, Finka A, Cicekli C, Vigh L (2010) Membrane lipid composition affects plant heat sensing and modulates Ca^{2+} -dependent heat shock response. *Plant Signal Behav* 5(12):1530–3.
44. Balogh G, Horváth I, Nagy E, Hoyk Z, Benkő S, et al. (2005) The hyperfluidization of mammalian cell membranes acts as a signal to initiate the heat shock protein response. *FEBS J* 272:6077–6086.
45. Jung J, Shin JS, Lee SY, Hwang SW, Koo J, et al. (2004) Phosphorylation of vanilloid receptor 1 by Ca^{2+} /calmodulin-dependent kinase II regulates its vanilloid binding. *J Biol Chem* 279, 7048–7054.
46. Carratù L, Franceschelli S, Pardini CL, Kobayashi GS, Horvath I, et al. (1996) Membrane lipid perturbation modifies the set point of the temperature of heat shock response in yeast. *Proc Natl Acad Sci U S A* Apr 30;93(9):3870–5.
47. Harwood JL, Maresca B, Vigh L (1998) Does the membrane's physical state control the expression of heat shock and other genes? *Trends Biochem Sci* Oct;23(10): 369–74.
48. Vigh L, Horváth I, Maresca B, Harwood JL (2007) Can the stress protein response be controlled by 'membrane-lipid therapy'? *Trends Biochem Sci* Aug;32(8):357–63. Epub 2007 Jul 12.
49. Saidi Y, Finka A, Muriset M, Bromberg Z, Weiss YG, et al. (2009) The heat shock response in moss plants is regulated by specific calcium-permeable channels in the plasma membrane. *Plant Cell* 21(9):2829–43.
50. Saidi Y, Finka A, Chakhporanian M, Zryd JP, Schaefer DG, et al. (2005) Controlled expression of recombinant proteins in *Physcomitrella patens* by a conditional heat-shock promoter: a tool for plant research and biotechnology. *Plant Mol Biol* 59, 697–711.
51. Tominaga M, Caterina MJ, Malmberg AB, Rosen TA, Gilbert H, et al. (1998) The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron* Sep;21(3):531–43
52. Voets T, Droogmans G, Wissenbach U, Janssens A, Flockerzi V, et al. (2004) The principle of temperature-dependent gating in cold- and heat-sensitive TRP channels. *Nature* Aug 12;430(7001):748–54.
53. Vay L, Gu C, McNaughton PC (2012) The thermo-TRP ion channel family: properties and therapeutic implications. *Journal of Pharmacology* 165 787–801.
54. Clark N, Keeble J, Fernandes ES, Starr A, Liang L, et al. (2007) The transient receptor potential vanilloid 1 (TRPV1) receptor protects against the onset of sepsis after endotoxin. *FASEB J* 21(13):3747–55.
55. Stangl S, Gehrman M, Riegger J, Kuhs K, Riederer I, et al. (2011) Targeting membrane heat-shock protein 70 (Hsp70) on tumors by cmHsp70.1 antibody. *Proc Natl Acad Sci U S A* Jan 11;108(2):733–8.
56. Holzer P (2008) The pharmacological challenge to tame the transient receptor potential vanilloid-1 (TRPV1) nociceptor. *Br J Pharmacol* Dec;155(8):1145–62. Review
57. Pratt WB, Morishima Y, Osawa Y (2008) The Hsp90 chaperone machinery regulates signaling by modulating ligand binding clefts. *J Biol Chem* 283:22885–22889.
58. Kiang JG, Gist ID, Tsokos GC (1999) Biochemical requirements for the expression of heat shock protein 72 kDa in human breast cancer MCF-7 cells. *Mol Cell Biochem* 199(1-2), 179–188.
59. Sánchez AM, Sánchez MG, Malagarie-Cazenave S, Olea N, Díaz-Laviada I (2006) Induction of apoptosis in prostate tumor PC-3 cells and inhibition of xenograft prostate tumor growth by the vanilloid capsaicin. *Apoptosis* 11(1), 89–99.
60. Cui X, Yu ZY, Wang W, Zheng YQ, Liu W, et al. (2012) Co-Inhibition of HSP70/HSP90 Synergistically Sensitizes Nasopharyngeal Carcinoma Cells Thermotherapy. *Integr Cancer Ther* Mar;11(1):61–7.
61. Calderwood SK, Khaleque MA, Sawyer DB, Ciocca DR (2006) Heat shock proteins in cancer: chaperones of tumorigenesis. *Trends Biochem Sci* 31(3), 164–72. Review.
62. Franco-Cereceda A, Rydh M, Lou YP, Dalsgaard CJ, Lundberg JM (1991) Endothelin as a putative sensory neuropeptide in the guinea-pig: different properties in comparison with calcitonin gene-related peptide. *Regul Pept* Feb 26;32(3):253–65.
63. Weiss YG, Tazelaar J, Gehan B, Bouwman A, Christofidou-Solomidou M, et al. (2001) Adenoviral vector transfection into the pulmonary epithelium after cecal ligation and puncture (CLP) in rats. *Anesthesiology* 95:974–982.
64. Weiss YG, Maloyan A, Tazelaar J, Raj N, Deutschman CS (2002) Adenoviral transfer of hsp-70 into pulmonary epithelium improves histology and outcome in experimental acute respiratory distress syndrome. *J Clin Invest* 110:801–806.