

Original Article

Molecular Identification of Agents of Human Cutaneous Leishmaniasis and Canine Visceral Leishmaniasis in Different Areas of Iran Using Internal Transcribed Spacer 1 PCR-RFLP

Aref Teimouri^{1,2}, *Mehdi Mohebbali^{1,3}, Elham Kazemirad¹, *Homa Hajjaran¹

¹Department of Medical Parasitology and Mycology, Tehran University of Medical Sciences, Tehran, Iran

²Students Scientific Research Center, Tehran University of Medical Sciences, Tehran, Iran

³Center for Research of Endemic Parasites of Iran, Tehran University of Medical Sciences, Tehran, Iran

(Received 6 June 2017; accepted 16 May 2018)

Abstract

Background: Leishmaniasis is a major medical health problem and distributes in nearly half of 31 provinces of Iran. We aimed to identify cutaneous and visceral *Leishmania* spp. isolated from infected humans and domestic dogs in various regions of Iran, 2010–2013.

Methods: DNA was extracted from 108 lesion exudate samples of suspected patients to cutaneous leishmaniasis and nine liver and spleen aspirates of infected dogs cultured in RPMI-1640 and amplified using partial sequence of ITS1 gene. The PCR amplicons were digested using *Hae*III endonuclease enzyme and used in restriction fragment length polymorphism (RFLP) assay. Then, 48 amplicons representing various hosts were sequenced and compared to sequences from GenBank databases using BLAST.

Results: PCR-RFLP analysis showed that 60 and 48 CL patients were infected by *Leishmania tropica* and *L. major*, respectively. From nine canine visceral leishmaniasis (CVL) isolates, eight isolates were identified as *L. infantum* and one as *L. tropica*. The greatest similarity of 95.7% in ITS1 region was seen between *L. infantum* and *L. major*. Furthermore, the lowest similarity with 65.7% was seen between *L. tropica* and *L. major*. Intra-species comparison of ITS1 region in *L. infantum*, *L. major* and *L. tropica* isolates were showed 100%, 98.2% and 72.4 % similarities, respectively.

Conclusion: PCR-RFLP based on ITS1 region is an appropriate method to distinguish three *Leishmania* spp. of *L. major*, *L. tropica*, and *L. infantum*. In intra-species comparison of ITS1 region, genotypic variations showed that *L. tropica* isolates were more heterogeneous than *L. major* and *L. infantum* isolates.

Keywords: *Leishmania*, ITS1 gene, PCR- RFLP, Iran

Introduction

Cutaneous (CL) and visceral leishmaniasis (VL) include some of the world most neglected diseases in tropical and subtropical areas with an estimated incidence of 0.6–1.0 and 0.05–0.09 million new cases of CL and VL occur worldwide each year, respectively (1). In Iran, CL is considered as a medical health problem in nearly half of 31 provinces (2). Both epidemiological forms of CL are reported in Iran, anthroponotic cutaneous leishmaniasis (ACL) and zoonotic cutaneous leishmaniasis (ZCL) caused by *Leishmania tropica* and *L. major*, respectively (3). Nearly 20000 CL

cases are annually reported and the prevalence of leishmaniasis in provinces of Iran suggestively ranges 1.8–37.9% (2). Furthermore, VL is caused by *L. donovani* complex and *L. infantum* is the main agent of VL in endemic areas of Iran with about 100–300 new cases annually. Infections due to *L. infantum* have been reported in humans, domestic dogs and phlebotomine vectors (4, 5). Human, rodents (mainly great Gerbils (*Rhombomys opimus*)) and domestic canines (*Canis familiaris*) are the main reservoir hosts of ACL, ZCL and VL, respectively (6–8).

*Corresponding authors: Dr Mehdi Mohebbali, E-mail: mohebbali@tums.ac.ir, Dr Homa Hajjaran, Email: hajjaranh@tums.ac.ir

Recently, a few cases of viscerotropic leishmaniasis caused by *L. tropica* and *L. major* have been reported in immunocompromised patients in Iran (9, 10). Furthermore, *L. infantum* is involved in sporadic CL in the endemic areas of VL in Northwestern Iran (11). Because of wide clinical diversities of leishmaniasis, various responses of patients to treatment, various reservoir hosts of the parasite and to develop effective control strategies in endemic areas, distinguish between *Leishmania* spp. and find dominant spp. in provinces are critical. Since *Leishmania* spp. are morphologically identical; therefore, species identification using microscopy or culture methods is not easily possible. For species characterization, additional methods must be used. One of these methods, MLEE (multilocus enzyme electrophoresis) analysis, remains the current gold standard but requires mass culture of the parasites which is not simply carried out (12). Relatively, molecular methods are sensitive for the detection of low amounts of the parasite.

Nowadays, molecular based analyses have extensively been used for the identification of *Leishmania* spp. as well as other parasites. Identification and phylogenetic implication with various targets such as kDNA genes and introns are routine (13). Furthermore, the ITS region of rDNA has been used in several studies to resolve taxonomic questions and to determine phylogenetic affinities among closely related *Leishmania* species (14, 15). The ITS1 gene is considered as the most appropriate region and gives the best results in *Leishmania* differentiation in the old world (16). The ITS1 locates between 18S and 5.8S rRNA genes and possesses conservative loci targets for the PCR (17). Additionally, it includes satisfactory polymorphisms to facilitate species identification.

The aim of this study was to achieve a better understanding of the current status of various isolates of *Leishmania* spp. in Iranian provinces using ITS1-PCR-RFLP. Furthermore,

molecular findings of this study were used to assess phylogenetic relationships between the isolates.

Materials and Methods

Sampling

Samples were collected from 148 lesion exudates of suspected patients to CL referred to Leishmaniasis Laboratory of the School of Public Health, Tehran University of Medical Sciences, Tehran, Iran, and District Health Centers of Iran, 2010–2013. Most of the patients were referred from endemic areas in north-east, southeast, center, west, and southwest of Iran (Fig. 1). Serosity materials of the lesions were smeared on a microscope slide, air-dried, fixed with absolute methanol and stained with Giemsa in 10% phosphate buffer (pH 7.4) for 25min and then examined for amastigotes using light microscope with 1000× magnification. Serosity materials from lesions of CL were cultured in RPMI-1640. Twelve dogs suspected to CVL living in endemic areas were tested serologically using direct agglutination test (DAT). Liver and spleen aspirates of the infected dogs positive for DAT were cultured in RPMI-1640. The parasitological positive dogs showed clinical manifestations of VL including dermatological wounds, ocular variations, weight loss, laziness, lymphadenopathy and splenomegaly (18, 19).

Ethical approval

The study was approved by the Ethics Committee of the Tehran University of Medical Sciences (Approval No. 13462). No information of the patients was revealed in the study. Verbal informed consents were received from the patients and the dog owners.

Direct agglutination test (DAT)

Anti *Leishmania* antibodies in dog sera were detected using DAT. Briefly, promastigotes of *L. infantum*, Iranian strain (MCAN/IR/07/Moheb-gh, GenBank Acces-

sion No. FJ555210), were cultured in RPMI-1640 with 10% heat-inactivated fetal bovine serum (FBS), trypsinized, fixed with 1.2% formaldehyde (Merck, Germany) and stained with Coomassie brilliant blue (Sigma, USA). Negative and positive controls were used in each experiment. Antibody titers $\geq 1:320$ were considered as positive in canines (18).

Parasite culture and cryopreservation

All samples including lesion exudates of the suspected patients to CL and liver and spleen aspirates of the infected dogs were cultured in RPMI-1640 media supplemented with 10–15% heat-inactivated FBS, 100U/ml of penicillin and 100 μ g/ml of streptomycin (Gibco, Life technologies, Germany) and incubated at 24–25 °C. Five days after the last sub-culture, parasites were harvested, washed in sterile phosphate buffered saline (PBS, pH 7.2–7.4), stored at -20 °C until use and were preserved in liquid nitrogen for further studies.

DNA extraction, PCR, and RFLP

Genomic DNA was extracted from pellets of the *Leishmania* cultures using High Pure PCR Template Preparation Kit (Roche Diagnostics, Germany) according to the manufacturer's instructions. Samples were stored at -20 °C until use. DNA samples from Iranian reference strains of *L. tropica* (MHOM/IR/02/Mash10/Accession No. EF653267), *L. major* (MRHO/IR/11/GOL-2/ Accession No. JN860745) and *L. infantum* (MCAN/IR/07/Mash-ir1/ Accession No. EU810776) were used as positive controls. The DNA samples were assessed for the *Leishmania*-specific ribosomal internal transcribed spacer 1 region (ITS1) by PCR amplification using primer pairs of LITSR (F: 5'-CTGGATCATTTTCCGATG-3') and L5.8S (R: 5'-TGATACCACTTATCGCACTT-3'). Amplification was carried out using PCR-Ready Premix (Roche, Germany) in a 25 μ l reaction. The amplification conditions included those described previously (20). PCR products (8 μ l) were digested with

the restriction endonuclease enzyme HaeIII (BsuRI) (Fermentas, Germany) for the species identification according to the manufacturer's instructions. Amplicons of nearly 300–350bp and restriction fragments were analyzed using 1.5–3% agarose gels containing safe stain, visualized under UV and compared with those from reference strains of *L. tropica*, *L. major* and *L. infantum*.

Nucleotide sequence and phylogenetic analysis

PCR products from 44 CL and four CVL samples were sequenced using LITSR as forward primers. DNA sequences obtained from the current study were compared to sequences from GenBank database using Basic Local Alignment Search Tool (BLAST). These sequences have been annotated to GenBank and aligned using Clustal X Software and MEGA Software v.6.0 (21). Various statistical methods (maximum likelihood, UPGMA and Neighbor-Joining (NJ) trees) were digitalized for phylogenetic analysis of the aligned sequences. The Tamura 3-parameter option of the neighbor-joining method was shown as the best phylogenetic tree (22). Inter and intra-species similarities of ITS1 region were calculated for *L. major*, *L. infantum* and *L. tropica* isolates using MEGA Software v.6.0. Bootstrap analysis was carried out with 1000 replicates. Accession numbers of 48 *Leishmania* isolates used in this study are recorded in GenBank as follows: JX289844-JX289881, JN860713-JN860714 and JN860718-JN860725.

Results

Microscopic, culture and DAT analysis

Microscopic study demonstrated *Leishmania* amastigotes in 112 out of 148 smears (75.6%) collected from patients suspected to CL. Furthermore, 108 samples (72.9%) were positive for *Leishmania* parasites in culture method. DAT analysis of domestic dog sera




Fig. 3. The neighbor-joining tree constructed from ITS1 regions of 48 isolates using Tamura 3-parameter. Numbers above branches correspond to bootstrap valued based on 1000 replicates. Branches without numbers include values of less than 70%. Genbank accession numbers are shown in parentheses




Fig. 4. Intra-species mean similarities for *Leishmania* spp. isolated from CL and VL cases based on ITS1 region sequence, 2010–2013




Fig. 5. Inter-species mean similarities among *Leishmania* spp. isolated from CL and VL cases based on ITS1 region sequence, 2010–2013

Discussion

Species identification is critical in diagnosis, treatment and epidemiological studies of *Leishmania* spp. Several studies have amplified ITS-1 region of the ribosomal DNA repeat unit (rDNA-ITS1) for the recognition of Iranian *Leishmania* spp. (8, 23-25). In the present study, a number of unknown *Leishmania* spp. isolated from CL and CVL cases in various endemic regions of Iran was identified by PCR-RFLP and sequencing based on ITS1 region of rRNA gene. In the current study, the major foci of 60 *L. tropica* CL isolates belonged to Khorasan Razavi Province, Bam City (Kerman Province), Kashan City (Isfahan Province). Moreover, one isolate belonged to a CL lupoid patient residing in Shahre-Rey in southern Tehran (ACC, JX289854). Forty-eight *L. major* isolates belonged to Golestan, Isfahan, Kermanshah, Khuzestan, Ilam and Semnan Provinces of Iran (Fig. 1). Two isolates recovered from diffuse cutaneous leishmaniasis (DCL) patients in Damghan City of Semnan Province in central (ACC, JN860713) and Dehloran City of Ilam Province in Southwestern Iran (ACC, JN860714) were identified as *L. major*. Of nine CVL isolates, eight isolates were identified as *L. infantum* (six isolates from Ardabil, one isolate from Golestan and one isolate from Tehran provinces) and one isolate as *L. tropica* (from Golestan Province). The phylogenetic analysis revealed 21 haplotypes within the isolates. No correlations were seen between the haplotypes and geographic distribution of the species complexes. In similar studies in Iran, no correlations were demonstrated between the intra-species divergence and geographical distribution based on RAPD-PCR and ITS1 and N-acetylglucosamine-1-phosphate transferase (NAGT) genes with PCR-RFLP methods (3, 26). In the phylogenetic analysis, *L. tropica* isolates showed 13 haplotypes (H1–H13) indicating a significant divergence between *L. tropica* isolates from the

other two species (Fig. 3). Although the ITS region is one of the best candidates for the differentiation of *Leishmania* at species and strain levels, limited studies have used ITS sequence analysis for *L. tropica* isolates to compare (27-29).

Leishmania tropica (a diploid microorganism) is known as a heterogenous species, quite observed in alignment of the PCR product sequences and in low bootstrap frequencies found in the phylogenetic tree (30-32). The existence of at least two alleles for ITS in rDNA of *Leishmania* spp. may be a good explanation for this phenomenon (33). In the current study, sequences from three *L. tropica* isolates were technically unreadable excluded from the phylogenetic analysis. However, unreadable sequences of ITS-rDNA fragments belonged to positive leishmaniasis cases (24). This can prove high heterogeneity or sometimes mixed infections with two or three *Leishmania* spp. (23, 24, 34, 35).

Similar to pilot studies, a high degree of heterogeneity was seen in *L. tropica* in our study (30, 33, 36, 37). In intra-species analysis, a varied heterogeneity was found at various levels in *Leishmania* spp. using ITS-PCR-RFLP analysis. This included a highest to lowest order of *L. tropica* > *L. major* > *L. infantum* with similarities of 72.4%, 98.2% and 100%, respectively. Relatively, *L. donovani*, *L. infantum* and *L. major* were reported as the less and *L. tropica*, *L. turanica* and *L. gerbilli* as the most divergent complexes (38). In 2005, a heterogeneity variation at various levels in Old World *Leishmania* spp. was reported with a highest to lowest order of *L. tropica* > *L. aethiopica* > *L. major* > *L. donovani* (39). These findings are similar to studies reported that *L. tropica* isolates included the highest divergence in ITS1 genes (26). Similar to a study on *L. major* with 98.2% similarity, results of the current study showed a limited genetic variation with seven haplotypes seen in *L.*

major (H1–H7) (25). Four haplotypes of *L. major* were found in a distinct clade in nine rodent isolates in Central Iran (23) and six haplotypes of *L. major* were identified through PCR-RFLP of rodent samples using ITS1 markers (8).

Results of the present study on *L. infantum* isolates showed no diversities while formed independent and equal clusters with the highest bootstrap values (94%) in the phylogenetic analysis. Moreover, heterogeneity of *L. tropica* with various groups was detected in the current phylogenetic analysis while *L. infantum* isolates consisted of only one haplotype. In a similar study in Turkey using an ITS1 based real-time PCR, genotypic variations of *Leishmania* spp. at species and intra-species levels were observed and heterogeneities were found in *L. tropica* isolates while *L. infantum* isolates formed single group (40). In the present study, the lowest and highest inter-species varieties were seen between *L. major* and *L. infantum* and between *L. tropica* and *L. major* with 4.3% and 34.3% variations, respectively. In the tree, *L. major* was more associated to *L. infantum* than *L. tropica* was. These findings were in contrast to those from studies based on NAGT genes, in which, the lowest inter-species similarity of 95.7% were reported between *L. major* and *L. infantum* (3).

Conclusion

Sequencing results of *Leishmania* spp. isolated from CL and CVL cases showed 93–98% similarities with other annotated sequences in GenBank database. Therefore, PCR-RFLP based on ITS1 region can be suggested as an adequate method to distinguish *L. major*, *L. tropica* and *L. infantum* which are the most prevalent *Leishmania* spp. in Middle East. Furthermore, genotypic variations based on ITS1 region within inter and intra-species of *Leishmania* spp. have shown that *L. tropica* includes more heterogeneity than *L. major* and *L. infantum* do.

Acknowledgements

The authors would like to thank all staff within the leishmaniose laboratory for their collaboration in isolation of *Leishmania* spp., School of Public Health, Tehran University of Medical Sciences, Tehran, Iran. Another thank to Tehran University of Medical Science for financial support (Project Nos.: 13462, 17293 and 34213). The authors declare that there is no conflict of interests.

References

1. World Health Organization (2016) “Leishmaniasis Fact Sheet”. Available at: <http://www.who.int/mediacentre/factsheets/fs375/en>
2. Shirzadi MR, Esfahani SB, Mohebbali M, Yaghoobi-Ershadi MR, Gharachorlo F, Razavi MR, Postigo JAR (2015) Epidemiological status of leishmaniasis in the Islamic Republic of Iran, 1983–2012. *East Mediterr Health J.* 21(10): 736–742.
3. Hajjaran H, Mohebbali M, Teimouri A, Oshaghi MA, Mirjalali H, Kazemi-Rad E, Shiee MR, Naddaf SR (2014) Identification and phylogenetic relationship of Iranian strains of various *Leishmania* species isolated from cutaneous and visceral cases of leishmaniasis based on N-acetyl glucosamine-L-phosphate transferase gene. *Infect Genet Evol.* 26(2014): 203–212.
4. Mohebbali M (2013) Visceral leishmaniasis in Iran: Review of the Epidemiological and Clinical Features. *Iran J Parasitol.* 8(3): 348–358.
5. Yaghoobi-Ershadi MR (2012) Phlebotomine sand flies (Diptera: Psychodidae) in Iran and their role on *Leishmania* transmission. *J Arthropod-Borne Dis.* 6(1): 1–17.
6. Edrissian Gh H, Nadim A, Alborzi AV, Ardehali S (1998) Visceral leishmaniasis: the Iranian experiences. *Arch Iran*

- Med. 1: 22–26.
7. Mohebbali M, Edrissian Gh H, Nadim A, Hajjaran H, Akhoundi B, Hooshmand B, Zarei Z, Arshi Sh, Mirsamadi N, Manouchehri Naeini K, Mamishi S, Sanati AA, Moshfe AA, Charehdar S, Fakhar M (2006) Application of direct agglutination test (DAT) for the diagnosis and seroepidemiological studies of visceral leishmaniasis in Iran. *Iran J Parasitol.* 1: 15–25.
 8. Hajjaran H, Mohebbali M, Abaei MR, Oshaghi MA, Zarei Z, Charehdar S, Mirjalali H, Sharifdini M, Teimouri A (2013) Natural infection and phylogenetic classification of *Leishmania* spp. Infecting *Rhombomys opimus*, a primary reservoir host of zoonotic cutaneous leishmaniasis in northeast Iran. *Trans R Soc Trop Med Hyg.* 107: 550–557.
 9. Alborzi AA, Shamsizadeh A (2006) *Leishmania tropica*—isolated patient with visceral leishmaniasis in southern Iran. *Am J Trop Med Hyg.* 74(2): 306–307.
 10. Shafiei R, Mohebbali M, Akhoundi B, Sharifdini Galian M, Kalantar F, Ashkan S, Fata A, Hosseini Farash BR, Ghasemian M (2014) Emergence of coinfection of visceral leishmaniasis in HIV-positive patients in northeast Iran: A preliminary study. *Travel Med Infect Dis.* 12(2): 173–178.
 11. Badirzadeh A, Mohebbali M, Ghasemian M, Amini H, Zarei Z, Akhoundi B, Hajjaran H, Emdadi D, Molaei S, Kusha A, Alizadeh S (2013) Cutaneous and post kala-azar dermal leishmaniasis caused by *Leishmania infantum* in endemic areas of visceral leishmaniasis, northwestern Iran 2002–2011: a case series. *Pathog Glob Health.* 107(4): 194–197.
 12. Reithinger R, Dujardin JC (2007) Molecular diagnosis of leishmaniasis: current status and future applications. *J Clin Microbiol.* 45(1): 21–25.
 13. Alvar J, Vélez ID, Bern C, Herrero M, Desjeux P, Cano J, Jannin J, Den Boer M (2012) WHO Leishmaniasis Control Team: leishmaniasis worldwide and global estimates of its incidence. *PLoS One.* 7: e35671.
 14. El Tai N, Osman O, El Fari M, Presber W, Schonian G (2000) Genetic heterogeneity of ribosomal internal transcribed spacer in clinical samples of *Leishmania donovani* spotted on filter paper as revealed by single-strand conformation polymorphisms and sequencing. *Trans R Soc Trop Med Hyg.* 94(5): 575–579.
 15. Schonian G, Schnur L, El Fari M, Oskam L, Kolesnikov AA, Sokolowska-Kohler W, Presber W (2001) Genetic heterogeneity in the species *Leishmania tropica* revealed by different PCR-based methods. *Trans R Soc Trop Med Hyg.* 95(2): 217–224.
 16. Odiwuor SO, Saad AA, De Doncker S, Maes I, Laurent T, El Safi S, Mbuchi M, Büscher P, Dujardin JC, Van der Auwera G (2011) Universal PCR assays for the differential detection of all old world *Leishmania* species. *Eur J Clin Microbiol Infect Dis.* 30(2): 209–218.
 17. Roelfsema JH, Nozari N, Herremans T, Kortbeek LM, Pinelli E (2011) Evaluation and improvement of two PCR targets in molecular typing of clinical samples of *Leishmania* patients. *Exp Parasitol.* 127(1): 36–41.
 18. Mohebbali M, Hajjaran H, Hamzavi Y, Mobedi I, Arshi S, Zarei Z, Akhoundi B, Naeini KM, Avizeh R, Fakhar M (2005) Epidemiological aspects of canine visceral leishmaniasis in the Islamic Republic of Iran. *Vet Parasitol.* 129(3–4): 243–251.
 19. Solano-Gallego L, Koutinas AF, Miro G, Cardoso L, Pennisi MG, Ferrer L, Bourdeau P, Oliva G, Baneth G (2009) Directions for the diagnosis, clinical staging, treatment and prevention of canine leishmaniasis. *Vet Parasitol.* 165: 1–18.

20. El Tai NO, El Fari M, Mauricio I, Miles MA, Oskam L, El Safi SH, Presber WH, Schönian G (2001) *Leishmania donovani*: Intraspecific Polymorphisms of Sudanese Isolates Revealed by PCR-based Analyses and DNA Sequencing. *Exp Parasitol.* 97(1): 35–44.
21. Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol.* 30: 2725–2729.
22. Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol.* 4: 406–425.
23. Parvizi P, Ready PD (2008) Nested PCRs and sequencing of nuclear ITS-rDNA fragments detect three *Leishmania* species of gerbils in sandflies from Iranian foci of zoonotic cutaneous leishmaniasis. *Trop Med Int Health.* 13(9): 1159–1171.
24. Mirzaie F, Eslami G, Yosefi MH, Pestehchian N (2013) Molecular identification of *Leishmania* isolates obtained from patients suspected as having cutaneous leishmaniasis referred to reference laboratories from Yazd province in central Iran. *Adv Biomed Res.* 2: 92.
25. Dabirzadeh M, Hashemi M, Maroufi Y (2016) Study of Genetic Variation of *Leishmania major* Based on Internal Transcribed Spacer1 (ITS1) in Chabahar, Iran. *Jundishapur J Microbiol.* 9(6): e33498.
26. Hajjarian H, Mohebbali M, Mamishi S, Vasigheh F, Oshaghi MA, Naddaf SR, Teimouri A, Edrissian GH, Zarei Z (2013) Molecular identification and polymorphism determination of cutaneous and visceral leishmaniasis agents isolated from human and animal hosts in Iran. *Biomed Res Int.* 2013: 789326.
27. Khanra S, Bandopadhyay S, Chakraborty P, Datta S, Mondal D, Chatterjee M, Naskar K, Roy S, Manna M (2011) Characterization of the recent clinical isolates of Indian Kala-azar patients by RAPD-PCR method. *J Parasit Dis.* 35(2): 116–122.
28. Mahdy MA, Al-Mekhlafi HM, Al-Mekhlafi AM, Lim YA, Shuaib NOB, Azazy AA, Mahmud R (2010) Molecular characterization of *Leishmania* species isolated from cutaneous leishmaniasis in Yemen. *PLoS One.* 5(9): e12879
29. Ghatee M, Sharifi I, Mirhendi H, Kannejad Z, Hatam G (2013) Investigation of double-band electrophoretic pattern of ITS-rDNA region in Iranian isolates of *Leishmania tropica*. *Iran J Parasitol.* 8(2): 264–272.
30. Le Blancq SM, Peters W (1986) *Leishmania* in the old world: heterogeneity among *L. tropica* zymodemes. *Trans R Soc Trop Med Hyg.* 80(1): 113–119.
31. Yeo M, Mauricio IL, Messenger LA, Lewis MD, Llewellyn MS, Acosta N, Bhattacharyya T, Diosque P, Carrasco HJ, Miles MA (2011) Multilocus sequence typing (MLST) for lineage assignment and high resolution diversity studies in *Trypanosoma cruzi*. *PLoS Negl Trop Dis.* 5(6): e1049.
32. Azmi K, Schnur L, Schonian G, Nasereddin A, Pratlong F, El Baidouri F, Ravel C, Dedet JP, Ereqat S, Abdeen Z (2012) Genetic, serological and biochemical characterization of *Leishmania tropica* from foci in northern Palestine and discovery of zymodeme MON-307. *Parasit Vectors.* 5: 121.
33. Mauricio IL, Stothard JR, Miles MA (2004) *Leishmania donovani* complex: genotyping with ribosomal internal transcribed spacer and mini-exon. *Parasitology.* 128 (03): 263–267.
34. Strelkova MV, Eliseev LN, Ponirovsky EN, Dergacheva TI, Annacharyeva DK, Erokhin PI, Evans DA (2001) Mixed Leishmanial infections in *Rhombomys opimus*: a key to the persistence of *Leishmania major* from one transmis-

- sion season to the next. *Ann Trop Med Parasitol.* 95(8): 811–819.
35. Parvizi P, Moradi G, Akbari G, Farahmand M, Ready PD, Piazak N, Assmar M, Amirkhani A (2008) PCR detection and sequencing of parasite ITS-rDNA gene from reservoir host of zoonotic cutaneous leishmaniasis in central Iran. *Parasitol Res.* 103(6): 1273–1278.
 36. Schönian G, Mauricio I, Cupolillo E (2010) Is it time to revise the nomenclature of *Leishmania*. *Trends Parasitol.* 26(10): 466–469.
 37. Sallouma T, Khalifeh I, Tokajian S (2016) Detection, molecular typing and phylogenetic analysis of *Leishmania* isolated from cases of leishmaniasis among Syrian refugees in Lebanon. *Parasite Epidemiol Control.* 1: 159–168.
 38. Waki K, Dutta S, Ray D, Kolli BK, Akman L, Kawazu SI, Lin CP, Chang KP (2007) Trans membrane molecules for phylogenetic analyses of pathogenic protists: *Leishmania*-specific informative sites in hydrophilic loops of transendoplasmic reticulum N-acetylglucosamine-1-phosphate transferase. *Eukaryot Cell.* 6(2): 198–210.
 39. Gomez-Marin JE (2005) DNA Fingerprinting in Protozoan Infectious Diseases, In: Read MM, editor. *Trends in DNA Fingerprinting Research.* Hauppauge: NOVA Science Publishers Inc. pp. 62–65.
 40. Toz SO, Nasereddin A, Ozbel Y, Ertabaklar H, Culha G, Sevil N, Ziya Alkan M, Jaffe CL (2009) leishmaniasis in Turkey: molecular characterization of *Leishmania* from human and canine clinical samples. *Trop Med Int Health.* 14(11): 1401–1406.