

POSTER PRESENTATION

Open Access

Antifungal activity of flavonoids and modulation of expression of genes of fatty acid synthesis in the dermatophyte *Trichophyton rubrum*

Tamires Aparecida Bitencourt*, Tatiana Takahasi Komoto, Mozart Marins, Ana Lúcia Fachin

From 5th Congress of the Brazilian Biotechnology Society (SBBIOTEC)
Florianópolis, Brazil. 10-14 November 2013

Background

Dermatophytosis are fungal infections caused by keratinophilic fungi known as dermatophytes and classified in three genera: *Trichophyton*, *Epidermophyton* and *Microsporum*. *Trichophyton rubrum* is the most frequent species associated to dermatophytosis worldwide [1]. The infections caused by dermatophytes are not lethal, but are difficult to treat and uncomfortable. In the case of *T. rubrum*, they tend to be chronic, and although the superficial infections are more common, cases of deep infection have been reported in immunocompromised patients [2][3]. The number of antifungal drugs are still limited, and the acquired resistance for some of clinical antifungal have been shown as well as the side effects that have been promoted by them. Reasons for the challenge in development of new antifungal drugs are the similarities shared by fungal and mammalian cells and the lack of knowledge about the biology of these pathogens. Recent evidences have shown that the fatty acid synthase (FAS) is an interesting antifungal target [4] because of marked differences between human and fungal cells. The aim of this study was to evaluate the antifungal activity of four flavonoids described as FAS inhibitors and verify the modulation of genes in the pathway of fatty acid synthesis in *T. rubrum* growth in presence of the most effective one as FAS inhibitor.

Methods

The susceptibility assay was carried out using the micro-dilution test in RPMI medium in 96-well plates with the flavonoids quercetin, ellagic acid, galangin and genistein in a range of 1000-1.9 µg/mL toward the strain ATCC MYA-3108 of *T. rubrum* [5]. The modulation of genes

fatty acid synthesis(FAS 1p, FAS 2p subunits), acetyl-COA carboxylase 2p subunit and fatty acid transporter protein (*FAT1*) was analyzed by quantitative PCR using Sybr Green after 16h of incubation of strain of *T. rubrum* with MIC of quercetin in liquid Sabouraud medium.

Results and conclusion

Quercetin showed the most effective antifungal activity with MIC of 125 µg/mL, ellagic acid presented MIC of 250 µg/mL, galangin and genistein were ineffective toward *T. rubrum* (MIC >1000 µg/mL). The positive controls fluconazole and cerulenin presented MICs of 63 and 125 µg/mL, respectively. The analyse of gene expression of the fatty acid synthetic pathway showed the majority of the genes were downregulated by quercetin, fluconazole and cerulenin. However, cerulenin caused a low upregulation of *FAS2p* gene. Thus, the results suggest the activity of quercetin could be due the modulation of genes of the pathway of fatty acid synthesis, which is a fungal specific target for development of antifungal drugs.

Acknowledgements

This study was supported by grants from Fundação de Amparo a Pesquisa do Estado de São Paulo (2012/02920-7 and 2009/12419-0) and CAPES.

Published: 1 October 2014

References

1. Martinez DA, Oliver BG, Graser Y, Goldberg JM, Li W, Martinez-Rossi NM, Monod M, Shelest E, Barton RC, Birch E, et al: Comparative genome analysis of *Trichophyton rubrum* and related dermatophytes reveals candidate genes involved in infection. *mBio* 2012, 3:1-14.
2. Achterman RR, White TC: Dermatophyte virulence factors: identifying and analyzing genes that may contribute to chronic or acute skin infections. *International journal of microbiology* 2012, 2012:1-8.
3. Nir-Paz R, Elinav H, Pierard GE, Walker D, Maly A, Shapiro M, Barton RC, Polacheck I: Deep infection by *Trichophyton rubrum* in an immunocompromised patient. *Journal of clinical microbiology* 2003, 41:5298-5301.

University of Ribeirão Preto, Ribeirão Preto, São Paulo, 14096900, Brazil

4. Li XC, Joshi AS, ElSohly HN, Khan SI, Jacob MR, Zhang Z, Khan IA, Ferreira D, Walker LA, Brodel SE, et al: Fatty acid synthase inhibitors from plants: isolation, structure elucidation, and SAR studies. *Journal of natural products* 2002, 65:1909-1914.
5. Rex JH, Alexander BD, Arthington-Skaggs B, Brown SD, Chatuveli V, Espinel-Ingroff A, Ghannoum MA, Knapp CC, Motyl MR, Ostrosky-Zeichner L, et al: Reference method for broth dilution antifungal susceptibility testing of conidium forming filamentous fungi. *NCCLS* 2008, 28:1-9.

doi:10.1186/1753-6561-8-S4-P53

Cite this article as: Bitencourt et al.: Antifungal activity of flavonoids and modulation of expression of genes of fatty acid synthesis in the dermatophyte *Trichophyton rubrum*. *BMC Proceedings* 2014 8(Suppl 4):P53.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

