

Research article

Open Access

Unexpected diversity of cnidarian integrins: expression during coral gastrulation

Brent A Knack^{†1}, Akira Iguchi^{†1}, Chuya Shinzato¹, David C Hayward², Eldon E Ball^{*2} and David J Miller^{*1}

Address: ¹ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Queensland, 4811, Australia and ²Centre for the Molecular Genetics of Development and Research School of Biological Sciences, Australian National University, P. O. Box 475 Canberra, ACT, 2601, Australia

Email: Brent A Knack - brent.knack@jcu.edu.au; Akira Iguchi - akira.iguchi@jcu.edu.au; Chuya Shinzato - chuya.shinzato@jcu.edu.au; David C Hayward - david.hayward@anu.edu.au; Eldon E Ball* - eldon.ball@anu.edu.au; David J Miller* - david.miller@jcu.edu.au

* Corresponding authors †Equal contributors

Published: 9 May 2008

Received: 26 October 2007

BMC Evolutionary Biology 2008, **8**:136 doi:10.1186/1471-2148-8-136

Accepted: 9 May 2008

This article is available from: <http://www.biomedcentral.com/1471-2148/8/136>

© 2008 Knack et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Adhesion mediated through the integrin family of cell surface receptors is central to early development throughout the Metazoa, playing key roles in cell-extra cellular matrix adhesion and modulation of cadherin activity during the convergence and extension movements of gastrulation. It has been suggested that *Caenorhabditis elegans*, which has a single β and two α integrins, might reflect the ancestral integrin complement. Investigation of the integrin repertoire of anthozoan cnidarians such as the coral *Acropora millepora* is required to test this hypothesis and may provide insights into the original roles of these molecules.

Results: Two novel integrins were identified in *Acropora*. *Amltg α 1* shows features characteristic of α integrins lacking an I-domain, but phylogenetic analysis gives no clear indication of its likely binding specificity. *Amltg β 2* lacks consensus cysteine residues at positions 8 and 9, but is otherwise a typical β integrin. In situ hybridization revealed that *Amltg α 1*, *Amltg β 1*, and *Amltg β 2* are expressed in the presumptive endoderm during gastrulation. A second anthozoan, the sea anemone *Nematostella vectensis*, has at least four β integrins, two resembling *Amltg β 1* and two like *Amltg β 2*, and at least three α integrins, based on its genomic sequence.

Conclusion: In two respects, the cnidarian data do not fit expectations. First, the cnidarian integrin repertoire is more complex than predicted: at least two β s in *Acropora*, and at least three α s and four β s in *Nematostella*. Second, whereas the bilaterian α s resolve into well-supported groups corresponding to those specific for RGD-containing or laminin-type ligands, the known cnidarian α s are distinct from these. During early development in *Acropora*, the expression patterns of the three known integrins parallel those of amphibian and echinoderm integrins.

Background

Integrins are a large family of cell surface transmembrane receptors known only from metazoans, which function in intracellular signalling as well as cell-cell and cell-extracel-

lular matrix (ECM) adhesion [1]. As the main mediators of cell-ECM interactions they are key players in early development [2] functioning in gastrulation by rapid modulation of their own adhesion between low and high

affinity states and by modulating the activities of adhesion molecules (e.g. cadherins) in cell layers undergoing convergence and extension [3].

Integrins function as $\alpha\beta$ heterodimers with several subunits of each type being present in most animals. Analyses of the whole genome sequence of *Caenorhabditis elegans* [4,5] indicate that it has a single β subunit of the $\beta 1$ type that is capable of associating with two α subunits, which confer specificity for either laminin- or RGD-containing ligands, and it has been suggested that this may reflect the ancestral state. *Drosophila melanogaster* has five α and two β subunits [5] and mammals, eighteen α and eight β integrin subunits [6]. In each case, however, integrin subunits above and beyond likely orthologs of the two α s and one β of *Caenorhabditis* are clearly lineage-specific. "Lower" animals are of particular significance in terms of understanding the ancestral state, but have not been extensively studied. Both α and β integrin subunits have been identified in sponges [7-9], and cnidarians [7,10,11], but the extent of integrin diversity and the range of functions of these molecules in "lower" animals are unknown.

Anthozoan cnidarians such as the coral *Acropora millepora* and the sea anemone *Nematostella vectensis* appear to have retained much of the genetic complexity of the metazoan common ancestor [12,13]; hence these animals are likely to be highly informative with respect to the ancestral integrin complement and may provide insights into the original roles of these molecules. Known cnidarian integrins include a β integrin from *Acropora millepora* [7] and single α and β subunits (IntA and IntB) from the hydrozoan jellyfish *Podocoryne carnea* [11]. Here we report the characterisation of novel β and α integrins from *Acropora*. The known *Acropora* integrins are expressed during gastrulation in patterns like those seen at the corresponding stages of echinoderm and amphibian development. However, we know from morphological observation [14,15] that *Acropora* gastrulation is not a simple epithelial to mesenchymal transition in which the expressing cells lose their adhesivity and invaginate. Instead, in *Acropora* it is clear that changing cell shape also plays a major role [14]. Two further implications of this work are that the cnidarian integrin complement is significantly more complex than was predicted, and that functional diversification of α integrins may have occurred independently in Cnidaria and Bilateria.

Results

Identification of novel integrins

Three unigenes encoding integrin subunits were identified during an ongoing EST analysis of *Acropora millepora* [13,16]. One of these corresponds to the previously known *Acropora* integrin β Cn1 [7]; to simplify compara-

tive analyses, this integrin is henceforth referred to as AmItg β 1. Complete sequences were determined for cDNA clones corresponding to the other two integrin unigenes; comparative analyses indicated that an EST clone corresponding to a second β subunit (AmItg β 2) encodes a full length protein of 771 amino acids, whilst an α integrin EST clone lacked the 5' end of the open reading frame. To complete the 1021 amino acid integrin α coding sequence, overlapping clones were isolated from a cDNA library, enabling the determination of the complete open reading frame for a molecule designated AmItg α 1. These sequences have been submitted to GenBank under the following accession numbers: [EU239371](#) (AmItg α 1) and [EU239372](#) (AmItg β 2).

AmItg β 2 is a possible coral ortholog of a known jellyfish integrin β

Database comparisons identified AmItg β 2 as a possible ortholog of integrin β (IntB; Q9GSF3) from *Podocoryne* [11]. Previously Reber-Muller et al. [11] suggested that *Podocoryne* IntB was orthologous with AmItg β 1; however, the former not only has higher overall sequence identity with AmItg β 2 (44% amino acid identity compared to <40%), but also shows the same atypical pattern of cysteine residues (Fig. 1, 2) Whereas the β integrin extracellular domain characteristically contains 56 cysteine residues arranged in a specific pattern [7], in both *Podocoryne* IntB [11] and AmItg β 2, cysteine residues at positions 8 and 9 in the canonical structure are absent as occurs in vertebrate $\beta 4$ -type integrins. In terms of most other structural features, however, both AmItg β 2 and *Podocoryne* IntB are typical integrin β s - the MIDAS domain, cysteine-rich stalk and transmembrane region are all clearly present. In both cases, the DxSxS motif of the MIDAS cation-binding domain [17] is completely conserved, whereas the DDL motif of the ADMIDAS is changed to EDL in AmItg β 2 (Fig. 1). The cytoplasmic domains contain the conserved membrane proximal sequence KLLxxxxD and two NPxY/F motifs (NPIF and NPTY in AmItg β 2, NPIY and NPMY in IntB). Whereas the degree of similarity between these *Acropora* and *Podocoryne* sequences implies that these might be orthologs, one complicating factor is that the *Nematostella* genome appears to encode two integrins of the AmItg β 2 type (see below). Hence orthology relationships will only be clear when more complete datasets are available for *Acropora* and *Podocoryne*.

AmItg α 1 is a cnidarian integrin resembling the vertebrate $\alpha 4/9$ -type

Comparative analyses indicate that AmItg α 1 shares some characteristics with those integrin α subunits that lack an α -A (I) domain. Database comparisons revealed that AmItg α 1 is most similar to mouse integrin $\alpha 9$ (MmItg $\alpha 9$; Q91YD5; 28% identity and 48% similarity). Whereas the *Acropora* and *Podocoryne* β integrins that are possible




Figure 1
β integrin alignments (amino terminal end of the molecules). Amino acid sequence of Amltgβ2 aligned with representative β integrin sequences. Atypical absence of cysteines (yellow, numbered) from positions 8 and 9 suggests orthology between Amltgβ2 and *Podocoryne* IntB (PcIntB). Structural features including the MIDAS motif (DLXS, underlined), transmembrane region (long wavy line), membrane proximal motif (short wavy line), and two NPXY/F motifs (double underlined) are conserved. The ADMIDAS motif (DDL, underlined) is changed to EDL in Amltgβ2. An arrow indicates the position where a deletion was made in the sponge sequence (Otltgβ1) to facilitate alignment. Abbreviations and database accession numbers for sequences used in the alignment are: *Acropora* Amltgβ2 (Amltgβ2; EU239372); *Podocoryne* IntB (PcIntB; AAG25994); *Acropora* Amltgβ1 (Amltgβ1; AAB66910); Human β1 (HsItgβ1; P05556); *Strongylocentrotus* βG (Urchin SpItgβG; AAB39739); *Strongylocentrotus* βL (SpItgβL; AAC28382); *Ophlitaspongia* βPoI (Sponge Otltgβ1; AAB66911).

orthologs (Amltgβ2 and PcIntB) are 44% identical, the α subunits Amltgα1 and *Podocoryne* IntA have a much lower amino acid identity (27%), and there are several differences between these that are likely to have functional and/or structural significance. Both proteins are typical in terms of the presence of FG-GAP repeats, transmembrane regions, membrane proximal KxGFFKR motifs and extracellular cleavage sites fitting the RxK/RR consensus (Fig. 3,

4). However, whereas Amltgα1 is typical in having three cation binding motifs (DxD/NxD/NxxD; [18]) within FG-GAP repeats V, VI, and VII, the three cation binding sites in *Podocoryne* IntA are in FG-GAP repeats VI, VII and immediately C-terminal of repeat VII. Both proteins are atypical in terms of the positions of cysteine residues relative to the consensus; Amltgα1 is missing Cys residues at positions 9, 10 and 17, whereas IntB is missing Cys resi-




Figure 2
 β integrin alignments (carboxy terminal end of the molecules). See legend for Fig. 1.

dues at positions 13 and 14, but both proteins have a novel Cys pair between consensus positions 10 and 11. A corresponding extra pair of Cys residues is also present at the same position in both the *Nematostella* α integrin predicted from the genome sequence (NvItgα1; see below) and the atypical *Drosophila* integrin PS3.

Phylogenetic analyses of the novel integrin sequences

To better understand relationships between the *Acropora* sequences and the major integrin types of higher animals, maximum likelihood (ML) phylogenetic analyses were undertaken using MolPhy version 2.3 [19]. Integrin phylogenetics is complicated by high levels of primary sequence divergence and homoplasmy, leading to difficulties in unambiguous alignment of sequences. The analyses presented here are therefore based on extensively edited alignments. Sequences were aligned using ClustalW via the EBI website, manually edited using JalView, and used for phylogenetic analyses. In the case of the integrin α alignment, the output from ClustalW consisted of 2383 positions and was manually edited to 1091 posi-

tions (20 sequences). The corresponding figures for the integrin β alignments were 2159 positions prior to editing and 991 after editing (24 sequences).

The ML phylogenetic analyses of integrin α sequences (Fig. 5A) are broadly consistent with previous studies; the resolution of bilaterian sequences into two major clades corresponding to the major ligand classes RGD (PS2) and laminin (PS1) is strongly supported. The fact that these two clades each contain protostome (fly, worm) and deuterostome (human, sea urchin) sequences indicates that the functional divergence of α integrins had already occurred in Urbilateria – the common ancestor of bilaterian (higher) animals. However, the ML phylogenetic analyses give no clear indication of the likely ligand specificity of the known cnidarian α integrins. The cnidarian (*Podocoryne*, *Acropora* and *Nematostella*) α integrins group together with high bootstrap support (albeit on long branches which reflects their divergence), the sister group of this cnidarian integrin α clade being the α4/9 type integrins (Fig. 5A).




Figure 3
α integrin alignments (amino terminal end of the molecules). The major structural features of alpha integrins lacking an alpha-A domain are conserved in AmItgα1 including seven FG-GAP repeats (underlined, roman numerals), three DxD/NxD/NxxxD cation binding sites (double underline), the transmembrane region (long way line) and the cytosolic membrane proximal domain (short way line). The position of a putative fourth cation binding site in the *Podocoryne* sequence is indicated in red. Arrows mark the positions where regions that could not be unambiguously aligned were removed from the *Drosophila* (DmPS2; 219 residues), *Caenorhabditis* (CePat2; 132 residues) and human (HsItgαV; 6 residues) sequences. Abbreviations and database accession numbers for sequences used in the alignment are: *Acropora* AmItgα1 (AmItgα1; EU239371); *Podocoryne* IntA (PcIntA; AAG25993); *Drosophila* αPS2 (DmPS2; P12080); *Mouse* α9 (MmItgα9; NP_598482); *Human* αV (HsItgαV; P06756); *Caenorhabditis* αPat2 (CePat2; P34446).

ML analyses of the integrin β sequences (Fig. 5B) confirmed the possible orthology of *Podocoryne* IntB with the novel *Acropora* β sequence (AmItgβ2) reported here. Preliminary surveys of the genome sequence of *Nematostella* suggest the presence of at least four β integrins, and gene models (the gene as predicted in the genome assembly, including the open reading frame, introns and untranslated regions) of these were sufficiently complete for them to be included in phylogenetic analyses (Fig. 5B) which group the *Nematostella* β integrins with the *Acropora* sub-

units – two with AmItgβ1 and two with AmItgβ2. It remains to be seen whether each member of these pairs of *Nematostella* genes has an *Acropora* ortholog. As in the integrin α phylogeny, the sponge sequences were relatively distant to those from the Cnidaria. Again, the analyses were broadly consistent with previous studies [20-22], resolving the vertebrate sequences into three clades known as β1 (integrin β1/2/7), β3 (integrin β3/5/6/8) and β4 in the Hughes [20] phylogeny. Unlike the α integrins, there is no evidence for divergence of β subunits





Figure 4
α integrin alignments (carboxy terminal end of the molecules). See legend for Fig. 3.

prior to the protostome/deuterostome split. Rather, it appears more likely that β integrins have diverged independently in several bilaterian lineages.

Expression of integrins during coral gastrulation

RT-PCR analysis demonstrates that mRNAs encoding each of the coral integrins (AmItga1, AmItgβ1 and AmItgβ2) are present in eggs (Knack et al., unpublished data; [23]) and early developmental stages (Fig. 6) at relatively uni-

form levels. No specific pattern could be detected via in situ hybridization prior to the prawn chip stage however, implying that the maternal mRNA is uniformly distributed until this time. In the early development of *Acropora*, a clear and specific in situ hybridization pattern for AmItga1 is first seen in late prawn chip stage embryos (early gastrula) (Fig. 7A), the mRNA being localized in a characteristic pattern on the concave side of the flattened cell bilayer corresponding to Fig. 4b in Hayward et al.

**Figure 5**

Maximum likelihood phylogenetic analysis of representative α and β integrin proteins. Numbers at branch points indicate the percentage of 1000 bootstrap replicates supporting the topology shown (using MolPhy version 2.3; see [14]). (A) α integrins. Whereas integrins from Bilateria group in a ligand specific manner, consistent with previous phylogenies, the cnidarian sequences form an independent clade, reflecting their early divergence. These groupings suggest that functional divergence of α integrins had already occurred in the Urbilateria. Sequences aligned, abbreviations, and accession numbers are: *Lytechinus* SU2 (LvSU2; AAC23572); *Strongylocentrotus* α P (Sp α P; AAD55724); *Drosophila* α PS2 (DmPS2; P12080); Human α 5 (Hslt α 5; P08648); Human α 7 (Hslt α 7; P06756); Human α 8 (Hslt α 8; P53708); Human α 3 (Hslt α 3; P26006); *Drosophila* α PS3 (DmPS3; O44386); *Acropora* Amltg α 1 (Amltg α 1; EU239371); Human α 4 (Hslt α 4; P13612); Human α 9 (Hslt α 9; Q13797); *Nematostella* Nvltg α 1 (Nvltg α 1; XP_001641435); *Caenorhabditis* α Pat2 (CePat2; P34446); *Drosophila* α PS1 (DmPS1; Q24247); *Podocoryne* IntA (PcIntA; AAG25993); *Caenorhabditis* α lna1 (Clna1; Q03600); *Geodia* α (Gcltg α ; CAA65943). (B) β integrins. Major clades resolved here are consistent with previous phylogenies. The position of sequences within the cnidarian clade is consistent with orthology between *Podocoryne* IntB (PcIntB) and Amltg β 2, and groups two *Nematostella* β s with each *Acropora* β . Unlike the α integrins, the β integrins appear to have diverged independently in several bilaterian lineages. Sequences aligned, abbreviations, and accession numbers are: Human β 3 (Hslt β 3; P05106); Human β 5 (Hslt β 5; P18084); Human β 6 (Hslt β 6; P18564); Human β 2 (Hslt β 2; P05107); Human β 7 (Hslt β 7; P26010); Human β 1 (Hslt β 1; P05556); *Strongylocentrotus* β G (Urchin Spltg β G; AAB39739); *Strongylocentrotus* β L (Spltg β L; AAC28382); *Strongylocentrotus* β C (Spltg β C; AAB39740); *Drosophila* β PS (Dm β PS P11584); *Caenorhabditis* β Pat3 (CePat3; Q27874); *Acropora* Amltg β 2 (Amltg β 2; EU239372); *Nematostella* β 1 (Nvltg β 1; XP_001641468); *Nematostella* β 2 (Nvltg β 2; XP_001627336); *Podocoryne* IntB (PcIntB; AAG25994); *Acropora* Amltg β 1 (Amltg β 1; AAB66910); *Nematostella* β 3 (Nvltg β 3; XP_001637894); *Nematostella* β 4 (Nvltg β 4; XP_001621822); *Ophlitaspongia* β Po1 (Sponge Oltg β 1; AAB66911); *Suberites* β (Sponge Sdltg β ; CAB38100); *Geodia* β (Sponge Gcltg β ; CAA77071); Human β 4 (Hslt β 4; P16144); *Drosophila* β -nu (Dm β -nu; Q27591); Human β 8 (Hslt β 8; P26012).

[14]. Expression is maintained in these presumptive endodermal cells as the concavity deepens and they are internalized. Staining remains strong in the presumptive endoderm until blastopore closure is complete (Fig. 7C), after which only weak endodermal staining of Amltg α 1 is observed.

The early expression patterns of both Amltg β 1 and Amltg β 2 (Fig. 7) were broadly similar to that of Amltg α 1, but with the following differences. First, it was only possible to visualize the localization of transcripts corresponding to the β integrins at slightly later stages of development. Second, whereas Amltg α 1 and Amltg β 1 transcripts were tightly restricted at the area of the blast-

opore lip in early gastrulae (Fig 7B and 7E), Amltg β 2 was also expressed more generally (Fig 7G–I).

Discussion

Whereas Reber-Muller et al. [11] hypothesized the presence of only single α and β integrin subunits in cnidarians, the integrins identified to date in *Acropora* are likely to be only a subset of those present. Preliminary surveys of the genome of *Nematostella* imply that at least four β integrins (two resembling Amltg β 1 and two more similar to Amltg β 2), and at least three distinct α types are present (data not shown). However, gene models for only one α integrin were sufficiently complete to allow its inclusion in the phylogenetic analyses shown as Fig 5A. The integrin




Figure 6
Reverse transcriptase PCR analysis of integrin expression during *Acropora* development. Time points: 1-Prawn Chip, 2-Gastrula, 3-Pear, 4-Planula, 5-Settlement. All three *Acropora* integrin subunits show constant levels of expression throughout development.

repertoire of this morphologically simple animal is therefore considerably more complex than that of *Caenorhabditis* (one β and two α s), which has often been assumed to reflect the ancestral metazoan state.

Due to structural constraints on integrin proteins during evolution, phylogenetic analyses are likely to be complicated by homoplasy effects. The position of the cnidarian sequences in the α integrin phylogeny (Fig. 5A) is likely due to both the early divergence of the Cnidaria and homoplasy effects. Hence, in this case, phylogenetics is not informative as to ligand binding properties. It is likely that in cnidarians, as in higher animals, distinct α integrin types participate in binding to laminin and RGD-containing ligands, but functional analyses are required to verify this hypothesis.

Despite having a common pattern of cysteine loss, phylogenetic analyses (Fig 5B) indicate that AmItg β 2 is only distantly related to the vertebrate β 4-type. Those cysteine residues (positions 8 and 9) in the consensus absent from the *Acropora* and *Podocoryne* sequences form the c8-c9 loop of the β A domain, which has been implicated in determining integrin-ligand specificity, specificity of α - β interactions, and signalling properties [24-28]. Whilst this loss has apparently occurred independently of that leading to the vertebrate β 4 type, it may result in common consequences for ligand and/or α -subunit specificity.

The presence of maternal integrin mRNAs and their relatively uniform expression through development reported




Figure 7
Comparison of Amltg α 1, Amltg β 1 and Amltg β 2 mRNA distribution patterns during gastrulation in *Acropora*. At the prawn chip stage, the Amltg α 1 and Amltg β 1 mRNAs are clearly restricted to one side of the flattened cell bilayer (A, A', D). During gastrulation, these mRNAs are tightly restricted to the area of the blastopore (asterisks) lip (B, B', C, C', E, F), and throughout development remain endodermal. The distribution of Amltg β 2 mRNA (G, H, I) is broadly similar to that of Amltg α 1, but is less tightly restricted, as indicated by weak general staining. Arrow heads in A, B, and C indicate the plane of the sections shown as A', B' and C'.

here for *Acropora* have precedents in *Podocoryne* [11] as well as in higher animals. Co-localization of mRNAs for Amltg β 1, Amltg β 2 and Amltg α 1 suggests that either or both of the β subunits associate with the α 1 subunit. There are many precedents from bilaterians for RGD-type α s associating with β 1-type β s (α 5 β 1; α V β 1; α 8 β 1; PS2 β PS). Of these, α 5 β 1 and PS2 β PS have been implicated as regulators of gastrulation in vertebrates [29,30] and *Drosophila* [31] respectively. In *Podocoryne*, IntA is assumed to associate with IntB (a possible Amltg β 2 ortholog) since they are co-expressed in a wide variety of locations over a range of life cycle stages [11], suggesting that Amltg β 2 may associate with the Amltg α 1 subunit.

Although integrins clearly play important roles in gastrulation in several animal groups, including vertebrates [29], *Drosophila* [31], and sea urchins [32], the interactions that have been demonstrated are heterogeneous

with respect to both the ligands and types of integrins involved. The expression patterns of α and β integrin genes observed in the coral are reminiscent of expression during both amphibian and sea urchin gastrulation. Modulation of integrin adhesion to the RGD domain of fibronectin plays a central role during *Xenopus* gastrulation [33], and in the sea urchin *Lytechinus* changes in laminin adhesion mediated by the epithelial α integrin α SU2 are likewise important [34]. In both the sea urchin and *Xenopus*, changes in integrin-mediated cell adhesion during gastrulation occur independently of transcription and translation. In *Acropora*, integrin mRNAs are present in eggs [23] and the extent to which those visualized in the presumptive endoderm reflect zygotic transcription is unknown.

The simplest interpretation of the patterns of integrin expression is that they reflect increases in cell adhesion in the presumptive endoderm. Expression on the concave side of the "fat prawn chip" stage embryo, which is essentially a flat bilayer of cells (for a description of gastrulation in *Acropora*, see [14]), suggests that integrin-based adhesion may constrain the presumptive endoderm whilst the cells of the presumptive ectoderm move upward and inward around them. Whilst the cells that will end up inside the embryo may adhere to each other more tightly than they do to the putative ectoderm, as is consistent with many forms of gastrulation, the inferred increased adhesion in the putative endoderm is not consistent with a typical epithelial to mesenchymal transition.

The early expression patterns of the α and β integrins in *Acropora* are very similar to those of two transcription factors, *snailA* [15] and *otxB* [35]. Snail genes have central and conserved roles in gastrulation in *Drosophila* [36] and vertebrates [37] and members of the broader class of related genes are regulators of other epithelial to mesenchymal transitions (EMTs). Whilst the best understood means by which snail genes regulate cell adhesion is by acting as repressors of E-cadherin expression [36,38,39], in human epidermal keratinocytes the snail-related gene Slug (Snail2) is a repressor of α 3, β 1 and β 4 integrin expression, leading to decreased cell-adhesion to fibronectin and laminin 5 [40]. Across the Bilateria, Otx genes are conserved anterior markers [41] and, whilst most Otx genes are expressed in the nervous system, evidence from a diverse range of metazoans suggests an ancient role as regulators of cell adhesion (eg. [42,43]). In *Hydra*, high levels of CnOtx expression correspond to regions where cells are undergoing rearrangements or movement [44]. Both *snailA* and *otxB* are thus candidate regulators of integrin expression in *Acropora*. Given the similarity of Amltg α 1 to diverged RGD-type mammalian integrins, it will be of particular interest to examine the expression of ECM proteins containing fibronectin type

III domains during cnidarian gastrulation in parallel with adhesion studies. Candidates identified in *Nematostella* include predicted proteins similar to vertebrate usherin and titin, and a likely homolog of *Drosophila* sidekick.

Conclusion

Whilst one might expect morphologically simple metazoans to have a correspondingly basic integrin complement, comprising perhaps just two α s and a single β subunit (as in *Caenorhabditis elegans*), the repertoire of these molecules in anthozoan cnidarians is considerably more complex. In the case of cnidarian α integrins, ligand specificity cannot be predicted by phylogenetic analysis, suggesting the possibility that specificity mechanisms arose independently in Cnidaria and Bilateria. During early development in *Acropora*, some of these adhesion/signalling molecules are expressed in patterns which parallel those of their amphibian and echinoderm counterparts, and which are inconsistent with gastrulation being a simple epithelial to mesenchymal transition.

Methods

Sample collection and RNA extraction

Developmentally staged *Acropora millepora* embryos were collected during annual spawning events. Embryos were staged based on Ball et al. (2002). Total RNA was extracted using RNeasy (Qiagen) according to the manufacturer's protocol.

RT-PCR analysis

RNA was treated with DNase (Fermentas) to remove contaminating genomic DNA. Single stranded cDNA was synthesised using the First-strand cDNA Synthesis Kit (Amersham Biosciences, Piscataway, NJ) using 1 μ g of total RNA. One μ l of this product was used as a polymerase chain reaction (PCR) template. For Amltg α 1, primers Amltg α 1RTF (5'-GCCAATGAAACAGCTACG-3') and Amltg α 1RTR (5'-TTGTCTCCAGCCITCAAC-3') were used to amplify a 130 bp product. For Amltg β 2, primers Amltg β 2RTF (5'-TGGGCATTTGTGGTGTGAG-3') and Amltg β 2RTR (5'-GCTTGTCTGATGAGTGATGG-3') were used to amplify a 219 bp product. For Amltg β 1 (Brower et al. 1997), primers IB1RTF (5'-CTTGTGTTGCCACTTATGGCTT-3') and IB1RTR (5'-CTGCTACTTGCAATTAACGCATC-3') were used to amplify a 144 bp product. The PCR protocol was 1 min at 94°C, then 40 cycles of 0.5 min at 94°C (denaturation), 0.5 min at 50°C (annealing), 2 min at 72°C (extension), followed by an additional extension for 2 min at 72°C. As a control, primers ADH-F (5'-AAGAAGACAAACATCAAGCCTCA-3') and ADH-R (5'-CACATCCAAGGTTCAAGACG-3') were used to amplify a portion of coral AdoHcyase (S-adenosyl-L-homocysteine hydrolase) cDNA (unpublished data).

Whole mount in situ hybridization

The basic procedures for fixation and hybridization with coral embryos were carried out as described [45]. Photographs were captured directly with a Spot digital camera. Digitised images were processed with Adobe Photoshop.

Authors' contributions

BAK was responsible for conducting DNA sequencing and RT-PCR analysis, and participated in both the in situ hybridisation experiments and preparing the manuscript. AI collected and prepared coral developmental stages, assisted in DNA sequencing and, with CS, carried out in situ hybridisation experiments and photographed the results. DCH was responsible for the initial identification of clones, assisted in sequence alignment and critically reviewed both the data and the manuscript. DJM conducted the phylogenetic analyses, and he and EEB drafted and reviewed the manuscript.

Acknowledgements

The authors thank the late Danny Brower for advice and critical input. This work was supported by grants from the Australian Research Council, both directly to DJM and EEB and via the Centre of Excellence for Coral Reef Studies and the Special Research Centre for the Molecular Genetics of Development. BAK acknowledges a James Cook University Postgraduate Research Scholarship, and AI and CS were supported by scholarships from the Okinawa International Exchange and Human Resources Development Council.

References

- Hynes RO: **Integrins: bidirectional, allosteric signaling machines.** *Cell* 2002, **110**:673-687.
- De Arcangelis A, Georges-Labouesse E: **Integrin and ECM functions: roles in vertebrate development.** *Trends Genet* 2000, **16**:389-395.
- Marsden M, DeSimone DW: **Integrin-ECM interactions regulate cadherin-dependent cell adhesion and are required for convergent extension in *Xenopus*.** *Curr Biol* 2003, **13**:1182-1191.
- consortium C: **Genome sequence of the nematode *C. elegans*: a platform for investigating biology.** *Science* 1998, **282**:2012-2018.
- Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG, Scherer SE, Li PW, Hoskins RA, Galle RF, George RA, Lewis SE, Richards S, Ashburner M, Henderson SN, Sutton GG, Wortman JR, Yandell MD, Zhang Q, Chen LX, Brandon RC, Rogers YH, Blazej RG, Champe M, Pfeiffer BD, Wan KH, Doyle C, Baxter EG, Helt G, Nelson CR, Gabor GL, Abril JF, Agbayani A, An HJ, Andrews-Pfannkoch C, Baldwin D, Ballew RM, Basu A, Baxendale J, Bayraktaroglu L, Beasley EM, Beeson KY, Benos PV, Berman BP, Bhandari D, Bolshakov S, Borkova D, Botchan MR, Bouck J, Brokstein P, Brottier P, Burtis KC, Busam DA, Butler H, Cadieu E, Center A, Chandra I, Cherry JM, Cawley S, Dahlke C, Davenport LB, Davies P, de Pablos B, Delcher A, Deng Z, Mays AD, Dew I, Dietz SM, Dodson K, Doup LE, Downes M, Dugan-Rocha S, Dunkov BC, Dunn P, Durbin KJ, Evangelista CC, Ferrez C, Ferriera S, Fleischmann W, Fosler C, Gabrielian AE, Garg NS, Gelbart WM, Glasser K, Glodek A, Gong F, Gorrell JH, Gu Z, Guan P, Harris M, Harris NL, Harvey D, Heiman TJ, Hernandez JR, Houck J, Hosten D, Houston KA, Howland TJ, Wei MH, Ibegwam C, Jalali M, Kalush F, Karpen GH, Ke Z, Kennison JA, Ketchum KA, Kimmel BE, Kodira CD, Kraft C, Kravitz S, Kulp D, Lai Z, Lasko P, Lei Y, Levitsky AA, Li J, Li Z, Liang Y, Lin X, Liu X, Mattei B, McIntosh TC, McLeod MP, McPherson D, Merkulov G, Milshina NV, Mobarry C, Morris J, Moshrefi A, Mount SM, Moy M, Murphy B, Murphy L, Muzny DM, Nelson DL, Nelson DR, Nelson KA, Nixon K, Nusskern DR, Pacleb JM, Palazzolo M, Pittman GS, Pan S, Pollard J, Puri V, Reese MG, Reinert K, Remington K, Saunders RD, Scheeler F, Shen H, Shue BC, Sidani
- Kiamos I, Simpson M, Skupski MP, Smith T, Spier E, Spradling AC, Stapleton M, Strong R, Sun E, Svirskas R, Tector C, Turner R, Venter E, Wang AH, Wang X, Wang ZY, Wassarman DA, Weinstock GM, Weissenbach J, Williams SM, Woodage T, Worley KC, Wu D, Yang S, Yao QA, Ye J, Yeh RF, Zaveri JS, Zhan M, Zhang G, Zhao Q, Zheng L, Zheng XH, Zhong FN, Zhong W, Zhou X, Zhu S, Zhu X, Smith HO, Gibbs RA, Myers EW, Rubin GM, Venter JC: **The genome sequence of *Drosophila melanogaster*.** *Science* 2000, **287**:2185-2195.
- Whittaker CA, Hynes RO: **Distribution and evolution of von Willebrand/integrin A domains: widely dispersed domains with roles in cell adhesion and elsewhere.** *Mol Biol Cell* 2002, **13**:3369-3387.
- Brower DL, Brower SM, Hayward DC, Ball EE: **Molecular evolution of integrins: genes encoding integrin beta subunits from a coral and a sponge.** *Proc Natl Acad Sci U S A* 1997, **94**:9182-9187.
- Pancer Z, Kruse M, Muller I, Muller WE: **On the origin of Metazoan adhesion receptors: cloning of integrin alpha subunit from the sponge *Geodia cydonium*.** *Mol Biol Evol* 1997, **14**:391-398.
- Wimmer W, Perovic S, Kruse M, Schroder HC, Krasko A, Batel R, Muller WE: **Origin of the integrin-mediated signal transduction. Functional studies with cell cultures from the sponge *Suberites domuncula*.** *Eur J Biochem* 1999, **260**:156-165.
- Muller WE: **Origin of metazoan adhesion molecules and adhesion receptors as deduced from cDNA analyses in the marine sponge *Geodia cydonium*: a review.** *Cell Tissue Res* 1997, **289**:383-395.
- Reber-Muller S, Studer R, Muller P, Yanze N, Schmid V: **Integrin and talin in the jellyfish *Podocoryne carnea*.** *Cell Biol Int* 2001, **25**:753-769.
- Ball EE, Hayward DC, Saint R, Miller DJ: **A simple plan--cnidarians and the origins of developmental mechanisms.** *Nat Rev Genet* 2004, **5**:567-577.
- Technau U, Rudd S, Maxwell P, Gordon PM, Saina M, Grasso LC, Hayward DC, Sensen CW, Saint R, Holstein TW, Ball EE, Miller DJ: **Maintenance of ancestral complexity and non-metazoan genes in two basal cnidarians.** *Trends Genet* 2005, **21**:633-639.
- Hayward DC, Samuel G, Pontynen PC, Catmull J, Saint R, Miller DJ, Ball EE: **Localized expression of a dpp/BMP2/4 ortholog in a coral embryo.** *Proc Natl Acad Sci U S A* 2002, **99**:8106-8111.
- Hayward DC, Miller DJ, Ball EE: **snail expression during embryonic development of the coral *Acropora*: blurring the diploblast/triploblast divide?** *Dev Genes Evol* 2004, **214**:257-260.
- Kortschak RD, Samuel G, Saint R, Miller DJ: **EST analysis of the cnidarian *Acropora millepora* reveals extensive gene loss and rapid sequence divergence in the model invertebrates.** *Curr Biol* 2003, **13**:2190-2195.
- Tozer EC, Hughes PE, Loftus JC: **Ligand binding and affinity modulation of integrins.** *Biochem Cell Biol* 1996, **74**:785-798.
- Tuckwell DS, Brass A, Humphries MJ: **Homology modelling of integrin EF-hands. Evidence for widespread use of a conserved cation-binding site.** *Biochem J* 1992, **285** (Pt 1):325-331.
- Adachi J, Hasegawa M: **Instability of quartet analyses of molecular sequence data by the maximum likelihood method: the Cetacea/Artiodactyla relationships.** *Mol Phylogenet Evol* 1996, **6**:72-76.
- Hughes AL: **Evolution of the integrin alpha and beta protein families.** *J Mol Evol* 2001, **52**:63-72.
- Ewan R, Huxley-Jones J, Mould AP, Humphries MJ, Robertson DL, Boot-Handford RP: **The integrins of the urochordate *Ciona* intestinalis provide novel insights into the molecular evolution of the vertebrate integrin family.** *BMC Evol Biol* 2005, **5**:31.
- Huhtala M, Heino J, Casciari D, de Luise A, Johnson MS: **Integrin evolution: insights from ascidian and teleost fish genomes.** *Matrix Biol* 2005, **24**:83-95.
- Iguchi A, Marquez LM, Knack B, Shinzato C, van Oppen MJ, Willis BL, Hardie K, Catmull J, Miller DJ: **Apparent involvement of a beta1 Type Integrin in Coral Fertilization.** *Mar Biotechnol (NY)* 2007.
- Bunch TA, Miller SV, Brower DL: **Analysis of the *Drosophila* betaPS subunit indicates that regulation of integrin activity is a primal function of the C8-C9 loop.** *Exp Cell Res* 2004, **294**:118-129.
- Takagi J, Kamata T, Meredith J, Puzon-McLaughlin W, Takada Y: **Changing ligand specificities of alphavbeta1 and alphavbeta3**

- integrins by swapping a short diverse sequence of the beta subunit. *J Biol Chem* 1997, **272**:19794-19800.
26. Lin CS, Chen Y, Huynh T, Kramer R: **Identification of the human alpha6 integrin gene promoter.** *DNA Cell Biol* 1997, **16**:929-937.
 27. Takagi J, DeBottis DP, Erickson HP, Springer TA: **The role of the specificity-determining loop of the integrin beta subunit I-like domain in autonomous expression, association with the alpha subunit, and ligand binding.** *Biochemistry* 2002, **41**:4339-4347.
 28. Miao H, Li S, Hu YL, Yuan S, Zhao Y, Chen BP, Puzon-McLaughlin W, Tarui T, Shyy JY, Takada Y, Usami S, Chien S: **Differential regulation of Rho GTPases by beta1 and beta3 integrins: the role of an extracellular domain of integrin in intracellular signaling.** *J Cell Sci* 2002, **115**:2199-2206.
 29. Davidson LA, Hoffstrom BG, Keller R, DeSimone DW: **Mesoderm extension and mantle closure in *Xenopus laevis* gastrulation: combined roles for integrin alpha(5)beta(1), fibronectin, and tissue geometry.** *Dev Biol* 2002, **242**:109-129.
 30. Whittaker CA, DeSimone DW: **Integrin alpha subunit mRNAs are differentially expressed in early *Xenopus* embryos.** *Development* 1993, **117**:1239-1249.
 31. Roote CE, Zusman S: **Functions for PS integrins in tissue adhesion, migration, and shape changes during early embryonic development in *Drosophila*.** *Dev Biol* 1995, **169**:322-336.
 32. Marsden M, Burke RD: **The betaL integrin subunit is necessary for gastrulation in sea urchin embryos.** *Dev Biol* 1998, **203**:134-148.
 33. Ramos JW, DeSimone DW: ***Xenopus* embryonic cell adhesion to fibronectin: position-specific activation of RGD/synergy site-dependent migratory behavior at gastrulation.** *J Cell Biol* 1996, **134**:227-240.
 34. Hertzler PL, McClay DR: **alphaSU2, an epithelial integrin that binds laminin in the sea urchin embryo.** *Dev Biol* 1999, **207**:1-13.
 35. de Jong DM, Hislop NR, Hayward DC, Reece-Hoyes JS, Pontynen PC, Ball EE, Miller DJ: **Components of both major axial patterning systems of the Bilateria are differentially expressed along the primary axis of a 'radiate' animal, the anthozoan cnidarian *Acropora millepora*.** *Dev Biol* 2006, **298**:632-643.
 36. Ip YT, Gridley T: **Cell movements during gastrulation: snail dependent and independent pathways.** *Curr Opin Genet Dev* 2002, **12**:423-429.
 37. Carver EA, Jiang R, Lan Y, Oram KF, Gridley T: **The mouse snail gene encodes a key regulator of the epithelial-mesenchymal transition.** *Mol Cell Biol* 2001, **21**:8184-8188.
 38. Barrallo-Gimeno A, Nieto MA: **The Snail genes as inducers of cell movement and survival: implications in development and cancer.** *Development* 2005, **132**:3151-3161.
 39. Nieto MA: **The snail superfamily of zinc-finger transcription factors.** *Nat Rev Mol Cell Biol* 2002, **3**:155-166.
 40. Turner FE, Broad S, Khanim FL, Jeanes A, Talma S, Hughes S, Tselepis C, Hotchin NA: **Slug regulates integrin expression and cell proliferation in human epidermal keratinocytes.** *J Biol Chem* 2006, **281**:21321-21331.
 41. Lichtneckert R, Reichert H: **Insights into the urbilaterian brain: conserved genetic patterning mechanisms in insect and vertebrate brain development.** *Heredity* 2005, **94**:465-477.
 42. Bellipanni G, Murakami T, Doerre OG, Andermann P, Weinberg ES: **Expression of Otx homeodomain proteins induces cell aggregation in developing zebrafish embryos.** *Dev Biol* 2000, **223**:339-353.
 43. Perea-Gomez A, Lawson KA, Rhinn M, Zakin L, Brulet P, Mazan S, Ang SL: **Otx2 is required for visceral endoderm movement and for the restriction of posterior signals in the epiblast of the mouse embryo.** *Development* 2001, **128**:753-765.
 44. Smith KM, Gee L, Blitz IL, Bode HR: **CnOtx, a member of the Otx gene family, has a role in cell movement in hydra.** *Dev Biol* 1999, **212**:392-404.
 45. Hayward DC, Catmull J, Reece-Hoyes JS, Berghammer H, Dodd H, Hann SJ, Miller DJ, Ball EE: **Gene structure and larval expression of cnox-2Am from the coral *Acropora millepora*.** *Dev Genes Eval* 2001, **211**:10-19.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

