

# Acupuncture and Drug Combination Therapy for Abnormal Glucose Metabolism: Exploring Synergistic Enhancement and Reduced Toxicity Mechanisms

Xinyi Tian<sup>1</sup>, Wenjun Wang<sup>2</sup>, Lu Zhang<sup>3</sup>, Liuqing Wang<sup>4</sup>, Kaiqi Zhang<sup>5</sup>, Xiaolei Ge<sup>5</sup>, Zhengrong Luo<sup>5</sup>, Yaqian Zhao<sup>1</sup>, Xu Zhai<sup>3</sup>, Chunjing Li<sup>1</sup>

<sup>1</sup>School of Acupuncture-Moxibustion and Tuina, Shandong University of Traditional Chinese Medicine, Jinan, 250014, People's Republic of China; <sup>2</sup>Xiyuan Hospital, China Academy of Chinese Medical Sciences, Beijing, 100700, People's Republic of China; <sup>3</sup>Graduate School, China Academy of Chinese Medical Sciences, Beijing, 100700, People's Republic of China; <sup>4</sup>Institute of Chinese Medical History and Literatures, China Academy of Chinese Medical Sciences, Beijing, 100700, People's Republic of China; <sup>5</sup>Wangjing Hospital, China Academy of Chinese Medical Sciences, Beijing, 100700, People's Republic of China

Correspondence: Xu Zhai; Chunjing Li, Email [zhaixu@mail.cintcm.ac.cn](mailto:zhaixu@mail.cintcm.ac.cn); [cherry\\_lcj@163.com](mailto:cherry_lcj@163.com)

**Abstract:** This review examines the impact of combining acupuncture with drug therapy on abnormal glucose metabolism and investigates their underlying mechanisms. Conditions like diabetes pose significant health risks due to irregular glucose metabolism. Traditional drug treatments often encounter challenges related to side effects and drug resistance. Acupuncture, as a non-pharmacological intervention, is thought to enhance glucose metabolism and mitigate medication side effects. We selected the relevant studies of acupuncture or electroacupuncture combined with drugs in the treatment of abnormal glucose metabolism in the past five years, and the results indicate that the combination of acupuncture or electroacupuncture and drug therapy markedly enhances glucose metabolism and mitigates medication-related side effects such as gastrointestinal discomfort and hypoglycemia. Overall, this review underscores the synergistic benefits of acupuncture and drug therapy in improving treatment efficacy and reducing adverse effects, offering promising new approaches for managing abnormal glucose metabolism. Our review provides evidence for the potential benefits of combining acupuncture with drug therapy for abnormal glucose metabolism, which could lead to improved treatment outcomes and reduced side effects for patients with type 2 diabetes mellitus.

**Keywords:** acupuncture, drug, medicine, combination therapy, abnormal glucose metabolism, synergistic enhancement, reduced toxicity

## Introduction

Abnormal glucose metabolism is becoming more prevalent as a type of metabolic disorder with significant global health implications. These abnormalities not only affect patients' quality of life directly but also elevate the risk of serious conditions like cardiovascular disease and diabetes.<sup>1</sup> Abnormal glucose metabolism encompasses a range of conditions such as diabetes, insulin resistance, and metabolic syndrome.<sup>2</sup> Type 2 diabetes mellitus (T2DM) is particularly noteworthy due to its widespread prevalence and significant impact on global health.<sup>3</sup> According to data from the World Health Organization, the global adult diabetes prevalence has quadrupled since 1980. This increase is most pronounced in low- and middle-income countries, particularly in urbanized and rapidly developing economic regions.<sup>4</sup> Risk factors for T2DM include genetic predisposition, unhealthy dietary habits (such as high-sugar and high-fat diets), physical inactivity, obesity, age, and ethnicity. The global surge in obesity, particularly among children and adolescents, has contributed to the rising incidence of T2DM. This condition not only heightens the risk of severe complications such as cardiovascular diseases, stroke, blindness, and kidney disease but also diminishes patients' quality of life, imposing

substantial burdens on families and societal economies.<sup>5</sup> Hence, effective prevention and management strategies are crucial. These include health education initiatives, promoting healthy diets and lifestyle improvements, encouraging physical activity, and timely medical interventions.<sup>6</sup> Comprehensive health policies and global cooperation can effectively mitigate the adverse impacts of abnormal glucose metabolism and its complications worldwide. This approach promises to enhance global population health and improve overall quality of life.<sup>7</sup>

Chemical drug intervention remains the primary approach for managing glucose metabolism disorders; however, it also has limitations such as drug side effects, tolerance issues, individual variability, and suboptimal long-term efficacy. Adverse effects, including gastrointestinal discomfort and the risk of hypoglycemia, often impact patient adherence and may exacerbate metabolic burdens with prolonged use.<sup>8</sup> Consequently, an increasing number of studies are focusing on alternative methods to improve treatment outcomes, aiming to control blood glucose levels while reducing adverse drug reactions and enhancing patients' quality of life.

Acupuncture, as one of the traditional Chinese medical therapies, is gradually recognized as a complementary and alternative treatment for glucose metabolism disorders. Numerous studies have demonstrated that acupuncture can positively influence T2DM and other glucose metabolism disorders through multiple pathways, such as modulating the neuroendocrine-immune system.<sup>9,10</sup> Specifically, acupuncture may promote glucose metabolism by activating metabolic signaling pathways, such as AMPK, or by regulating the nervous system to affect insulin secretion and utilization efficiency, thereby assisting in blood glucose control. Furthermore, the mechanisms of acupuncture also include the regulation of serotonin levels, bile acid metabolism, and gut microbiota, which can alleviate gastrointestinal discomfort caused by certain medications.<sup>11,12</sup>

The mechanism of the combined effects of acupuncture and pharmacotherapy is a complex and synergistic process that integrates the strengths of both treatments, enhancing therapeutic efficacy and reducing side effects through interaction and complementarity. Many clinical and basic studies have confirmed that the combined use of acupuncture and medication demonstrates superior effectiveness in lowering blood glucose and alleviating medication-related side effects.<sup>13</sup> Specifically, acupuncture can enhance the therapeutic pathways of drugs to strengthen their efficacy while also modulating metabolic pathways or blocking certain routes to mitigate potential side effects. By interacting with medications, acupuncture not only improves the safety of treatments but also significantly enhances patient efficacy and tolerance over the long term.<sup>14</sup>

In summary, the combination of acupuncture and pharmacotherapy for treating glucose metabolism disorders shows promising prospects, potentially becoming an effective and viable integrative approach in managing diabetes and other metabolic diseases. Future research exploring the mechanisms of action and applicability across different patient populations will facilitate the standardization and promotion of this approach in clinical practice.

## Conventional Therapies for Intervening in Abnormal Glucose Metabolism

### Pharmacological Interventions for Abnormal Glucose Metabolism

Chemical drug intervention remains the primary approach for treating abnormal glucose metabolism. Commonly prescribed drug classes include Biguanides, sulfonylureas, meglitinides, thiazolidinediones (TZDs), alpha-glucosidase inhibitors, dipeptidyl peptidase inhibitors (DDP-4 inhibitors), sodium-glucose cotransporter 2 inhibitors (SGLT2 inhibitors), GLP-1 receptor agonists, and insulin. Biguanides like metformin exert robust hypoglycemic effects by reducing hepatic glucose output, enhancing peripheral tissue insulin sensitivity, and improving glucose uptake and utilization. They are typically recommended as first-line therapy for type 2 diabetes due to their favorable safety profile and relatively low cost.<sup>15</sup> Sulfonylureas such as gliclazide and glimepiride stimulate insulin release from pancreatic  $\beta$ -cells, improve insulin sensitivity, reduce insulin resistance, and thereby lower hepatic glucose production. They are prescribed when metformin alone proves insufficient in controlling blood glucose levels and are also considered a first-line therapy for treating type 2 diabetes mellitus.<sup>16</sup> In addition, GLP-1 receptor agonists like liraglutide and exenatide replicate the effects of GLP-1. They enhance insulin release by binding to GLP-1 receptors, suppress glucagon secretion, reduce blood glucose levels, and contribute to weight loss.<sup>17</sup> SGLT-2 inhibitors such as empagliflozin and dapagliflozin are newer additions to antidiabetic medications. They lower blood glucose levels by blocking renal glucose reabsorption, which

facilitates the excretion of excess glucose through urine.<sup>18</sup> Alongside traditional first-line drug therapies,  $\alpha$ -glucosidase inhibitors such as acarbose delay carbohydrate absorption by inhibiting  $\alpha$ -glucosidase in the small intestine's brush border. This mechanism helps to decrease postprandial hyperglycemia and is suitable for individuals resistant to metformin or experiencing significant side effects.<sup>19</sup> Insulin is appropriate for all patients with type 1 diabetes and for those with type 2 diabetes who do not respond adequately to oral hypoglycemic drugs and dietary management. It is typically administered through regular subcutaneous injections.<sup>20</sup> Meglitinides lower blood glucose levels by stimulating the release of insulin from the beta cells in the pancreas, with their effects primarily focused on controlling postprandial glucose levels. Thiazolidinediones (TZDs) primarily reduce blood glucose by increasing the sensitivity of target cells to insulin. Dipeptidyl Peptidase-4 (DPP-4) inhibitors lower blood glucose by inhibiting the activity of the DPP-4 enzyme, thereby prolonging the half-life of incretin hormones such as glucagon-like peptide-1 (GLP-1) and gastric inhibitory polypeptide (GIP).<sup>21</sup> The specific drug classification and side effects are shown in Table 1.

Furthermore, these drugs exert their effects through various mechanisms, including improvements in intermittent inflammation,<sup>22</sup> activation of the AMPK signaling pathway,<sup>23</sup> modulation of intestinal microbiota,<sup>24</sup> regulation of cellular autophagy,<sup>25,26</sup> binding with cytokines to enhance cellular function,<sup>27,28</sup> and other pathways to improve abnormal glucose metabolism.

**Table 1** Classification of T2DM Medications and Their Mechanisms of Action and Side Effects

Drug Class	Representative Drugs	Mechanism of Action	Related Side Effects
Biguanides	Metformin	Reduces blood glucose by decreasing hepatic glucose production, increasing muscle insulin sensitivity, and improving lipid metabolism.	Gastrointestinal discomfort (eg, diarrhea, nausea), lactic acidosis (rare).
Sulfonylureas	Glimepiride, Glyburide	Stimulates insulin release from pancreatic $\beta$ -cells, thereby lowering blood glucose levels.	Hypoglycemia, weight gain, rash.
Meglitinides	Repaglinide, Nateglinide	Stimulates insulin release from pancreatic $\beta$ -cells, mainly used for postprandial glucose control.	Hypoglycemia, weight gain.
Thiazolidinediones (TZDs)	Rosiglitazone, Pioglitazone	Activates PPAR- $\gamma$ to increase insulin sensitivity in target cells.	Edema, weight gain, increased risk of heart disease (to be used with caution).
$\alpha$ -Glucosidase Inhibitors	Acarbose, Miglitol	Inhibits the activity of $\alpha$ -glucosidase in the small intestine, slowing carbohydrate absorption and reducing postprandial blood glucose.	Bloating, diarrhea, gastrointestinal discomfort.
Dipeptidyl Peptidase-4 Inhibitors (DPP-4 Inhibitors)	Sitagliptin, Vildagliptin	Inhibits the activity of DPP-4 enzyme, prolonging the half-lives of GLP-1 and GIP, promoting insulin secretion and inhibiting glucagon.	Gastrointestinal discomfort, hypoglycemia (rare), rash.
Sodium-Glucose Cotransporter-2 Inhibitors (SGLT-2 Inhibitors)	Dapagliflozin, Empagliflozin	Inhibits SGLT-2 in the renal tubules, reducing glucose reabsorption and promoting glucose excretion.	Urinary tract infections, ketoacidosis, dehydration.
GLP-1 Receptor Agonists	Liraglutide, Exenatide	Mimics the action of GLP-1, promoting insulin secretion, inhibiting glucagon, and slowing gastric emptying.	Gastrointestinal discomfort (eg, nausea, vomiting), hypoglycemia (rare).
Insulin	Regular insulin, long-acting insulin (eg, glargine insulin)	Directly replaces the insulin deficiency in the body, promoting cellular uptake and utilization of glucose.	Hypoglycemia, weight gain, allergic reactions.

## Limitations of Pharmacological Therapy

Traditional pharmacological therapies for treating abnormal glucose metabolism have limitations, including drug side effects, tolerance issues, individual variability in response, and questions about long-term efficacy.

### Drug Side Effects

Traditional pharmacological therapies for treating abnormal glucose metabolism face limitations, such as drug side effects, tolerance issues, individual variability in response, and concerns regarding long-term efficacy.<sup>29</sup> GLP-1 receptor agonists like exenatide, liraglutide, and semaglutide have been noted to induce gastrointestinal discomfort as a side effect.<sup>30–32</sup> While metformin is widely acknowledged for its efficacy and safety, approximately 30% of patients using this medication may experience severe adverse effects such as nausea, vomiting, and diarrhea.<sup>33–35</sup>

According to a clinical trial, nearly half of the patients experienced hypoglycemia after taking sulfonylurea drugs such as glimepiride.<sup>36</sup> Case reports also suggest that using non-sulfonylurea insulin secretagogues like repaglinide alone or in combination with clopidogrel increases the risk of hypoglycemia.<sup>37,38</sup> Since the discovery of insulin, hypoglycemia has been a well-documented complication associated with its use.<sup>39</sup>

### Drug Tolerance, Individual Differences, and Long-Term Efficacy

Long-term use of the same drug may lead to drug resistance, where patients gradually become less responsive to the medication. This often requires dose adjustments or switching to alternative treatments to maintain therapeutic effectiveness. Individual differences in drug absorption, distribution, metabolism, and excretion directly impact the long-term efficacy and tolerance of medications. Thus, understanding these factors is crucial in managing abnormal glucose metabolism effectively.<sup>40</sup> Other studies have shown taxonomic and functional differences in the intestinal microbiota between patients who are intolerant to metformin and those who tolerate the medication well.<sup>41</sup> Differences in patients' intestinal microbiota directly or indirectly influence drug tolerance, thereby affecting the long-term efficacy of medications.

## Application and Research of Acupuncture and Medicine Combination in Intervening Abnormal Glucose Metabolism

### Clinical and Basic Research on Acupuncture and Moxibustion Intervention of Abnormal Glucose Metabolism

Acupuncture, valued for its simplicity, affordability, safety, and effectiveness,<sup>42,43</sup> is recognized as a complementary and alternative therapy for abnormal glucose metabolism.<sup>44–47</sup> Acupuncture can positively influence type 2 diabetes mellitus (T2DM) and other metabolic disorders by regulating the neuroendocrine immune system, enhancing insulin sensitivity, and adjusting metabolic functions.

In clinical research, a recent 2024 study indicates that electroacupuncture can reduce plasma glucose levels by enhancing insulin production and improving insulin sensitivity. Additionally, it lowers sympathetic nervous activity by promoting the secretion of endogenous  $\beta$ -endorphin and serotonin.<sup>48</sup> A three-arm randomized controlled trial demonstrated that acupuncture improves glucose metabolism more effectively than metformin and has a lower incidence of gastrointestinal reactions compared to metformin. Acupuncture may therefore serve as a low-risk non-pharmacological treatment option for patients with polycystic ovary syndrome.<sup>49</sup> Additionally, several meta-analyses have demonstrated that acupuncture, as an adjunctive therapy to antidiabetic medications, has a statistically significant effect on lowering fasting blood glucose and improving insulin resistance. Although its effects on HbA1c, 2-hour blood glucose, and fasting insulin are less certain, acupuncture is generally considered safe for patients with mild diabetes. As a multifaceted intervention, acupuncture is recommended as a complementary or alternative therapy for T2DM, particularly for patients with obesity or metabolic disorders. In summary, acupuncture and electroacupuncture, as supplementary therapies, offer effective intervention for abnormal glucose metabolism with a favorable safety profile.<sup>43,44</sup>

In basic research, a study indicated that abdominal acupuncture regulates the FGF21 signaling pathway and related adipokines in obese rats with type 2 diabetes mellitus (T2DM). This intervention resulted in reduced fasting blood glucose levels, decreased body weight, and improved insulin sensitivity.<sup>50</sup> Multiple studies have demonstrated that

acupuncture or electroacupuncture achieves glycemic control by modulating the intestinal microbiota,<sup>51,52</sup> regulating the Akt/FoxO1 signaling pathway,<sup>53</sup> activating the AMPK signaling pathway,<sup>54,55</sup> regulating the macrophage-islet adipocyte-islet  $\beta$ -cell network,<sup>56</sup> and inhibiting the PKC $\beta$ /P66shc signaling pathway and oxidative stress.<sup>57</sup>

## Clinical and Basic Research on Intervention of Abnormal Glucose Metabolism by Combination of Acupuncture and Medicine

The mechanism of combining acupuncture with medicine is a complex and synergistic process that integrates the benefits of both therapies. Through interaction and complementation, this approach aims to enhance therapeutic efficacy and minimize side effects.<sup>58,59</sup> Acupuncture can enhance the efficacy of drugs by facilitating their pathways of action, while also potentially reducing drug-related side effects by blocking specific pathways.

In terms of enhancing efficacy, several basic studies have demonstrated that electroacupuncture at the “Tianshu” acupoint (ST 25) promotes the activation of AMPK in the liver tissue of rats with type 2 diabetes mellitus (T2DM) treated with metformin. This intervention helps mitigate the negative effects of metformin on pancreatic tissue AMPK and enhances its glucose-lowering effects. The mechanism may involve upregulation of AMPK expression and inhibition of the pancreatic intrinsic nervous system,<sup>55,60</sup> and activation of GLUT4 activation.<sup>61</sup> Furthermore, electroacupuncture can enhance the glucose-lowering ability of acarbose in T2DM rats and alleviate insulin resistance, with superior effects compared to acarbose alone.<sup>62</sup> Clinical studies have shown that knife-edge acupuncture combined with metformin hydrochloride tablets can improve blood glucose, clinical symptoms, and quality of life in patients with T2DM, possibly through modulation of inflammatory responses.<sup>63</sup> Furthermore, electroacupuncture has been shown to enhance the glucose-lowering efficacy of acarbose in rats with type 2 diabetes mellitus (T2DM) and alleviate insulin resistance. This combined approach demonstrates superior effects compared to acarbose treatment alone,<sup>64</sup> this effect may be achieved through improvements in weight reduction and reduction of inflammation, thereby alleviating insulin resistance.<sup>65,66</sup> Additionally, it may improve lipid metabolism and regulate adipokines.<sup>67</sup>

In terms of reducing toxicity, a systematic review indicated that acupuncture effectively alleviates nausea and vomiting caused by glucagon-like peptide-1 receptor agonists,<sup>68</sup> possibly through mechanisms involving 5-HT,<sup>69</sup> intestinal microbiota,<sup>70</sup> and other factors.<sup>71</sup> Moreover, acupuncture has been shown to alleviate drug-induced liver damage. A basic study indicated that acupuncture at Zusanli and Taichong points may have a therapeutic effect on carbon tetrachloride-induced liver damage in rats, effectively reducing biochemical and pathological indicators of liver damage,<sup>72</sup> possibly through mechanisms involving apoptosis and ferroptosis.<sup>73</sup> Additionally, acupuncture combined with herbal medicine is commonly used to treat various complications of diabetes, such as diabetic nephropathy,<sup>74</sup> diabetic peripheral neuropathy,<sup>75</sup> diabetic stroke,<sup>76</sup> and diabetic cardiovascular diseases<sup>77</sup> and so on.

## Exploring the Mechanisms of Acupuncture and Medicine Combination Therapy for Abnormal Glucose Metabolism

### Potential Mechanisms of Synergistic Enhancement

AMP-activated protein kinase (AMPK) has garnered significant attention as a potential therapeutic target for metabolic disorders. Mammalian AMPK is activated in response to decreased cellular energy levels, indicated by an increase in AMP/ATP and ADP/ATP ratios. Activation of AMPK can transcriptionally regulate metabolic pathways, enhancing glucose uptake and utilization, and inhibiting hepatic glycogen synthesis, thereby lowering blood glucose levels.<sup>78</sup> Earlier high-quality studies have demonstrated that metformin, when combined with PEN2, initiates signaling pathways where AMPK activation intersects with ATP6AP1, specifically involving the lysosomal glucose sensing pathway,<sup>23</sup> Metformin induces energy stress by inhibiting mitochondrial respiratory complex I, thereby altering ATP/AMP ratios and activating the AMPK pathway. This process inhibits hepatic gluconeogenesis, enhances insulin sensitivity in the liver, and increases glucose uptake in muscles, thereby achieving glycemic control.<sup>79,80</sup>

Furthermore, numerous studies have indicated that electroacupuncture or acupuncture can improve insulin resistance and regulate energy metabolism by activating AMPK and related pathways. For example, electroacupuncture intervention improves lipid metabolism in middle-aged and elderly obese rats, promotes browning of white adipose tissue,

possibly through activation of the AMPK/Sirt1 signaling pathway, and upregulation of Nrg4 levels.<sup>81</sup> Electroacupuncture effectively improves insulin resistance in Zucker diabetic fatty (ZDF) rats and regulates energy metabolism. Potential mechanisms underlying these effects include modulation of AMPK/mTORC1/p70S6K signaling and related molecules.<sup>82</sup> Moxibustion at ST36 improves lipid levels in Sprague-Dawley (SD) rats with hyperlipidemia by activating the AMPK/mTOR signaling pathway. This activation initiates the transcription of autophagy genes like LC3 and promotes lysosomal synthesis, thereby enhancing autophagy levels in liver cells and improving hepatic fat accumulation.<sup>83</sup> Electroacupuncture effectively promotes browning of white adipose tissue, which may involve several mechanisms. These include activation of GLP-1 neurons in the solitary nucleus, promotion of AMPK phosphorylation in the hypothalamic arcuate nucleus, and upregulation of UCP1.<sup>84</sup> Electroacupuncture can enhance brain glucose metabolism by promoting AMPK $\alpha$  phosphorylation in the caudate putamen (CPu), motor cortex (MCTX), and somatosensory cortex (SCTX) regions.<sup>85</sup> Electroacupuncture can increase insulin sensitivity in insulin-resistant rats, with one of the mechanisms potentially involving the positive regulation of the skeletal muscle AMPK/ACC pathway.<sup>86</sup> Electroacupuncture can improve lipid metabolism disorders in insulin-resistant rats, potentially by reducing the activity of enzymes involved in fatty acid synthesis and regulating the AMPK/p38MAPK/PPAR $\gamma$  signaling pathway to enhance insulin sensitivity.<sup>87</sup> Electroacupuncture can improve diet-induced insulin resistance, likely through the activation of the AMPK signaling pathway in skeletal muscle.<sup>88</sup>

By recognizing that both metformin and electroacupuncture activate the AMPK signaling pathway through common mechanisms, we can hypothesize that acupuncture enhances the glucose-lowering effects of metformin by aiding in the activation of the AMPK pathway.

It is well-known that glucagon-like peptide-1 receptor agonists (GLP-1RAs), such as semaglutide, liraglutide, and dulaglutide, target GLP-1 receptors to stimulate insulin secretion from pancreatic  $\beta$ -cells while inhibiting glucagon secretion, thereby achieving glycemic control and effectively reducing body weight.<sup>89</sup>

Multiple high-quality clinical and preclinical studies suggest that acupuncture may improve obesity and insulin resistance through GLP-1 receptor activation. Preclinical studies have shown that electroacupuncture activates GLP-1 receptors via the GLP-1–VTADA reward pathway, alleviating obesity and insulin resistance.<sup>90</sup> Clinical trials further demonstrate that acupuncture-activated acupoints contain somatic afferent nerves that send sensory signals to neurons in the spinal cord, brainstem, and hypothalamus, activating GLP-1 receptors to produce glucose-lowering effects.<sup>91</sup> Additionally, electroacupuncture has been observed to help restore islet morphology, reduce fasting blood glucose levels, and decrease insulin resistance markers.<sup>92</sup>

Given the shared GLP-1 receptor activation mechanism between GLP-1 receptor agonists and acupuncture, we hypothesize that acupuncture may enhance the glucose-lowering effects of GLP-1RAs by supporting GLP-1 receptor activation.

## Possible Mechanisms of Reducing Toxicity

Common side effects of metformin include gastrointestinal reactions such as nausea, vomiting, and diarrhea. These symptoms may be related to metformin's impact on the energy metabolism of intestinal cells. Activation of AMPK can influence the metabolic function of intestinal cells, potentially contributing to these side effects.

Studies have demonstrated that electroacupuncture at Zhongwan, Tianshu, and Shangjuxu can effectively alleviate symptoms such as diarrhea in rats with ulcerative colitis. This effect may be attributed to the promotion of autophagy in the colon, increased levels of AMPK phosphorylation, and reduced levels of mTOR phosphorylation.<sup>93</sup> Another study suggests that regulating the AMPK/TSC2/Rheb-mediated mTOR inhibition pathway to alter ghrelin levels is a significant mechanism for electroacupuncture in the treatment of functional dyspepsia.<sup>94</sup> Electroacupuncture at ST36 can promote gastrointestinal motility in rats with functional dyspepsia, potentially by downregulating the AMPK/ULK1 signaling pathway, which inhibits excessive autophagy of interstitial cells of Cajal.<sup>88</sup>

We have reason to believe that electroacupuncture alleviates the gastrointestinal reactions caused by metformin through regulation of the AMPK pathway.

Among the numerous neurotransmitters involved in the digestive process, serotonin (5-HT) and 5-hydroxytryptophan (5-HTP) are known to play crucial roles. Studies indicate that metformin shares structural similarities with known

selective 5-hydroxytryptamine 3 (5-HT<sub>3</sub>) receptor agonists.<sup>95</sup> Metformin induces the non-dependent release of 5-HT<sub>3</sub> receptors in the duodenal mucosa through both neuronal and non-neuronal mechanisms. This can lead to gastrointestinal reactions such as nausea, vomiting, and diarrhea.<sup>96</sup>

A basic experiment has demonstrated that electroacupuncture can downregulate the expression of 5-HT<sub>1A</sub> receptors (5-HT<sub>1AR</sub>) and c-fos protein in the hypothalamus and colon tissues of rats with functional dyspepsia.<sup>97</sup> Another head-to-head trial has demonstrated that both electroacupuncture and moxibustion treatment significantly reduce the expression of serotonin (5-HT), 5-HT<sub>3</sub> receptors (5-HT<sub>3R</sub>), and 5-HT<sub>4</sub> receptors (5-HT<sub>4R</sub>) in the mucosal tissues of the sigmoid colon in patients with diarrhea-predominant irritable bowel syndrome. These treatments were associated with significant improvements in abdominal pain and bloating.<sup>98</sup>

We have reason to believe that electroacupuncture alleviates gastrointestinal reactions caused by metformin by potentially reducing elevated levels of intestinal serotonin (5-HT) following metformin intake.

Additionally, metformin may inhibit bile acid reabsorption by altering the function of sodium-dependent bile acid transporters, thereby increasing the pool of bile acids in the intestine.<sup>99</sup> This alteration in bile acid reabsorption may affect stool consistency, secretion of GLP-1, cholesterol levels, and the composition of the microbiota,<sup>35</sup> which could be a direct cause of metformin-induced diarrhea.

Nuclear receptors (NRs) play crucial roles in bile acid signaling. Studies suggest that acupuncture can modulate changes in relevant NRs and influence the circulation of bile acids.<sup>100</sup> Mild moxibustion combined with separated moxibustion has been shown to improve colonic symptoms in rats with segmental enteritis, suppress inflammation, and repair colonic mucosal damage. Additionally, moxibustion can improve abnormal expression of bile acids (BAs) in the colon, liver, and serum.<sup>101</sup>

We have reason to believe that acupuncture alleviates gastrointestinal reactions caused by metformin by potentially regulating abnormal expression of nuclear receptors (NRs) induced by metformin and modulating bile acid circulation.

Other studies suggest that gastrointestinal adverse effects caused by metformin may be associated with changes in the intestinal microbiota induced by the medication.<sup>102</sup> Metformin may affect the intestinal microbiota by potentially regulating inflammation, intestinal permeability, glucose homeostasis, and the abundance of bacteria producing short-chain fatty acids.<sup>103</sup> An analysis focusing solely on the therapeutic impact of metformin reveals a notable rise in Enterobacteriaceae levels and a decrease in Enterococcaceae. These shifts in microbiota are closely linked to gastrointestinal side effects associated with metformin treatment. These effects include diminished absorption of intestinal lipids, localized inflammation triggered by lipopolysaccharides, and increased butyrate production. Notably, elevated butyrate levels have been observed to enhance the expression of genes involved in intestinal gluconeogenesis, potentially directly contributing to the gastrointestinal reactions induced by metformin.<sup>104</sup>

Several studies have demonstrated acupuncture's efficacy in alleviating gastrointestinal symptoms such as nausea, vomiting, and diarrhea. Additionally, a retrospective study suggests that acupuncture may ameliorate abdominal discomfort associated with irritable bowel syndrome by influencing the composition of intestinal microbiota.<sup>105</sup> Furthermore, several basic and clinical studies indicate that acupuncture can alleviate various gastrointestinal discomfort symptoms by regulating the intestinal microbiota.<sup>106–109</sup>

There is reason to believe that acupuncture alleviates gastrointestinal reactions caused by metformin by modulating metformin-induced changes in gut microbiota.

GLP-1 receptor agonists, such as semaglutide, may lead to gastrointestinal adverse effects like nausea and vomiting due to their direct effects on the gastrointestinal tract and the delay in gastric emptying. Studies suggest that the delayed gastric emptying induced by semaglutide may be associated with the inhibition of gastrin, motilin, and cholecystokinin secretion. Gastrin promotes gastric acid secretion, stimulates gastrointestinal smooth muscle, and enhances gastrointestinal motility;<sup>110</sup> motilin initiates the “migrating motor complex” (MMC), which helps clear residuals from the stomach and small intestine, thereby promoting gastric emptying;<sup>111</sup> cholecystokinin promotes bile secretion, facilitates gastric emptying, and regulates appetite.<sup>112</sup>

Multiple basic studies have shown that electroacupuncture at the Zusanli (ST36) acupoint can improve gastric motility in stress-induced rats by increasing levels of vasoactive intestinal peptide (VIP) and nitric oxide (NO).<sup>113</sup> Additionally, acupuncture at local and distal points can exert antagonistic or synergistic effects on specific physiological

and biochemical indicators related to gastric motility, such as ghrelin, gastrin, and growth factor-1, thus alleviating symptoms of nausea and vomiting.<sup>114</sup> Acupuncture may also relieve nausea and vomiting by promoting cholecystokinin secretion, enhancing bile secretion, and indirectly stimulating gastrointestinal motility.<sup>115,116</sup>

We believe that acupuncture may improve GLP-1RA-induced gastrointestinal reactions by modulating changes in gastrin, motilin, and cholecystokinin levels caused by GLP-1RA use.

## Discussion and Prospects

Acupuncture, a traditional Chinese medical therapy, offers substantial benefits when combined with conventional drug treatments. Research indicates that integrating acupuncture with medication effectively enhances blood glucose control in patients with diabetes. This combined approach not only improves medication tolerance but also reduces the occurrence of common drug side effects, such as gastrointestinal discomfort and hypoglycemia.<sup>78</sup> Acupuncture may specifically enhance drug efficacy through various pathways, including activation of the AMPK pathway, a critical energy sensor in cellular metabolism. Additionally, acupuncture has shown potential to mitigate gastrointestinal side effects of medications by regulating serotonin (5-HT) levels, bile acids, and the composition of intestinal microbiota. Understanding these mechanisms is crucial for elucidating the role of acupuncture combined with medication in diabetes treatment. Furthermore, acupuncture has shown potential in managing other conditions associated with glucose metabolism disorders, such as obesity and metabolic syndrome. Studies indicate that acupuncture significantly contributes to weight control in obese patients<sup>117</sup> and improves metabolic syndrome,<sup>118</sup> suggesting its broader applicability in managing metabolic diseases. According to the latest research advancements in 2024, acupuncture may help regulate weight and improve metabolic health by influencing neurocircuits within metabolic pathways,<sup>119</sup> offering new therapeutic approaches for a variety of metabolism-related diseases.

However, as an adjunct therapy, acupuncture also presents certain limitations and challenges. For instance, its therapeutic outcomes may vary significantly among individuals, and the standardization of personalized treatment protocols remains difficult.<sup>120</sup> Future research should therefore weigh its potential efficacy against practical limitations to thoroughly assess its suitability across diverse patient groups. Additionally, acupuncture therapy may increase treatment duration and resource demands, which are important factors to consider in clinical application. In the future, further research should focus on several key areas. First, addressing the standardization and normalization of acupuncture combined with drug therapy in clinical applications requires additional studies on acupuncture protocol standardization, optimal treatment duration, and efficacy evaluation to enable clinicians to implement acupuncture-drug combination therapy more effectively in practice.<sup>121</sup> Second, further investigation into the mechanisms of acupuncture combined with different categories of drugs and the differential effects of these combination therapies across various types of diabetes and patient profiles is essential. Moreover, large-scale, multicenter clinical trials are necessary to validate the long-term efficacy and safety of acupuncture combined with medication in sustained glycemic control, complication prevention, and in the treatment of insulin resistance and insulin resistance-related diseases. Expanding research in these areas will also help explore acupuncture's potential applicability in treating other metabolic disorders.

In conclusion, acupuncture combined with drug therapy represents a comprehensive approach that enhances drug efficacy and reduces the incidence of side effects in patients with type 2 diabetes. This combined treatment shows promising clinical application prospects. With a deeper understanding of its mechanisms and growing clinical evidence, acupuncture-drug combination therapy is poised to gain wider recognition and adoption in the management of diabetes and related metabolic disorders. This approach offers new perspectives and methods for improving patient outcomes in these conditions.

## Ethics Statement

This medical review does not require approval from an ethics committee due to its nature and scope. It primarily involves a synthesis and analysis of existing medical literature and research, without conducting any original research studies or clinical trials that would require ethical review. As a result, the ethical considerations that typically apply to research studies do not apply to this medical review.

## Author Contributions

Wenjun Wang and Lu Zhang are co-first authors. All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

## Funding

This work was supported by Fundamental Research Funds for the China Academy of Chinese Medical Sciences (BJZYB-2023-28).

## Disclosure

The authors declare no conflict of interest.

## References

1. Ormazabal V, Nair S, Elfeky O, Aguayo C, Salomon C, Zuñiga FA. Association between insulin resistance and the development of cardiovascular disease. *Cardiovasc Diabetol*. 2018;17(1):122. doi:10.1186/s12933-018-0762-4
2. Du H, Zhao Y, Yin Z, Wang DW, Chen C. The role of miR-320 in glucose and lipid metabolism disorder-associated diseases. *Int J Biol Sci*. 2021;17(2):402–416. doi:10.7150/ijbs.53419
3. Ong KL, Stafford LK, McLaughlin SA, et al. Global, regional, and national burden of diabetes from 1990 to 2021, with projections of prevalence to 2050: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet*. 2023;402(10397):203–234.
4. Taylor R. Type 2 diabetes: etiology and reversibility. *Diabetes Care*. 2013;36(4):1047–1055. doi:10.2337/dc12-1805
5. Zheng Y, Ley SH, Hu FB. Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. *Nat Rev Endocrinol*. 2018;14(2):88–98. doi:10.1038/nrendo.2017.151
6. Magkos F, Hjorth MF, Astrup A. Diet and exercise in the prevention and treatment of type 2 diabetes mellitus. *Nat Rev Endocrinol*. 2020;16(10):545–555. doi:10.1038/s41574-020-0381-5
7. Carbone S, Del Buono MG, Ozemek C, Lavie CJ. Obesity, risk of diabetes and role of physical activity, exercise training and cardiorespiratory fitness. *Prog Cardiovasc Dis*. 2019;62(4):327–333. doi:10.1016/j.pcad.2019.08.004
8. Andreadis P, Karagiannis T, Malandris K, et al. Semaglutide for type 2 diabetes mellitus: a systematic review and meta-analysis. *Diabetes Obes Metab*. 2018;20(9):2255–2263. doi:10.1111/dom.13361
9. Li Q, Ren Q, Luo Q, et al. Research trends of acupuncture therapy on postoperative nausea and vomiting from 2011 to 2023: a bibliometric analysis. *Complement Ther Med*. 2023;78:102987. doi:10.1016/j.ctim.2023.102987
10. Hahm TS. Electroacupuncture. *Korean J Anesthesiol*. 2009;57(1):3–7. doi:10.4097/kjae.2009.57.1.3
11. Tan MY, Shu SH, Liu RL, Zhao Q. The efficacy and safety of complementary and alternative medicine in the treatment of nausea and vomiting during pregnancy: a systematic review and meta-analysis. *Front Public Health*. 2023;11:1108756. doi:10.3389/fpubh.2023.1108756
12. Park MS, Park WS, Nam D, Min SY, Chae S. Efficacy of electroacupuncture in preventing nausea and vomiting after thyroidectomy: a prospective randomized controlled trial. *Asian J Surg*. 2023;46(9):3480–3484. doi:10.1016/j.asjsur.2022.10.080
13. Wu XK, Gao JS, Ma HL, et al. Acupuncture and doxylamine-pyridoxine for nausea and vomiting in pregnancy: a randomized, controlled, 2 × 2 factorial trial. *Ann Intern Med*. 2023;176(7):922–933. doi:10.7326/M22-2974
14. Kim YH, Kim KS, Lee HJ, Shim JC, Yoon SW. The efficacy of several neuromuscular monitoring modes at the P6 acupuncture point in preventing postoperative nausea and vomiting. *Anesth Analg*. 2011;112(4):819–823. doi:10.1213/ANE.0b013e31820f819e
15. LaMoia TE, Shulman GI. Cellular and molecular mechanisms of metformin action. *Endocr Rev*. 2021;42(1):77–96. doi:10.1210/endo/bnaa023
16. Tomlinson B, Li YH, Chan P. Evaluating gliclazide for the treatment of type 2 diabetes mellitus. *Expert Opin Pharmacother*. 2022;23(17):1869–1877. doi:10.1080/14656566.2022.2141108
17. Jacobsen LV, Flint A, Olsen AK, Ingwersen SH. Liraglutide in type 2 diabetes mellitus: clinical pharmacokinetics and pharmacodynamics. *Clin Pharmacokinet*. 2016;55(6):657–672. doi:10.1007/s40262-015-0343-6
18. Heerspink HJL, Kosiborod M, Inzucchi SE, Cherney DZI. Renoprotective effects of sodium-glucose cotransporter-2 inhibitors. *Kidney Int*. 2018;94(1):26–39. doi:10.1016/j.kint.2017.12.027
19. Charpentier G, Riveline JP, Dardari D, Varroud-Vial M. Should postprandial hyperglycaemia in prediabetic and type 2 diabetic patients be treated? *Drugs*. 2006;66(3):273–286. doi:10.2165/00003495-200666030-00001
20. Heise T. The future of insulin therapy. *Diabet Res Clin Pract*. 2021;175:108820. doi:10.1016/j.diabres.2021.108820
21. Arvind A, Memel ZN, Philpotts LL, Zheng H, Corey KE, Simon TG. Thiazolidinediones, alpha-glucosidase inhibitors, meglitinides, sulfonyleureas, and hepatocellular carcinoma risk: a meta-analysis. *Metabolism*. 2021;120:154780. doi:10.1016/j.metabol.2021.154780
22. Foretz M, Guigas B, Viollet B. Understanding the glucose regulatory mechanisms of metformin in type 2 diabetes mellitus. *Nat Rev Endocrinol*. 2019;15(10):569–589. doi:10.1038/s41574-019-0242-2
23. Ma T, Tian X, Zhang B, et al. Low-dose metformin targets the lysosomal AMPK pathway through PEN2. *Nature*. 2022;603(7899):159–165. doi:10.1038/s41586-022-04431-8
24. Wang Z, Wang J, Hu J, Chen Y, Dong B, Wang Y. A comparative study of acarbose, vildagliptin and saxagliptin intended for better efficacy and safety on type 2 diabetes mellitus treatment. *Life Sci*. 2021;274:119069. doi:10.1016/j.lfs.2021.119069
25. Yang K, Cao F, Wang W, Tian Z, Yang L. The relationship between HMGB1 and autophagy in the pathogenesis of diabetes and its complications. *Front Endocrinol*. 2023;14:1141516. doi:10.3389/fendo.2023.1141516

26. Ge X, Wang L, Fei A, Ye S, Zhang Q. Research progress on the relationship between autophagy and chronic complications of diabetes. *Front Physiol.* 2022;13:956344. doi:10.3389/fphys.2022.956344
27. Hanley DA, Adachi JD, Bell A, Brown V. Denosumab: mechanism of action and clinical outcomes. *Int J Clin Pract.* 2012;66(12):1139–1146. doi:10.1111/ijcp.12022
28. Zhang H, Bu P, Xie YH, et al. Effect of repaglinide and gliclazide on glycaemic control, early-phase insulin secretion and lipid profiles in. *Chin Med J.* 2011;124(2):172–176.
29. Wang B, Zhao J, Zhan Q, et al. Acarbose for postprandial hypotension with glucose metabolism disorders: a systematic review and meta-analysis. *Front Cardiovasc Med.* 2021;8:663635. doi:10.3389/fcvm.2021.663635
30. Karagiannis T, Avgerinos I, Liakos A, et al. Management of type 2 diabetes with the dual GIP/GLP-1 receptor agonist tirzepatide: a systematic review and meta-analysis. *Diabetologia.* 2022;65(8):1251–1261. doi:10.1007/s00125-022-05715-4
31. Rosenstock J, Frias J, Jastreboff AM, et al. Retatrutide, a GIP, GLP-1 and glucagon receptor agonist, for people with type 2 diabetes: a randomised, double-blind, placebo and active-controlled, parallel-group, Phase 2 trial conducted in the USA. *Lancet.* 2023;402(10401):529–544. doi:10.1016/S0140-6736(23)01053-X
32. Frias JP, Davies MJ, Rosenstock J, et al. Tirzepatide versus semaglutide once weekly in patients with type 2 diabetes. *N Engl J Med.* 2021;385(6):503–515. doi:10.1056/NEJMoa2107519
33. Takemori H, Hamamoto A, Isogawa K, et al. Mouse model of metformin-induced diarrhea. *BMJ Open Diabetes Res Care.* 2020;8(1):e000898. doi:10.1136/bmjdr-2019-000898
34. Şahin K, Şahintürk Y, Köker G, et al. Metformin with versus without concomitant probiotic therapy in newly diagnosed patients with type 2 diabetes or prediabetes: a comparative analysis in relation to glycemic control, gastrointestinal side effects, and treatment compliance. *Turk J Gastroenterol.* 2022;33(11):925–933. doi:10.5152/tjg.2022.211063
35. McCreight LJ, Bailey CJ, Pearson ER. Metformin and the gastrointestinal tract. *Diabetologia.* 2016;59(3):426–435. doi:10.1007/s00125-015-3844-9
36. Zullo AR, Riestler MR, Hayes KN, Munshi MN, Berry SD. Comparative safety of sulfonylureas among U.S. nursing home residents. *J Am Geriatr Soc.* 2023;71(4):1047–1057. doi:10.1111/jgs.18160
37. Hantson P, El Balkhi S, Haufroid V, Laterre PF. Sustained hypoglycemia with therapeutic use of repaglinide. *Acta Diabetol.* 2020;57(6):751–753. doi:10.1007/s00592-019-01456-w
38. Wei Y, Lin FJ, Lin SY, Wang CC. Risk of hypoglycemia and concomitant use of repaglinide and clopidogrel: a population-based nested case-control study. *Clin Pharmacol Ther.* 2019;106(6):1346–1352. doi:10.1002/cpt.1556
39. Amiel SA. The consequences of hypoglycaemia. *Diabetologia.* 2021;64(5):963–970. doi:10.1007/s00125-020-05366-3
40. Vallianou NG, Stratigou T, Tsagarakis S. Metformin and gut microbiota: their interactions and their impact on diabetes. *Hormones.* 2019;18(2):141–144. doi:10.1007/s42000-019-00093-w
41. Diaz-Perdigones CM, Muñoz-Garach A, Álvarez-Bermúdez MD, Moreno-Indias I, Tinahones FJ. Gut microbiota of patients with type 2 diabetes and gastrointestinal intolerance to metformin differs in composition and functionality from tolerant patients. *Biomed Pharmacother.* 2022;145:112448. doi:10.1016/j.biopha.2021.112448
42. Bao P, Mi J, Yu Z, et al. Efficacy and safety of acupuncture combined with Chinese herbal medicine in the treatment of type 2 diabetes mellitus: a protocol for a systematic review and meta-analysis. *Medicine.* 2021;100(43):e27658. doi:10.1097/MD.00000000000027658
43. Li SQ, Chen JR, Liu ML, Wang YP, Zhou X, Sun X. Effect and safety of acupuncture for type 2 diabetes mellitus: a systematic review and meta-analysis of 21 randomised controlled trials. *Chin J Integr Med.* 2022;28(5):463–471. doi:10.1007/s11655-021-3450-2
44. Chen C, Liu J, Sun M, Liu W, Han J, Wang H. Acupuncture for type 2 diabetes mellitus: a systematic review and meta-analysis of randomized controlled trials. *Complement Ther Clin Pract.* 2019;36:100–112. doi:10.1016/j.ctcp.2019.04.004
45. Wei JP, Wang QH, Zheng HJ, Wei F. Research progress on non-drug treatment for blood glucose control of type 2 diabetes mellitus. *Chin J Integr Med.* 2018;24(10):723–727. doi:10.1007/s11655-018-2844-2
46. Li X, Jia HX, Yin DQ, Zhang ZJ. Acupuncture for metabolic syndrome: systematic review and meta-analysis. *Acupunct Med.* 2021;39(4):253–263. doi:10.1177/0964528420960485
47. Xiao E, Luo L. Alternative therapies for diabetes: a comparison of western and Traditional Chinese Medicine (TCM) approaches. *Curr Diabetes Rev.* 2018;14(6):487–496. doi:10.2174/1573399813666170519103230
48. Zhou W, Lee A, Zhou A, Lombardo D. Integrative care: acupuncture based neuromodulation therapy for diabetes and heart failure. *Front Neurosci.* 2024;18:1332957. doi:10.3389/fnins.2024.1332957
49. Wen Q, Hu M, Lai M, et al. Effect of acupuncture and metformin on insulin sensitivity in women with polycystic ovary syndrome and insulin resistance: a three-armed randomized controlled trial. *Hum Reprod.* 2022;37(3):542–552. doi:10.1093/humrep/deab272
50. Xihui Q, Jianli P, Guan X, Jie F. Bo's abdominal acupuncture improves disordered metabolism in obese type 2 diabetic rats through regulating fibroblast growth factor 21 and its related adipokines. *J Tradit Chin Med.* 2023;43(6):1200–1208. doi:10.19852/j.cnki.jtcm.20231008.002
51. Wang H, Chen X, Chen C, et al. Electroacupuncture at lower he-sea and front-mu acupoints ameliorates insulin resistance in type 2 diabetes mellitus by regulating the intestinal flora and gut barrier. *Diabetes Metab Syndr Obes.* 2022;15:2265–2276. doi:10.2147/DMSO.S374843
52. An J, Wang L, Song S, et al. Electroacupuncture reduces blood glucose by regulating intestinal flora in type 2 diabetic mice. *J Diabetes.* 2022;14(10):695–710. doi:10.1111/1753-0407.13323
53. Dong Y, Li R, Guo WX, et al. Effect of electroacupuncture on liver Akt/FoxO1 signaling pathway in rats with diabetic fatty. *Zhongguo Zhen Jiu.* 2023;43(6):679–683. doi:10.13703/j.0255-2930.20220505-k0006
54. Duan HR, Li R, Song SS, Hu SQ, Zhuang ST, Li QY. Electroacupuncture improves glucose and lipid metabolism by regulating APN/AMPK/PPAR $\alpha$  signaling of skeletal muscle in Zucker diabetic obese rats. *Zhen Ci Yan Jiu.* 2021;46(11):907–913. doi:10.13702/j.1000-0607.20210201
55. Shen XT, Zhang SS, Chen XY, Yu Z, Xu B. Hypoglycemic effect of electroacupuncture at “Tianshu” (ST 25) combined with metformin on rats with type 2 diabetes mellitus based on AMPK. *Zhongguo Zhen Jiu.* 2023;43(1):53–59. doi:10.13703/j.0255-2930.20211106-k0003
56. Liu Y, Xu TC, Yu Z, Xu B. Investigation on the mechanism of acupuncture in treatment of diabetes mellitus type 2 based on the network of islet macrophages-pancreatic adipose cells-islet  $\beta$  cells. *Zhongguo Zhen Jiu.* 2022;42(4):433–436. doi:10.13703/j.0255-2930.20210126-k0004
57. Yin G, Shen GM, Jiang AJ, Li JY. Acupuncture intervention induced improvement of oxidative stress by regulating PKC $\beta$ /P66shc signaling in obese diabetic rats. *Zhen Ci Yan Jiu.* 2021;46(8):642–648. doi:10.13702/j.1000-0607.201318

58. Yang NN, Lin LL, Li YJ, et al. Potential mechanisms and clinical effectiveness of acupuncture in depression. *Curr Neuropharmacol.* 2022;20(4):738–750. doi:10.2174/1570159X19666210609162809
59. Wang J, Liao Y, You Y, et al. Acupuncture and Chinese herbal medicine for menopausal mood disorder: a randomized controlled trial. *Climacteric.* 2023;26(4):392–400. doi:10.1080/13697137.2023.2187284
60. Xu T, Yu Z, Liu Y, et al. Hypoglycemic effect of electroacupuncture at ST25 through neural regulation of the pancreatic intrinsic nervous system. *Mol Neurobiol.* 2022;59(1):703–716. doi:10.1007/s12035-021-02609-1
61. Liao HY, Sun MF, Lin JG, Chang SL, Lee YC. Electroacupuncture plus metformin lowers glucose levels and facilitates insulin sensitivity by activating MAPK in steroid-induced insulin-resistant rats. *Acupunct Med.* 2015;33(5):388–394. doi:10.1136/acupmed-2014-010724
62. Chung YC, Chen YI, Lin CM, et al. Electroacupuncture combined with acarbose improves insulin sensitivity via peroxisome proliferator-activated receptor  $\gamma$  activation and produces a stronger glucose-lowering effect than acarbose alone in a rat model of steroid-induced insulin resistance. *Acupunct Med.* 2020;38(5):335–342. doi:10.1177/0964528419901135
63. Zhu T, Ji L, Gao Z, Ren J, Ji Y, Miao J. Acupotomy combined with metformin hydrochloride tablet for type 2 diabetes mellitus and its effect on serum inflammatory factors. *Zhongguo Zhen Jiu.* 2024;44(3):245–250. doi:10.13703/j.0255-2930.20230705-k0002
64. Chen X, Lan Y, Yang L, et al. Acupuncture combined with metformin versus metformin alone to improve pregnancy rate in polycystic ovary syndrome: a systematic review and meta-analysis. *Front Endocrinol.* 2022;13:978280. doi:10.3389/fendo.2022.978280
65. Liu Y, Fan HY, Hu JQ, Wu TY, Chen J. Effectiveness and safety of acupuncture for insulin resistance in women with polycystic ovary syndrome: a systematic review and meta-analysis. *Heliyon.* 2023;9(3):e13991. doi:10.1016/j.heliyon.2023.e13991
66. Muharam R, Ph D, Srilestari A, Mihardja H, Juvanni Callestya L, Kemal Harzif A. Combination of electroacupuncture and pharmacological treatment improves insulin resistance in women with polycystic ovary syndrome: double-blind randomized clinical trial. *Int J Reprod Biomed.* 2022;20(4):289–298. doi:10.18502/ijrm.v20i4.10900
67. Firouzjaei A, Li GC, Wang N, Liu WX, Zhu BM. Comparative evaluation of the therapeutic effect of metformin monotherapy with metformin and acupuncture combined therapy on weight loss and insulin sensitivity in diabetic patients. *Nutr Diabetes.* 2016;6(5):e209. doi:10.1038/nutd.2016.16
68. Ding N, Li L, Zhu X, Huang X, Wang L, Yue R. Acupuncture for patients with glucagon-like peptide-1 receptor agonists-induced nausea and vomiting: a systematic review protocol. *Medicine.* 2020;99(21):e20343. doi:10.1097/MD.0000000000020343
69. Zhou X, Cao SG, Tan XJ, et al. Effects of transcutaneous electrical acupoint stimulation (TEAS) on postoperative recovery in patients with gastric cancer: a randomized controlled trial. *Cancer Manag Res.* 2021;13:1449–1458. doi:10.2147/CMAR.S292325
70. Sun H, Zhang B, Qian HH, Chen ZC. Effect of warm-needle moxibustion intervention on immune function and intestinal flora in patients after colorectal cancer radical operation. *Zhen Ci Yan Jiu.* 2021;46(7):592–597. doi:10.13702/j.1000-0607.200647
71. Jing X, Ou C, Chen H, et al. Electroacupuncture reduces weight gain induced by rosiglitazone through PPAR $\gamma$  and leptin receptor in CNS. *Evid Based Complement Alternat Med.* 2016;2016:8098561. doi:10.1155/2016/8098561
72. Liu HJ, Hsu SF, Hsieh CC, et al. The effectiveness of Tsu-San-Li (St-36) and Tai-Chung (Li-3) acupoints for treatment of acute liver damage in rats. *Am J Chin Med.* 2001;29(2):221–226. doi:10.1142/S0192415X01000253
73. Zhu KX, Wu M, Bian ZL, et al. Growing attention on the toxicity of Chinese herbal medicine: a bibliometric analysis from 2013 to 2022. *Front Pharmacol.* 2024;15:1293468. doi:10.3389/fphar.2024.1293468
74. Yu Z, Zhang W, Li B, et al. Efficacy and safety of acupuncture combined with Chinese herbal medicine for diabetic nephropathy: a protocol for systematic review and meta-analysis. *Medicine.* 2021;100(35):e27087. doi:10.1097/MD.0000000000027087
75. Shi Y, Liu L, Sun X, Jiao J. Efficacy and safety of acupuncture combined Chinese herbal medicine for diabetic peripheral neuropathy: a protocol for systematic review and meta-analysis. *Medicine.* 2021;100(50):e28086. doi:10.1097/MD.0000000000028086
76. Zhang A, Han F, Piao C. Comparative effectiveness of acupuncture and pharmacological interventions in treating diabetic stroke: a protocol for a systematic review and network meta-analysis. *Medicine.* 2022;101(46):e31823. doi:10.1097/MD.0000000000031823
77. Wang JY, Zhang J, Liu Y, Gao YM, Sun JJ, Liu XM. Acupuncture combined with western medication in treatment of type 2 diabetes mellitus with angina pectoris of coronary heart disease: a randomized controlled study. *Zhongguo Zhen Jiu.* 2021;41(4):371–375. doi:10.13703/j.0255-2930.20200329-k0001
78. Lin SC, Hardie DG. AMPK: sensing glucose as well as cellular energy status. *Cell Metab.* 2018;27(2):299–313. doi:10.1016/j.cmet.2017.10.009
79. Herzig S, Shaw RJ. AMPK: guardian of metabolism and mitochondrial homeostasis. *Nat Rev Mol Cell Biol.* 2018;19(2):121–135. doi:10.1038/nrm.2017.95
80. Liu L, Patnana PK, Nimmagadda SC. Low-dose metformin and PEN2-dependent lysosomal AMPK activation: benefits outnumber side effects. *Signal Transduct Target Ther.* 2022;7(1):178. doi:10.1038/s41392-022-01040-9
81. He XL, Li XZ, Xu DW, Li Y, Yang ZX. Electroacupuncture intervention improves lipid metabolism and promotes browning of white adipose tissue by activating AMPK/Sirt1 pathway and up-regulating Nrg4 content in middle-aged and aged obese rats. *Zhen Ci Yan Jiu.* 2023;48(8):764–772. doi:10.13702/j.1000-0607.20230187
82. Liu XX, Zhang LZ, Zhang HH, et al. Low-frequency electroacupuncture improves disordered hepatic energy metabolism in insulin-resistant Zucker diabetic fatty rats via the AMPK/mTORC1/p70S6K signaling pathway. *Acupunct Med.* 2022;40(4):360–368. doi:10.1177/09645284211070301
83. Xu Q, Wu H, Zhu H, et al. Grain-sized moxibustion at Zusanli (ST36) promotes hepatic autophagy in rats with hyperlipidemia by regulating the ULK1 and TFEB expression through the AMPK/mTOR signaling pathway. *Heliyon.* 2023;9(5):e15316. doi:10.1016/j.heliyon.2023.e15316
84. Zhu Y, Tian J, Shao YW, Zhao J, Jia SH, Shu Q. Electroacupuncture improves obesity and promotes white adipose tissue browning by regulating central glucagon-like peptide-1. *Zhen Ci Yan Jiu.* 2023;48(8):727–735. doi:10.13702/j.1000-0607.20221074
85. Wu J, Lin B, Liu W, et al. Roles of electro-acupuncture in glucose metabolism as assessed by 18F-FDG/PET imaging and AMPK $\alpha$  phosphorylation in rats with ischemic stroke. *Int J Mol Med.* 2017;40(3):875–882. doi:10.3892/ijmm.2017.3057
86. Li Z, Lan D, Zhang H, Zhang H, Chen X, Sun J. Electroacupuncture mitigates skeletal muscular lipid metabolism disorder related to high-fat-diet induced insulin resistance through the AMPK/ACC signaling pathway. *Evid Based Complement Alternat Med.* 2018;2018:7925842. doi:10.1155/2018/7925842
87. Li ZX, Zhang HH, Lan DC, Zhang HT, Sun J. Electroacupuncture improves lipid metabolic disorder by regulating hepatic AMPK/p38 MAPK/RRAR $\gamma$  signaling in rats with high-fat diet-induced insulin resistance. *Zhen Ci Yan Jiu.* 2019;44(1):8–12. doi:10.13702/j.1000-0607.170633
88. Tomimaga A, Ishizaki N, Naruse Y, Kitakoji H, Yamamura Y. Repeated application of low-frequency electroacupuncture improves high-fructose diet-induced insulin resistance in rats. *Acupunct Med.* 2011;29(4):276–283. doi:10.1136/acupmed-2011-010006

89. Yao H, Zhang A, Li D, et al. Comparative effectiveness of GLP-1 receptor agonists on glycaemic control, body weight, and lipid profile for type 2 diabetes: systematic review and network meta-analysis. *BMJ*. 2024;384:e076410. doi:10.1136/bmj-2023-076410
90. Zhu Y, Tian J, Wei X, Jia S, Shu Q. Electroacupuncture alleviates obesity and insulin resistance via the GLP-1-VTADA reward circuit. *Neuroendocrinology*. 2024;114(3):263–278. doi:10.1159/000535068
91. Chen YK, Liu TT, Teia FKF, Xie MZ. Exploring the underlying mechanisms of obesity and diabetes and the potential of Traditional Chinese Medicine: an overview of the literature. *Front Endocrinol*. 2023;14:1218880. doi:10.3389/fendo.2023.1218880
92. Cao B, Li R, Tian H, et al. Effect on glycemia in rats with type 2 diabetes induced by streptozotocin: low-frequency electro-pulse needling stimulated Weiwanshiashu (EX-B 3) and Zusanli (ST 36). *J Tradit Chin Med*. 2016;36(6):768–778. doi:10.1016/S0254-6272(17)30013-4
93. Wu RL, Xu HC, Jiang ZW, et al. Effect of electroacupuncture on colonic autophagy and AMPK/mTOR signaling pathway in rats with acute ulcerative colitis. *Zhen Ci Yan Jiu*. 2023;48(8):818–824. doi:10.13702/j.1000-0607.20220863
94. Tang L, Zeng Y, Li L, et al. Electroacupuncture upregulated ghrelin in rats with functional dyspepsia via AMPK/TSC2/Rheb-mediated mTOR inhibition. *Dig Dis Sci*. 2020;65(6):1689–1699. doi:10.1007/s10620-019-05960-5
95. Bouchoucha M, Uzzan B, Cohen R. Metformin and digestive disorders. *Diabetes Metab*. 2011;37(2):90–96. doi:10.1016/j.diabet.2010.11.002
96. Cubeddu LX, Bönisch H, Göthert M, et al. Effects of metformin on intestinal 5-hydroxytryptamine (5-HT) release and on 5-HT3 receptors. *Naunyn Schmiedebergs Arch Pharmacol*. 2000;361(1):85–91.
97. Wang Q, Wang Y, Niu WM, Yang XH, Liu ZB. Electroacupuncture improves intestinal function by down-regulating expression of 5-HT(1A)R and c-fos proteins in the hypothalamus and colon tissues via brain-gut axis in rats with functional diarrhea. *Zhen Ci Yan Jiu*. 2019;44(7):501–505. doi:10.13702/j.1000-0607.180023
98. Zhao JM, Lu JH, Yin XJ, et al. Comparison of electroacupuncture and moxibustion on brain-gut function in patients with diarrhea-predominant irritable bowel syndrome: a randomized controlled trial. *Chin J Integr Med*. 2015;21(11):855–865. doi:10.1007/s11655-015-2049-x
99. Napolitano A, Miller S, Nicholls AW, et al. Novel gut-based pharmacology of metformin in patients with type 2 diabetes mellitus. *PLoS One*. 2014;9(7):e100778. doi:10.1371/journal.pone.0100778
100. Shen JC, Qi Q, Han D, et al. Role of bile acids and nuclear receptors in acupuncture in improving Crohn's disease. *Evid Based Complement Alternat Med*. 2022;2022:5814048. doi:10.1155/2022/5814048
101. Shen JC, Qi Q, Han D, et al. Moxibustion improves experimental colitis in rats with Crohn's disease by regulating bile acid enterohepatic circulation and intestinal farnesoid X receptor. *J Integr Med*. 2023;21(2):194–204. doi:10.1016/j.joim.2023.01.001
102. Cheng M, Ren L, Jia X, Wang J, Cong B. Understanding the action mechanisms of metformin in the gastrointestinal tract. *Front Pharmacol*. 2024;15:1347047. doi:10.3389/fphar.2024.1347047
103. Lee CB, Chae SU, Jo SJ, Jerng UM, Bae SK. The relationship between the gut microbiome and metformin as a key for treating type 2 diabetes mellitus. *Int J Mol Sci*. 2021;22(7):3566.
104. Whang A, Nagpal R, Yadav H. Bi-directional drug-microbiome interactions of anti-diabetics. *EBioMedicine*. 2019;39:591–602. doi:10.1016/j.ebiom.2018.11.046
105. Yaklai K, Pattanakuhar S, Chattapakorn N, Chattapakorn SC. The role of acupuncture on the gut-brain-microbiota axis in irritable bowel syndrome. *Am J Chin Med*. 2021;49(2):285–314. doi:10.1142/S0192415X21500154
106. Sui H, Zhang L, Gu K, et al. YYFZBJS ameliorates colorectal cancer progression in Apc(Min/+) mice by remodeling gut microbiota and inhibiting regulatory T-cell generation. *Cell Commun Signal*. 2020;18(1):113. doi:10.1186/s12964-020-00596-9
107. Xu MM, Guo Y, Chen Y, Zhang W, Wang L, Li Y. Electro-acupuncture promotes gut motility and alleviates functional constipation by regulating gut microbiota and increasing butyric acid generation in mice. *J Integr Med*. 2023;21(4):397–406. doi:10.1016/j.joim.2023.05.003
108. Li X, He F, Tuo X, et al. Electroacupuncture ameliorates peptic ulcer disease in association with gastroduodenal microbiota modulation in mice. *Front Cell Infect Microbiol*. 2022;12:935681. doi:10.3389/fcimb.2022.935681
109. Cheng J, Li W, Wang Y, et al. Electroacupuncture modulates the intestinal microecology to improve intestinal motility in spinal cord injury rats. *Microb Biotechnol*. 2022;15(3):862–873. doi:10.1111/1751-7915.13968
110. Schubert ML, Rehfeld JF. Gastric Peptides-Gastrin and Somatostatin. *Compr Physiol*. 2019;10(1):197–228.
111. Kitazawa T, Kaiya H. Motilin comparative study: structure, distribution, receptors, and gastrointestinal motility. *Front Endocrinol*. 2021;12:700884. doi:10.3389/fendo.2021.700884
112. Liddle RA. Cholecystokinin cells. *Annu Rev Physiol*. 1997;59:221–242. doi:10.1146/annurev.physiol.59.1.221
113. Shen GM, Zhou MQ, Xu GS, Xu Y, Yin G. Role of vasoactive intestinal peptide and nitric oxide in the modulation of electroacupuncture on gastric motility in stressed rats. *World J Gastroenterol*. 2006;12(38):6156–6160. doi:10.3748/wjg.v12.i38.6156
114. Yu B, Sun M, Wang Z, et al. Effects of stimulating local and distal acupoints on diabetic gastroparesis: a new insight in revealing acupuncture therapeutics. *Am J Chin Med*. 2021;49(5):1151–1164. doi:10.1142/S0192415X21500555
115. Chang J, Liu Y, Jiang TC, Zhao L, Liu JW. Cholecystokinin and cholecystokinin-A receptor: an attractive treatment strategy for biliary dyskinesia? *World J Gastroenterol*. 2024;30(3):283–285. doi:10.3748/wjg.v30.i3.283
116. Güçel F, Bahar B, Demirtas C, Mit S, Cevik C. Influence of acupuncture on leptin, ghrelin, insulin and cholecystokinin in obese women: a randomised, sham-controlled preliminary trial. *Acupunct Med*. 2012;30(3):203–207. doi:10.1136/acupmed-2012-010127
117. Bian Z, Ren L, Bian J. Research trends of traditional Chinese non-pharmacological therapy in the management of overweight and obesity from 2004 to 2023: a bibliometric study. *Complement Ther Med*. 2024;87:103099. doi:10.1016/j.ctim.2024.103099
118. Stańczak NA, Grywalska E, Dudzińska E. The latest reports and treatment methods on polycystic ovary syndrome. *Ann Med*. 2024;56(1):2357737. doi:10.1080/07853890.2024.2357737
119. Jiang LY, Tian J, Yang YN, Jia SH, Shu Q. Acupuncture for obesity and related diseases: insight for regulating neural circuit. *J Integr Med*. 2024;22(2):93–101. doi:10.1016/j.joim.2024.03.001
120. Hong SH, Wu F, Ding SS, et al. Current status of standardization of acupuncture and moxibustion in China. *Qjm*. 2014;107(3):173–178. doi:10.1093/qjmed/hct240
121. Schnyer RN, Allen JJ. Bridging the gap in complementary and alternative medicine research: manualization as a means of promoting standardization and flexibility of treatment in clinical trials of acupuncture. *J Altern Complement Med*. 2002;8(5):623–634. doi:10.1089/10755302320825147

Diabetes, Metabolic Syndrome and Obesity

Dovepress

### Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-journal>