



Management and Outcomes of Low-Grade Gliomas in Africa: A Scoping Review Protocol

SETTHASORN ZHI YANG OOI

ROSALINE DE KONING

ABDULLAH EGIZ

DAVID ULRICH DALLE

MOUSSA DENOUE

MARVIN RICHIE DONGMO

TSOPMENE

MEHDI KHAN

RÉGIS TAKOUKAM

JAY KOTECHA

DAWIN SICHIMBA

YAO CHRISTIAN HUGUES DOKPONOU

ULRICK SIDNEY KANMOUNYE

NOUROU DINE ADENIRAN BANKOLE

*Author affiliations can be found in the back matter of this article

ABSTRACT

Background: Over the last decade, many advancements have been made in the management of low-grade gliomas (LGGs). Overall survival outcomes are correlated with factors such as postoperative residual volumes and specific tumor biomolecular profiles such as IDH mutation status. It is unclear whether these advancements have benefited LGG patients in Africa. This scoping review protocol outlines how the authors will evaluate the epidemiology, presentations, management and outcomes of LGGs in Africa.

Methods: MEDLINE, Embase and African Journals Online will be searched from database inception to date in order to identify the relevant studies. Patients of all ages with histologically and/or radiologically confirmed LGGs that were managed in an African country will be included. Surgical and chemoradiation management of LGG tumours will be considered. Original research, reviews, commentaries, editorials and case reports will be included.


Results: Primary outcomes of the review will include LGG management, morbidity and mortality. Secondary outcomes include epidemiology and recurrence of LGGs.

Discussion: This scoping review will be the first to evaluate the current landscape of LGG management and outcomes in Africa, highlighting pertinent themes that may be used to guide further research as well as health system strengthening efforts by policymakers and stakeholders.

Scoping Review Registration: The protocol has been registered on the Open Science Framework (OSF; registration link: <https://doi.org/10.17605/OSF.IO/E732G>).

Highlights:

- LGGs account for 17% to 22% of total brain tumours and have a median survival time between 5.6 and 13.3 years.



PROTOCOL



IJS Press

Part of the IJS Publishing Group

CORRESPONDING AUTHOR:

Nourou Dine Adeniran Bankole

Ibn Sina Teaching Hospital,
Rabat Morocco/Mohammed V
University of Rabat Morocco,
Department of Neurosurgery,
Rabat, Morocco

bankolenouroudine@yahoo.fr

KEYWORDS:

low-grade gliomas; Africa;
epidemiology; management;
outcomes; scoping review;
protocol

TO CITE THIS ARTICLE:

Ooi SZY, de Koning R, Egiz A,
Dalle DU, Denou M, Tsopmene
MRD, Khan M, Takoukam
R, Kotecha J, Sichimba D,
Dokponou YCH, Kanmounye
US, Bankole NDA. 2022.
Management and Outcomes
of Low-Grade Gliomas in
Africa: A Scoping Review
Protocol. *International Journal
of Surgery: Protocols*, 26(1),
pp. 1–6. DOI: [https://doi.
org/10.29337/ijsp.171](https://doi.org/10.29337/ijsp.171)

- Despite many recent advancements in the management of LGGs, there is a paucity in the data within the African landscape.
- This scoping review will be the first to evaluate the current landscape of LGG management and outcomes in Africa, highlighting pertinent themes that may be used to guide further research and policymaking efforts.

1. INTRODUCTION

Low-grade gliomas (LGGs) are grade I and II primary brain tumors, according to the World Health Organization (WHO) [1]. LGGs include astrocytomas, oligodendrogliomas and oligoastrocytomas, with the latter being designated as Not Otherwise Specified (NOS) in the new 2016 WHO classification of central nervous system tumours [2]. They develop from glial cells and are characterized by a low proliferation index (median, 4.4 mm/yr) [3]. Nevertheless, they have a high potential of malignant transformation over time and can cause considerable morbidity and lead to death [4]. The annual global age-standardized incidence of primary malignant brain tumors is estimated at 3.7 and 2.6 per 100,000 males and females, respectively [5]. There is a marked increase of these rates in high-income countries (HICs) (men, 5.8 and women, 4.1 per 100,000) more than in low- and middle-income countries (LMICs) (men 3.0 and women 2.1 per 100,000) [5], however this could be due to an under-diagnosis of LGGs in LMICs. In Sub-Saharan Africa, brain tumors represent 0.14% of disability-adjusted life years (DALYs) and 0.17% of deaths, while in the Northern African and the Middle Eastern regions brain tumors cause 0.44% of DALYs and 0.62% of deaths [6]. LGGs represent between 17% to 22% of brain tumors (approximately 20 000 cases per year) and have a median survival time between 5.6 and 13.3 years [7].

Anecdotal evidence suggests that the management of patients with LGGs is challenging in Africa. Poor access to neuroimaging facilities leads to delays in diagnosis and referral [8]. Diagnosis and subsequent management decisions require the use of advanced and costly imaging modalities, which are not available in many resource limited settings. Though basic neuroimaging tools such as computerized tomography (CT) scan and magnetic resonance imaging (MRI) are being acquired widely, they are distributed unevenly, along with the already poor neurosurgical workforce [9]. Gross total resection is also a daunting task because of the absence of supportive technology. Moreover, high out-of-pocket costs and lack of financial risk protection limit access when those facilities are available [10]. In addition, infrastructural deficiencies limit service delivery. For example, only 10% of awake craniotomy LGG operations are done in optimal conditions [9],

and there is a significant deficit in the literature on neurosurgical care for LGGs in Africa [11].

To our knowledge, there is no literature evaluating epidemiology, management and outcomes of patients with LGGs in Africa, hence necessitating the need for a scoping review.

2. AIMS

Primary aims

1. Assess the treatment modality and management plan (i.e., chemotherapy, radiotherapy and intraoperative surgical adjuncts), and
2. Assess the clinical outcome of patients defined as rates of mortality, morbidity, and recurrence.

Secondary aims

1. Assess the epidemiology of LGGs in different African countries,
2. Assess the availability of diagnostic modalities such as histopathology and/or molecular pathology testing and neuroimaging (MRI, CT scan), and
3. Assess the availability of specialized workforce, infrastructure and financial risk protection.

3. METHODS

The review will be conducted per the Arksey and O'Malley framework [12] and reported in accordance with the Preferred Reporting Items for Systematic Review and Meta-Analysis Extension for Scoping Reviews (PRISMA-ScR) guidelines [13]. A scoping review is chosen instead of a systematic review because the evidence relating to the epidemiology, presentation, management and outcomes of LGGs in Africa has not been comprehensively reviewed. Systematic reviews answer a focused research question with narrow parameters, strict endpoints, and eligibility criteria of the included studies defined at the outset, whereas scoping reviews can explore several questions in a broad sense. A scoping review is, therefore, more suitable for our investigation because it remains relatively unclear which specific questions should be asked and valuably addressed by a more precise systematic review and meta-analysis.

This protocol has been developed in accordance with the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) guidelines [14]. The review was registered on Open Science Framework [15]. Protocol amendments will be updated and published alongside the scoping review results (<https://doi.org/10.17605/OSF.IO/E732G>).

3.1. ELIGIBILITY CRITERIA

Inclusion criteria

Included will be any relevant article published in a peer-reviewed journal that discusses the epidemiology, presentation, management and outcomes of LGGs in Africa. Study types including journal articles, reviews, case reports, letters will be included. There will be no restrictions on the age of patients included nor the time period of publication. Publications in English and French languages will be considered.

Exclusion criteria

The following articles will be considered as outside the scope of this work and will be excluded:

- Do not include African patients or do not have disaggregated data about the African population,
- Do not discuss LGGs or do not have disaggregated data about LGGs,
- Do not discuss epidemiology, presentation, management or outcomes of patients with LGGs,
- Are neither written in English nor French, and
- Conference abstracts (due to the lack of in-depth information available).

3.2. INFORMATION SOURCES

The databases to be searched include: MEDLINE, Embase and African Journals Online.

3.3. SEARCH STRATEGY

A search strategy has been developed to identify studies related to the treatment and/or outcomes of patients with LGGs in Africa. Synonyms relating to terms describing individual African countries, LGG, therapy, and outcome will be used (Supplementary Figure 1).

3.4. DATA MANAGEMENT

Data records will be downloaded from respective databases in comma-separated values (CSV) formatted files. They will then be imported into Rayyan [16] where deduplication, title and abstract screening, and full-text screening will take place. Further data extraction and quality assessment will be carried out on Microsoft Excel (Microsoft, Richmond, Virginia, USA).

3.5. STUDY SELECTION

A calibration exercise will be carried out before title and abstract screening in order to ensure adequate

understanding of the inclusion criteria by study screeners. Deduplication will be undertaken on Rayyan. Each study will then be screened using title and abstract, by two independent reviewers, against the pre-defined inclusion and exclusion criteria. Potentially eligible studies will be further screened for full-text review. Disagreements will be discussed amongst the reviewers and in case of no resolution, an appeal will be made to a senior author (USK or NDAB).

3.6. DATA EXTRACTION

Full-text screened articles will be exported into a previously-made data extraction proforma on Microsoft Excel (Microsoft, Richmond, Virginia, USA). Data will be extracted on (i) study design, (ii) patient demographics, (iii) country of origin, (iv) tumour characteristics, (v) neuroimaging modality used, (vi) histopathology diagnosis, (vii) molecular pathology diagnosis, (viii) type of intervention, and (ix) outcomes of care. Data extraction will be performed in two stages, a pilot stage followed by a proper stage. The pilot stage will consist of having multiple authors, each going through the same 10 randomly selected articles to extract data. This is to assure the reliability of the proforma and that all participant authors were able to extract data accurately and homogeneously. Feedback from the pilot stage will inform any necessary changes to be made, upon discussion, in order to accurately capture the pertinent themes in the literature.

3.7. RISK OF BIAS ASSESSMENT

The purpose of this review is to produce a systematically conducted scoping review of the available literature to provide a comprehensive overview on the management and outcomes of LGGs in Africa, hence a formal bias assessment will not be conducted. Given the limited and heterogenous literature body, a formal bias risk assessment was deemed unnecessary for this emerging area of literature, as it will suffer from standard biases associated with new areas of clinical research.

3.8. DATA SYNTHESIS

Study characteristics will be summarised using descriptive statistics and presented in a table. Data relating to study characteristics will be grouped into categories where appropriate. Categorisation might be based on type of LGG tumours, countries/regions within Africa or will be finalised in discussions with the team wherever necessary if notable differences emerge in study findings.

Pertinent characteristics of the study population will be analysed particularly relating to age, all which contribute to the risk of being diagnosed with LGG and more importantly the subsequent outcomes. Particular features of the tumour will be noted, in order to gauge all the factors that have contributed to tumour prognosis in study participants, such as those related to tumour biology (histology and molecular pathology), tumour location, solitary or multiple-lesion tumours, and tumour

recurrence. The intraoperative adjuncts used during surgery and the extent of resection for these tumours will also be highlighted whenever applicable. Any notable surgical adjuncts used in improving the extent of resection will be reported. The treatment outcomes will be described: mortality rate, morbidity rate, and recurrence. An analysis of these outcomes will enable comparison and discussion of effectiveness of treatment techniques adopted in Africa with HICs and other LMICs.

4. ETHICS AND DISSEMINATION

This study will exclusively involve secondary data collection and no human participants will be involved in the design or dissemination of this research, hence ethical approval was not required. The results from this study will be disseminated through a peer-reviewed journal.

5. LIMITATIONS

There is an extensive amount of literature published in Arabic, Portuguese and Spanish that will not be addressed by this review.

6. CONCLUSION

The proposed scoping review aims to detail the current landscape of LGGs in Africa, highlighting parallels and differences among countries within the continent as well as with other countries of different income groups. This novel work will allow for a better understanding of the current situation in Africa, wherein important lessons can be drawn from the robust workforce environment and the use of advanced surgical adjuncts in HICs, and vice-versa from the ingenious thinking that is common in regions with limited resources. This would provide pertinent lessons to be learnt to improve patient outcomes and the quality of care of LGG management, especially given its poor prognosis of progressing to high-grade glioma and eventually death [4].

AMENDMENTS

Any amendments to this protocol will be prospectively updated on Open Science Framework.

ADDITIONAL FILE

The additional file for this article can be found as follows:

- **Supplementary Figure 1.** Search strategy. DOI: <https://doi.org/10.29337/ijsp.171.s1>

ACKNOWLEDGEMENTS

We would like to address special thanks to the Neurology and Neurosurgery Interest Group (NANSIG) and the Association of Future African Neurosurgeons (AFAN), who, in collaboration, made this research project possible. We would also like to thank Mr. Neal Thurley, whose input on the search strategy and methodology was invaluable.

COMPETING INTERESTS

The authors have no competing interests to declare.


AUTHOR CONTRIBUTIONS


Setthasorn Zhi Yang Ooi (project administration, methodology, writing – original draft, writing – review and editing, supervision), Rosaline de Koning (conceptualisation, methodology, writing – original draft, writing – review and editing), Abdullah Egiz (writing – review and editing), David Ulrich Dalle (writing – review and editing), Moussa Denou (writing – review and editing), Marvin Richie Dongmo Tsopmene (writing – review and editing), Mehdi Khan (writing – review and editing), Régis Takoukam (writing – review and editing), Jay Kotecha (writing – review and editing), Dawin Sichimba (writing – review and editing), Dokponou Yao Christian Hugues (writing – review and editing), Ulrick Sidney Kanmounye (conceptualisation, methodology, writing – review and editing, supervision), Nourou Dine Adeniran Bankole (conceptualisation, methodology, writing – original draft, writing – review and editing, supervision).


Setthasorn Zhi Yang Ooi and Rosaline de Koning are joint first authors.


Setthasorn Zhi Yang Ooi, Rosaline de Koning, and Nourou Dine Adeniran Bankole contributed equally to this paper.

AUTHOR AFFILIATIONS

Setthasorn Zhi Yang Ooi  orcid.org/0000-0002-7097-0948
Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Rosaline de Koning  orcid.org/0000-0002-6349-2220
Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Abdullah Egiz  orcid.org/0000-0003-0304-7982
Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

David Ulrich Dalle  orcid.org/0000-0003-3676-7847
Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Moussa Denou
Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Marvin Richie Dongmo Tsopmene  orcid.org/0000-0002-5652-1986

Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Mehdi Khan  orcid.org/0000-0003-0778-4955

Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Régis Takoukam  orcid.org/0000-0002-4856-7183

Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Jay Kotecha  orcid.org/0000-0002-6463-8658

Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Dawin Sichimba  orcid.org/0000-0002-9829-2325

Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Yao Christian Hugues Dokponou  orcid.org/0000-0002-5956-2070

Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Ulrick Sidney Kanmounye  orcid.org/0000-0001-6791-1018

Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon

Nourou Dine Adeniran Bankole  orcid.org/0000-0002-4277-9767

Research Department, Association of Future African Neurosurgeons, Yaounde, Cameroon; Ibn Sina Teaching Hospital, Rabat Morocco/Mohammed V University of Rabat Morocco, Department of Neurosurgery, Rabat, Morocco

REFERENCES

- Brown T, Bota D, van Den Bent M**, et al. Management of low-grade glioma: a systematic review and meta-analysis. *Neurooncol Pract*. 2018; 6(4): 249–258. DOI: <https://doi.org/10.1093/nop/np034>
- Louis D, Perry A, Reifenberger G**, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol*. 2016; 131(6): 803–820. DOI: <https://doi.org/10.1007/s00401-016-1545-1>
- Mandonnet E, Delattre J, Tanguy M**, et al. Continuous growth of mean tumor diameter in a subset of grade II gliomas. *Ann Neurol*. 2003; 53(4): 524–528. DOI: <https://doi.org/10.1002/ana.10528>
- Claus E, Walsh K, Wiencke J**, et al. Survival and low-grade glioma: the emergence of genetic information. *Neurosurg Focus*. 2015; 38(1): E6. DOI: <https://doi.org/10.3171/2014.10.FOCUS12367>
- Bondy M, Scheurer M, Malmer B**, et al. Brain tumor epidemiology: Consensus from the Brain Tumor Epidemiology Consortium. *Cancer*. 2008; 113(S7): 1953–1968. DOI: <https://doi.org/10.1002/cncr.23741>
- GBD Compare|IHME Viz Hub**. *Vizhub.healthdata.org*. <http://vizhub.healthdata.org/gbd-compare>. Accessed December 7, 2021.
- Ostrom Q, Gittleman H, Liao P**, et al. CBTRUS Statistical Report: Primary brain and other central nervous system tumors diagnosed in the United States in 2010–2014. *Neuro Oncol*. 2017; 19(suppl 5): v1–v88. DOI: <https://doi.org/10.1093/neuonc/nox158>
- Nguemu S, Kanmounye U, Tétinou F, Djiofack D, Takoukam R**. Barriers to Management of Nontraumatic Neurosurgical Diseases at 2 Cameroonian Neurosurgical Centers: Cross-Sectional Study. *World Neurosurg*. 2020; 139: e774–e783. DOI: <https://doi.org/10.1016/j.wneu.2020.04.162>
- Benyaich Z, Hajhouji F, Laghmari M**, et al. Awake Craniotomy with Functional Mapping for Glioma Resection in a Limited-Resource-Setting: Preliminary Experience from a Lower-Middle Income Country. *World Neurosurg*. 2020; 139: 200–207. DOI: <https://doi.org/10.1016/j.wneu.2020.04.039>
- Ndubuisi C, Mezue W, Nzegwu M, Okwunodulu O, Ejembi G, Ohaegbulam S**. The Challenges of Management of High-grade Gliomas in Nigeria. *J Neurosci Rural Pract*. 2017; 8(3): 407–411. DOI: https://doi.org/10.4103/jnrp.jnrp_18_17
- Weiss H, Garcia R, Omiye J**, et al. A Systematic Review of Neurosurgical Care in Low-Income Countries. *World Neurosurg X*. 2020; 5: 100068. DOI: <https://doi.org/10.1016/j.wnsx.2019.100068>
- Arksey H, O'Malley L**. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol*. 2005; 8(1): 19–32. DOI: <https://doi.org/10.1080/1364557032000119616>
- Tricco A, Lillie E, Zarin W**, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med*. 2018; 169(7): 467–473. DOI: <https://doi.org/10.7326/M18-0850>
- Moher D, Shamseer L, Clarke M**, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev*. 2015; 4(1). DOI: <https://doi.org/10.1186/2046-4053-4-1>
- Koning R, Adeniran B, Ooi S**, et al. Inventorying the Management and Outcomes of Low Grade Gliomas in Africa and Suggestions for Improvement: A Scoping Review Protocol. *OSF*. DOI: <https://doi.org/10.17605/OSF.IO/E732G>. Published 2021. Accessed December 7, 2021.
- Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A**. Rayyan—a web and mobile app for systematic reviews. *Syst Rev*. 2016; 5(1). DOI: <https://doi.org/10.1186/s13643-016-0384-4>

TO CITE THIS ARTICLE:

Ooi SZY, de Koning R, Egiz A, Dalle DU, Denou M, Tsopmene MRD, Khan M, Takoukam R, Kotecha J, Sichimba D, Dokponou YCH, Kanmounye US, Bankole NDA. 2022. Management and Outcomes of Low-Grade Gliomas in Africa: A Scoping Review Protocol. *International Journal of Surgery: Protocols*, 26(1), pp. 1–6. DOI: <https://doi.org/10.29337/ijsp.171>

Submitted: 20 December 2021 Accepted: 25 January 2022 Published: 02 February 2022

COPYRIGHT:

© 2022 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

International Journal of Surgery: Protocols is a peer-reviewed open access journal published by IJS Publishing Group.