

RESEARCH

Open Access



# Reducing drug-use harms among higher education students: MyUSE contextual-behaviour change digital intervention development using the Behaviour Change Wheel

Vasilis S. Vasiliou<sup>1\*</sup> , Samantha Dockray<sup>1</sup>, Samantha Dick<sup>2</sup>, Martin P. Davoren<sup>2,3</sup>, Ciara Heavin<sup>4</sup>, Conor Linehan<sup>1</sup> and Michael Byrne<sup>5</sup>

## Abstract

**Background:** Digital harm-reduction interventions typically focus on people with severe drug-use problems, yet these interventions have moderate effectiveness on drug-users with lower levels of risk of harm. The difference in effectiveness may be explained by differences in behavioural patterns between the two groupings. Harnessing behavioural theories to understand what is at the core of drug-use behaviours and mapping the content of new interventions, may improve upon the effectiveness of interventions for lower-risk drug-users. To the best of our knowledge, this is the first study to systematically apply the Behaviour Change Wheel (BCW) approach to understand the components, influencing capabilities, opportunities, and motivations (COM-B) of higher education students to change their drug-use behaviors. It is also the first study which identifies specific patterns of behaviours that are more responsive to harm reduction practices through the use of the Theoretical Domain Framework (TDF).

**Methods:** We employed an explanatory sequential mix-method design. We first conducted an on-line survey and a Delphi exercise to understand the factors influencing COM-B components of higher education students to change their drug-use. Subsequently, we mapped all evidence onto the COM-B components and the TDF domains to identify clusters of behaviours to target for change, using a pattern-based discourse analysis. Finally, a series of multidisciplinary group meetings identified the intervention functions—the means by which the intervention change targeted behaviours and the Behavioural Change Techniques (BCTs) involved using the behaviour change technique taxonomy (v.1).

**Results:** Twenty-nine BCTs relevant to harm-reduction practices were identified and mapped across five intervention functions (education, modelling, persuasion, incentivization, and training) and five policy categories (communication/marketing, guidelines, regulation, service provision, and environmental/social planning). These BCTs were distributed across eight identified saturated clusters of behaviours MyUSE intervention attempts to change.

\*Correspondence: v.vasiliou@ucc.ie

<sup>1</sup> School of Applied Psychology, University College Cork, N Mall, Kilbarry

Enterprise Centre, Cork Enterprise Centre, Cork, Ireland

Full list of author information is available at the end of the article



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Conclusions:** The BCTs, identified, will inform the development of a digitally delivered behaviour change intervention that focuses on increasing mindful decision-making with respect to drug-use and promotes alternatives to drug-use activities. The findings can also inform implementation scientists in applying context-specific harm-reduction practices in higher education. We present examples of how the eight identified clusters of target behaviours are mapped across the COM-B components and the TDF, along with suggestions of implementation practices for harm reduction at student population level.

**Keywords:** Higher education students, Illegal drug-use, Contextual-behaviour change intervention, Psychological Flexibility, Digital intervention

## Contribution to the literature

- Proposes a conceptual map of the influences of drug-use behaviours in higher education settings
- Suggests an implementation paradigm shift on how to address harms from drug-use among higher education students
- Indicates foci for harm reduction implementation practices in higher education settings

## Background

Illicit drug-use is becoming a public health concern among young adults [1]. The prevalence of drug-use reaches its peak among 18–25 years olds [2–4], with cannabis being the most common illicit drug used by more than 30% of higher education students in the US [5], followed by non-prescribed use of prescription medications, including drugs such as amphetamines [4]. Drug use among young adults coincides with the time when many individuals enter higher education and when experiencing neurodevelopmental processes in the prefrontal cortex that affect risk taking behaviours [6]. Drug use occurs during a period of transition for young adults who are gaining independence, with many living away from family for the first time [7]. In this transitional phase, experimentation with or regular use of drugs is seen as a normative behaviour by students, to either achieve some form of personal enhancements [8], develop new social support structures [2], or enhance a new experience [9, 10]. A period of drug use experimentation can sometimes drive individuals to form new drug-use habits, some of which can continue far into adulthood, leading to possible negative consequences in adult life [11–13].

The potential risks and adverse consequences of drug-use to student populations map across several life domains, including lower grade point averages, poor class attendance [14], heavy drinking [15], polydrug-use [16], and other high-risk behaviours, such as driving under the influence, unprotected sex, physical fights [17], or exacerbation of mental health problems [18]. These

consequences may be experienced concurrently, potentially reshaping trajectories of wellbeing across the life course [14, 19, 20]. Given the potential harms illicit drugs can cause, preventive and intervention programmes at higher education institutions are needed to respond effectively to drugs, used by students.

Harm reduction interventions to reduce injection-related harms among young population exhibit moderate-to-large effect for individuals with severe drug use problems [21]. For the population with less severe drug use problems, traditional harm-reduction interventions do not show such a strong evidence of effectiveness [21–24], including when these interventions incorporate personalized information feedback provision and corrections of norms; two components considered to be associated with reductions in harms in other areas, such as alcohol use. Even when these intervention are delivered, digitally, at low-cost, with high levels of acceptability among young adults [25, 26], they achieve only modest success in harm reduction [27, 28]. The reasons for this are the lack of a unified theory-driven behaviour change framework that informs the design and development of the behaviour change intervention. Also, these interventions lack theory-based contextual-driven approaches that incorporate skills training as vehicles to implement harm-reduction practices [21, 27, 28]. In order to effectively employ behaviour change practices, we first need to describe and understand both what is at the core of the target behaviour to change [29] and the contextual influences on this behaviour [30, 31]. We must also make use of process-based behavioural change practices [32–34] which can predict and influence behaviours (e.g. ongoing decisions and actions) with scope, precision, and sensitivity to the context where these behaviours occur [35].

For drug-use, there have been calls for research that goes beyond understanding the antecedents and consequences of use (e.g., motivation for use, perceived harms/benefits of drug use) and towards the explicit use of theories to understand what is at the core of drug-use behaviours [36] within a person's context [35, 37, 38]. The psychological theories in motivation for change and transtheoretical models of change [39]

have produced a plethora of perspectives that explain the phenomenon of drug-use. However, a synthesis of this knowledge to systematically map drug-use behaviours in the users' context, thus, inform current interventions development and implementation practices, has yet to be developed. Given that harm reduction theory inherently acknowledges that some drug use is likely within different populations [40], innovations in harm-reduction approaches at student population level should target to reduce drug use and minimize the harms occurring through the use of drugs [27, 28]. One framework that can provide a systematic way to use behavioural theories in developing such an intervention is the Behaviour Change Wheel (BCW) [29].

At the core of the BCW approach is the COM-B model [29, 41] which suggests that individuals need the capability (C), opportunity (O), and motivation (M) to change behaviours. In addition, there are nine intervention functions (e.g. modelling) via which an intervention exerts its effect and seven policy categories (e.g. regulation) that support the implementation of the intervention [29]. Also, the COM-B approach includes the behaviour change taxonomy (BCCTv1) [42] which allows the identification of the "active ingredients" of the intervention through a list of 93 possible theory-driven behaviour change techniques (BCTs). Coupled with the COM-B model is the Theoretical Domain Framework (TDF) [31]. This framework consists of 14 theoretical determinants (e.g. knowledge, skills, beliefs about consequences, etc.) which can enhance understanding of the cognitive, emotional, social, and environmental influences on target behaviours [31]. In turn, this knowledge can be translated into agile and effective behaviour change components [36, 43] that can reduce the harm of drug-use at a population-level [44]. In drug-use, the BCW approach and the TDF framework have guided the development and implementation of interventions that target drug-use related harms [45]. But have not yet been applied in the context of higher education populations.

To address this gap, we established the MyUSE (My Understanding of Drug Use Experiences) [46] project that aims to develop a theoretically based, digitally delivered behaviour change intervention to reduce harms from drug use in higher education students. MyUSE project adopted the BCW [29] approach that allows intervention researchers to understand the target behaviours via a behavioural analysis of the problem and then design an intervention on the basis of this analysis.

In this article, for the first time in the relevant literature, we use the BCW approach and the TDF framework, as guides to identify the BCTs that informed the MyUSE content. The specific aims of this study were to

synthesize the evidence from primary (survey) and secondary (three systematic reviews) data sources, gathered as part of the overall project, called MiUSE (My Understanding of Drug Use Experiences) [46] in order to (a) identify clusters of drug-use target behaviours among higher education students; (b) select specific intervention functions through which the intervention will exert its effect, and (c) choose specific BCTs to be operationalized within the MyUSE digital intervention.

## Methods

In January 2020, we established a multidisciplinary advisory team with experts in Behavioural Science (VSV, SD2, CL), Information Systems (CH), Public Health (SD1, MD), and Student Health Services (BM) to assist the completion of the BCW approach. The advisory team completed four open-sort grouping exercises, a Delphi-type exercise with two rounds, and a discourse pattern-based analysis, over eight-months to complete the BCW analysis of drug use behaviours. Prior to beginning the analyses, all members attended a training session on the BCW approach and consented to participate in a series of consensus-type meetings. Ethical approval was granted from the Social Research Ethics Committee and University College Cork (UCC) (SREC reference number no: 2018-072A). Written informed consent was obtained from all participants in this study. TIDieR checklist was used for developing this manuscript.

### Phase I—Identifying the problem in behavioural terms

As presented in the BCW approach [41], phase I consisted of four steps. In steps one to three, we analysed drug-use behaviours among higher education students in behavioural terms. In step four, we sought to identify what specific aspects need to change for targeted behaviours to occur.

#### Step 1—Define the problem in behavioural terms

The research group had previously conducted three systematic reviews [27, 28, 48] to identify, gather, and understand all relevant research in relation to the MyUSE project. The first systematic review examined the effectiveness of digital behavioural change interventions for drug use harm reduction in student populations, showing only modest success of these interventions in reducing the harm of drug use [27]. The second review was conducted to examine whether previous similar interventions had employed user-centered design (UCD) practices to inform the intervention development. Findings showed only limited consideration had been given to the end user experience (UX) in designing interventions through UCD practices; limiting their potential effectiveness and sustainability [28]. Finally, the third review

examined motivational factors for students' decision to lower or cease drug use, showing that the identification of the adverse consequences of drug use is not sufficient to prompt behavioural changes among this cohort [48].

Specific findings from these three systematic reviews were used to define the problem in behavioural term and later (step 2) to help us identify items as relevant to targeted behaviours to change. From the first systematic review, an important finding that guided our decision making was that we identified eight studies; of those four focused on cannabis use changes, three on multiple changes in health behaviours and one on changes in different drug use; making it difficult to isolate the mechanism of change. In the second review, we identified personalization and feedback provision as components that can drive behavioural changes. Finally, the third review indicated social factors as strong predictors of drug-use related behaviour, including students' concerns on how their peers would view them and the feeling of shame surrounding drug use. Both were regarded as important components in defining the targeted behaviour to change.

The advisory group participated in the first open sort grouping exercise, setting out the findings from the three systematic reviews in a template (whiteboard in a class). They worked in pairs and asked to complete a worksheet, focusing on two questions: (a) what is the target group of individuals involved in the behaviour? and (b) where does the behaviour occur? A consensus meeting followed and agreement on how to define the problem in behavioural term was reached.

### **Step 2—Select the target behaviour**

The first author (VSV) worked through the findings from the three systematic reviews [27, 28, 48] and the relevant literature to create a long list of items deemed relevant to the targeted behaviour(s). The group then participated in a Delphi-type exercise with two rounds. In each round, the members rated each behaviour using the APEASE criteria (Affordability, Practicability, Effectiveness, Acceptability, Side effects, Equity; see Table 1 for definitions of the APEASE criteria) [47]. The members of the expert advisory group rated the long-list of the targeted behaviours ( $n=57$ ), and these ratings were used to collate a shorter-list for the second round of ratings. Inter-rater agreements were calculated, using a 70% threshold as an agreement point for the first round (long-list rating; i.e. participants fall within two agreement categories on a Likert-type scale from 0 = not at all impactful to 5 = extremely impactful [49]), and a threshold of 3.25 median score in the second round (short-list) to resolve differences (convergence biases

of opinion) [47]. The final selection of the target behaviour was agreed upon by the expert group members in a consensus meeting.

### **Step 3—Specify the target behaviour**

Following the synthesis of the reviews [7–9], in Step 3 the expert advisory group participated in a third open sort grouping exercise to specify: the target behaviours, the population, and the context (when and where the behaviours will be performed).

### **Step 4—Identify what needs to change**

In step 4, we triangulated data to better understand what needs to happen for the target behaviour change to occur [41]. We utilized an explanatory sequential mixed-method design, as defined by Creswell [50]. In this process, we analysed and mapped the data from a quantitative on-line survey of student drug use, onto the COM-B model and TDF framework. The findings from this quantitative analysis informed the qualitative synthesis that followed (the pattern-based discourse analysis [51, 52]). We used the comprehensive set of theoretical construct domains (TDF) to select the domains most relevant to the targeting behaviours and then mapped those domains into the COM-B summative components.

In the first phase, we developed a survey to measure individual and contextual factors relevant to drug-use behaviours (see the Additional file 1; MyUSE drug use survey overview). A Public and Patient Involvement (PPI) group consisting of higher education students reviewed the survey for cultural context appropriateness. The clarity, acceptability, and relevance to students' context were piloted by a small group of students ( $n=6$ ) that made several modifications to items and phrasing in the survey. The final survey comprised of six sections (demographics, student life, patterns of drug-use, decision-making process, motivations for using, and behaviour change). The survey was then distributed electronically to a randomly selected representative sample of UCC students ( $n=3770$ ) via emails. The survey achieved a 30% response rate ( $n=1138$  responses) and a 66% completion rate. Following data clearance, descriptive analysis and reporting (SPSS, V22), a mapping exercise was used, to synthesize qualitatively, aspects of the targeted behaviour to change.

In the second phase, we used the on-line survey data through a large mapping exercise, following a pattern-based discourse analysis [51, 52]. We followed the Strauss and Corbin [53] procedure through a systematic analysis of the on-line survey data, generating categories of intervening (situational factors [54]) that were then translated into narrative statements [53]. Next, pattern-based

**Table 1** Behavioural specification of the target behaviours

Target behaviour	Who	What	When	Where
Increase students' behavioural awareness regarding their decision to take drugs	Higher education students (aged 18–25 yrs)	1. Pause automatic/habitual decision making in relation to drug-use	1. In night outs/ social media/ festive periods	Social contexts involving a decision to take drugs (e.g. at campus, in social media, streets around the university, etc.)
Reflect on alternatives to drug-use activities which fulfil personal enhancements	Full-time/ part time students, Erasmus/visiting students (bachelor to graduate studies)	2. Increase behavioural awareness in relation to drug-use decision making 3. Committing in specified goals which can fulfil personal enhancements	2. During induction weeks 3. Before, during & after the exam's periods 4. Randomly repeated during the semester	
Selected items derived from the Delphi-type exercise using the APEASE criteria				
[A1]	Users to compare their own use with the norm (descriptive norm correction; eg. % of students with higher use than their own)			
[A2]	Users to reflect on their close friends' perception about their own use or abstinence (injunctive norm correction)			
[A5]	Users to find out personally relevant primary motives for use, listing the personal enhancements drug use fulfil to them (underlying needs that drive the decision to take drugs)			
[A10]	Users to reflect on the possible negative consequences of using drugs on academic, athletic and social performance			

Note 1: Given that the four selected items (see Additional file 1: Table 2) are all clustered within the behavioural domain A, specifications are applied for both targeting behaviours

Note 2: The advisory group consulted the APEASE criteria to guide their decisions making. The APEASE criteria specify as to: (a) the impact of the selected behaviour on the intervention's desired outcomes; (b) the likelihood of behavioural change (c) the spillover effect of behaviour on related behaviours; (d) the accessibility of the targeted behaviour; and (e) the efficacy of the behaviour in bringing about the desired outcomes



discourse analysis was conducted in four steps. All derived statements were “grounded” and shaped by university students with previous drug use experiences (theoretical sampling; [54]).

In the first step (open coding), two members of the expert advisory group (coders; VSV, SD2) identified items from the on-line survey and mapped them in one of the three COM-B components, based on definitions of the components, provided by Michie et al. [41]. The items were analysed quantitatively (descriptive statistics), and then, qualitative descriptors were added to explain an aspect of the targeted behaviours through the COM-B lens [55]. During this process, the coders analysed the selected on-line survey items and extracted patterns (descriptors). In the second step (axial coding), the sequential mixing of data [56] occurred. Here, the coders considered the three COM-B components as the phenomena to focus, and the TDF as the main categories, assembled around the core phenomena [53]. The coders coded the data deductively, assembling the qualitative descriptors, generated previously, in the COM-B components, and then systematically relating these descriptors into the TDF categories, based on the definitions of the domains provided by Atkins et al. [31]. In the third step (selective), the coders selected codes that had been previously qualitatively described (descriptors), to develop narrative statements describing what needs to change for the targeting behaviours MyUSE intervention attempts to change [53]. In the final step, according to Strauss and Corbin [53], the advisory group participated in a meeting, to develop a template with narrative statements (clusters of behaviours to change) that elucidate the potential influences the COM-B and the TDF domains may have in the MyUSE intervention.

Credibility of the analysis was ensured through multiple mechanisms. The first author, a qualified Clinical Psychologist with experience working with young adults and drug addiction, immersed himself in the whole research process, engaging in an ongoing familiarization with the data and discussions with co-researchers and the Student Advisory Group (SAG) [57]. The second co-author, a behavioural scientist with experience in developmental psychology, examined the patterns for transparency issues [58] and reached agreement on the mapped items across the four steps of the analysis. Thirdly, coders included a detailed log with the coding data [51]. Finally, there were several opportunities for the multidisciplinary advisory team, involved in the BCW approach, to debate and reach consensus in relation to the iterative emergence of the findings.

For each one of the analytic steps of the discourse analysis (open, axial, selective, and paradigm coding), the

coders kept thorough memos, writing down ideas and coders’ personal impressions about the emerging narratives, acknowledging the role of reflexivity [57]. In relation to reflexivity, two members of the team (VSV and SD2) led the data interpretation and synthesis part of this analysis. Finally, in keeping with Fine’s recommendations [59] that researchers move beyond the data, we sought to validate the narratives by conducting another qualitative inquiry engaging non-users and users in a card-sorting exercise. Here, we examined the narratives in terms of their importance by presenting a series of cards and asking participants to engage in an open dialogue about their importance in relation to targeted harm-reduction and prevention practices [60].

This mapping exercise was undertaken to examine: (a) what function each TDF domain and COM-B components serve in the target behaviours and (b) what needs to change for the target behaviour change to occur.

#### **Phase II: Identifying intervention options**

Phase II consisted of two steps to guide our decision on intervention functions and policy categories.

##### ***Step 5 & 6—Identify intervention function and policy categories***

Using the APEASE criteria, three co-authors (VSV, SD1, MD) examined, first individually, and then as a group, the nine intervention functions. The goal was to examine whether the functions would serve each of the identified clusters of target behaviours. Further, they also identified policies that support the intervention functions. In each selection of functions, Fleiss’ kappa statistics [61] were calculated, to quantify the reliability of agreement between the raters.


#### **Phase III: Identify content and implementation options**

##### ***Step 7—Identify behaviour change techniques***

To identify specific BCTs, the first author employed the taxonomy matrix of 93 BCTs (BCTTv1) [42] to generate an extended list of BCTs that could be mapped on the eight identified clusters of techniques relevant to drug-use behaviours. Three members of the advisory group with expertise in behavioural science used the APEASE criteria to exclude non-relevant BCTs with the targeting behaviours to change. A shorter list of BCTs was generated, and a final selection of the targeted BCTs was consensually agreed.

##### ***Step 8—Identify modes of delivery***

The mode was predetermined as digital, so this step was not executed.



## Results

The results reported represent the application of the BCW approach. A summary of the BCW approach employed is illustrated in Fig. 1.

### Step 1—Define the problem in behavioural terms


Following a consensus meeting, the problem was defined as (a) the low level of awareness regarding drug-use decision making, (b) poor insight as to the consequences of drug-use behaviours, and (c) the lack of opportunities to consider alternatives to fulfil personal enhancement. Therefore, the group decided that the intervention should focus on (a) increasing mindful decision-making in relation to illicit drug-use behaviours, and (b) enhancing individuals’ insight for alternatives to drug-use behaviours as a mean to fulfil the students’ personal enhancement. We also decided that this intervention should be most effective for students declaring occasional drug use or no previous drug use. Finally, harm-reduction practices were deemed more useful in social events (e.g. parties, gatherings, etc.) and in places where alcohol is consumed (e.g. pubs, bars, night clubs).

Some of the identified influences on drug-use are related to the physical and social opportunities that may be afforded by the university context. Other factors are related to students’ reflective motivation (e.g.

how their peers would view them, feelings of shame) and automatic motivation (e.g. a desired outcome from the use). Finally, capabilities are not identified in the context of students’ drug-use behaviours. The scoping review [48] identified two contextual variables as potentially risky factors: the university context and the transition period from the second level (i.e. high school) to higher education. These factors require harm-reduction strategies at a systemic/policy-level (e.g. new public health responses to illicit drugs and alcohol use) [62]. However, our analyses of behavioural diagnostics indicated the value of individual-level focus.

### Step 2—Select the target behaviour

A long list of 67 potential behaviours (items) was derived from the synthesis of the relevant literature. Figure 2 illustrates five potential targets relevant to drug-use behaviours that can increase students’ awareness in relation to their decision to take drugs and can increase understanding of alternatives as means to fulfil the students’ personally relevant enhancements (the entire list is shown in Additional file 1: Table A1). In refining the long list of potential targets, one can see that current university service provisions address some of the targeting behaviours. For example, psychoeducation about the consequences of illicit drug-use (domain C) is one service



universities often provide as part of their health care policies [62]. The advisory group participated in a Delphi-type exercise with two ranks.

In round one from the 67 items identified, only 24 items reached the agreement point of 70% (see Additional file 1: Table A1). Seventeen items were from the targeted behaviour A (increase awareness of current illicit drug-use behaviours), four from B (maximize self-regulatory capacity and skill), one from C (resilient-related skills), and two from E (address personalized-contextual factors). None of the items from domain D (provide psychoeducation relevant to drug-use) reached the agreement point, and this behavioural target was excluded from round two.

In round two, from the 24 items selected, 15 items reached the agreement point of the median score, using the APEASE criteria (values > 3.25), as potential drug-use behaviours to target (10 from A; 4 for B; and one from E; see Additional file 1: Table A2). Finally, of these 15 items, only 4 reached the agreement of the mean score of the APEASE criteria (ranked > 70%) and these were selected as potential behaviours to target. All four items (targets relevant to drug-use behaviours) were from the behavioural target A (see Table 1 for the four identified behaviours). These items indicate that the targeted behaviours

should increase awareness in relation to contextual factors (e.g. peers) that influence drug-use decision making and enhance insight as to the internal motivations of the students to use drugs (e.g. expectations).

**Step 3—Specify the target behaviour**

Table 1 presents the specifications of the targeted behaviours and the four selected items derived from the Delphi-type exercise.

**Step 4—Identify what needs to change**

We present an overview of the findings from the on-line survey. We then present the findings from mapping the identified patterns of drug-use behaviours onto the COM-B components. We finally present the findings from the discourse-based analysis that show what needs to change for the targeted behaviours to occur, using the TDF domains. Table 2 below shows the patterns of drug-use related behaviours mapped on the COM-B components and the TDF domains. Table 3 summarizes the findings of the whole analyses arising from the step 4.

**Findings from the on-line survey**

Almost a third (32%) of respondents reported using an illicit drug in the last year (n = 236; current users) with







**Table 3** Intervention components targeting the eight identified patterns of drug-use related behaviours

Pattern of drug-use related target behaviours	COM-B mapped elements targeted	Intervention Function served	Coded identified (from BCTTv1)	BCTs selected	Translation of BCTs within MyUSE intervention
Cluster 1: Increase awareness about the real vs. anticipated effects drug-use can have on students' personally desired behaviours	Reflective motivation	Education	5.1	Information about health consequences	Provide a personalized animated feedback explaining the possible health-related risks and harms per drug-use type and frequency
			5.3	Information about social and environmental consequences	Present a gamified quiz showing the possible consequences of drugs in students' academic and emotional area of living (e.g. legal problems, etc.)
			5.6	Information about emotional consequences	
			6.3	Information about others' approval	Provide a subjunctive norm correction, in a gamified quiz way
			13.4	Valued self-identity	Include a story-telling exercise requesting from students to produce self-statements of their life they want while in college
			1.3	Goal setting (outcome)	Present an interactive value's identification and committed action exercise
Cluster 2: Promote identification of personally relevant activities (which are they?) which lead to positive expected outcomes in students' desired behaviours (fun & enjoyment)		Education	7.2	Cue signalling reward	Present a reflective- non-judgmental rhetorical question, prompting students to consider how their values match with drug-use behaviours
			13.2	Framing/reframing	Suggest the adoption of alternative to drug-use activities as means to enhance fun and enjoyment
			5.6	Information about emotional consequences	Present a mindful-based exercise ("how fully present am I?"), prompting students to visualize (mental representation) how they would feel after achieving college-related goals
			15.1	Verbal persuasion about capability	Use motivational cards to show how students can pursue value committed actions
			16.3	Vicarious learning	Present a scenario-based story showing a student pursuing his/her goals while enjoying college's years
			1.4	Action planning	Prompt students to use their e-calendar to plan implementing SMART defined goals

**Table 3** (continued)

Pattern of drug-use related target behaviours	COM-B mapped elements targeted	Intervention Function served	Coded identified (from BCTTv1)	BCTs selected	Translation of BCTs within MyUSE intervention	
Cluster 3: Increase perceived competence & optimism that an implementation plan of alternatives to drug-use activities can induce positive expected experiences (fun & enjoyment)		Education	1.3	Goal setting (outcome)	Use an animation to educate how commitment to SMART goals can work as alternative to drug-use behaviours and reinforce the deliberate adoption of several, non-drug-use behaviours, asserting that these behaviours can elicit as fun as drugs can, without putting students into risks or harms	
			1.9	Commitment		
			6.3	Information about others' approval		
			8.2	Behaviour substitutions		
			8.6	Generalization of a target behaviour		
			13.2	Framing/reframing		
			15.1	Verbal persuasion about capability		
		Education	2.3	Self-monitoring of behaviour	Explain the role of cues (triggers of use) within an ABC analysis (behavioural analysis), prompting students to identify (self-monitoring) their own antecedent triggers in relation to the context of use	
	Cluster 4: Increase awareness of the university context as a risk factor that increase the chances for drug-use behaviours and how this can restrict students from having positive college's experiences			4.2	Information about antecedents	
			Modelling	7.1	Prompt/cues	Show how self-talk can be used to help students recognize cues that can influence decision making in relation to drug-use
			15.4	Self-talk		

**Table 3** (continued)

Pattern of drug-use related target behaviours	COM-B mapped elements targeted	Intervention Function served	Coded identified (from BCTTv1)	BCTs selected	Translation of BCTs within MyUSE intervention
Cluster 5: Cultivate mindful awareness of the perceived reasons for using (why I use?) and increase insight as to whether the use leads to desired outcomes in goal-directed behaviours at a long-run		Education	1.6	Discrepancy between current behaviours and goals	Present a personalized feedback showing students' level of behavioural awareness and goal attainment (i.e. whether their decision to take drugs is influenced by others). Then, prompt students to monitor whether there is a discrepancy between goals and present behaviours
			2.4	Self-monitoring of the outcomes of behaviours	
			4.1	Instructions on how to perform a behaviour	Introduce via an animation the mindful decision-making skill (learn to pause step-back, notice and decide) and prompt students to think how this skill resonates with their role identity
			8.1	Behavioural practice/rehearsal	
			13.4	Valued self-identity	
			4.2	Information about antecedents	Prompt students to apply the new skills to identify cues
		Modelling	4.1	Instructions on how to perform a behaviour	Present via a story-narrative (a party house) how a student applying mindful decision-making skill (pausing- noticing- deciding) in a situation requiring drug-use decision making.@Heinforce students' capacity to apply the new skill in different situations
			6.1	Demonstration of the behaviour (modelling)	
			8.1	Behavioural practice/rehearsal	
			8.6	Generalization of a target behaviour	
			15.1	Verbal persuasion about capability	
		Training	6.1	Demonstration of the behaviour	Present a series of mindful cards demonstrating what mindful decision-making skill does and advice for a gradual implementation of this skill building activity
			8.7	Graded tasks	



**Table 3** (continued)

Pattern of drug-use related target behaviours	COM-B mapped elements targeted	Intervention Function served	Coded identified (from BCTTv1)	BCTs selected	Translation of BCTs within MyUSE intervention
Cluster 6: Resolve students' misleading expectations about the expected outcomes of drug-use in students' desired behaviours in the long-run	Automatic motivation	Education	5.2	Saliency of consequences	Provide a general performance score in a drug-use quiz activity. Wrong-answered responses will provide feedback that will target at correcting students' expectations about the effect of drugs in the long-run. They will also aim at increasing awareness about the potential regrets, students may experience from the use of drugs
			13.2	Framing/reframing	
Cluster 7: Increase procedural knowledge and practice skills on how harm-reduction practices are implemented within the university context	Capability physical	Persuasion	5.5	Anticipated regret	Present a series of harm-reduction practices and invite students to select the ones that best fit with their experiences (personalized plan)
			1.8	Behavioural contract	
		Education	4.1	Instructions on how to perform a behaviour	Show instructions in detail on how to perform selected harm-reduction practices via a series of illustrative cards
			8.6	Generalization of the target behaviour	Prompts students to generalize the new harm-reduction practices, including awareness of exposure to cues, to multiple different situation
		Modelling	12.3	Avoidance/ reducing exposure to cues for the behaviour	Show an animation illustrating a student to perform one harm reduction practice, highlighting his/her commitment
			1.9	Commitment	
			4.1	Instruction on how to perform a behaviour	
		Training	4.1	Instruction on how to perform a behaviour	
			6.1	Demonstration of the behaviour	

**Table 3** (continued)

Pattern of drug-use related target behaviours	COM-B mapped elements targeted	Intervention Function served	Coded identified (from BCTTv1)	BCTs selected	Translation of BCTs within MyUSE intervention	
Cluster 8: Promote behavioural awareness and behavioural regulation regarding drug-use decision making under the influence of peers	Social Opportunity	Education	1.6	Discrepancy between current behaviour and goal	Prompt students to reflect on their personalized feedback scores in questionnaires assessing levels of decision making influenced by others	
			4.1	Instruction on how to perform a behaviour	Present instructions on how to promote behavioural awareness in relation to decision making	
			4.2	Information about antecedents	Prompt students to think peers' influences as antecedent (cues) for them to use drugs	
			5.2	Saliency of consequences		
			8.2	Behavioural substitution	Promote a mindful decision-making, highlighting that influences from peers should be aware	
			8.7	Graded tasks	Provide suggestions on building the new behavioural regulation skill (behavioural awareness)	
			13.2	Framing/reframing	Prompt students to consider their feedback on subjective norm correction (how behaviours are approved by others) in relation to their valued identity	
			15.2	Mental rehearsal of successful performance	Advise students to imagine themselves taking a mindful decision, despite the presence of peers' influence	
			4.1	Modelling	Instruction on how to perform a behaviour	Illustrate an animation (a narrative story) with a student denying using drugs, while recognizing the potential influence of peers
			6.1	Demonstration of the behaviour		

Note 1: Physical opportunity was not targeted

Note 2: Translation of BCTs into digitally delivered components represents only concepts which are designed into ideations prototypes to be tested with students, rather than actual implementation practices for the MyUSE digital intervention

(86%), energy levels (62%), and decreases in irritability (70%), and distress (68%), with these effects reverting when the effects of the drugs wear off. For those declaring previous use, the five main motivations for change were: noticeable psychological impacts, financial burden, physical effects, impairments in executive functions, and concerns about how other people perceive their drug-use. Students also reported willingness to change their use if they were to socialize with other groups (18.7%), had alternatives to drug-use activities (24%) or had better ways to manage unwanted emotions (10.5%). From those declaring current use, 41.3% reported no confidence to use harm-reduction measures as means of protecting themselves from the effects of drugs, with 36.6% reporting “somewhat confidence” and 22.1% “confidence”.

#### **Findings from the pattern-based grounded discourse analysis**

During the first step of the coding (open coding), we identified, coded, and mapped onto the COM-B a total of 25 patterns of drug-use behaviours derived from the survey (see Table 2). 23 items were coded into motivation (17 reflective and 6 automatic), 5 in capability (3 in physical and 2 in psychological), and 7 in opportunity (5 in social and 2 in physical). Most of the selected items were first analysed quantitatively and then described, qualitatively (see Additional file 1: Table A3; step 1). They all focus on the role of reflective motivation. Data indicate how students’ behaviours are habitually driven by their motivation to take drugs in order to achieve a desired end-goal which is always relevant to college’s life (e.g. have fun, etc.). This decision occurs automatically and is partly grounded in the absence of students’ awareness of other means that lead to similar outcomes in desired behaviours, but with less risk of harms for them (e.g., valued-drive activated). Though descriptors provide salient categories -indicating students’ increased awareness about the risks and harms of drugs, what prevails is a habitual and automatic decision making on behalf of students, particularly when drugs are involved in their decision. Further, analysis showed that social opportunities increase decision making, favouring drug use. Finally, although the analysis showed sufficient psychological capability (i.e. knowledge, understanding) from students to reduce the use or harms from drugs, poor knowledge about how harm-reduction practices can be applied, limit capabilities towards the targeting behaviour.

In the second step of the analysis (axial coding), as presented in Table 2, we fit the descriptors of the initial coded data into the central phenomena (COM-B components) and the main categories (TDF). We synthesized the coding data into selective qualitative descriptors (see Additional file 1: Table A3; step 2). In the third step (selective coding), we developed narrative statements,

based on selective codes, exploring the role of each TDF domain (descriptor) in influencing the three behaviours MyUSE attempts to change (see Additional file 1: Table A3; step 3). In our analysis (see below), we examined what factors support the three targeting behaviours within each TDF domain, and also what competes or inhibits the desired behaviours to change.

#### **Reflective motivation**

**Beliefs about consequences:** students have strong expectations of the role of drug-use in enhancing personally relevant areas of interest. The immediate and potent effects of drugs (e.g. increased energy level, social interactions, confidence, reduction in anxiety, irritability, etc.) enhance students’ beliefs as to the long-term effects of drugs in fulfilling areas that matter the most to them (e.g. having fun with their friends). Although students report having concerns about the negative effects of drug use in the long-term (i.e. more than 80% agreed on that), these beliefs are buffered by the strong and immediate positive effects drug can have in students’ relevant areas of interest. Further, the immediate effects from drug use are directly and indirectly reinforced by contextual variables (peers, perceived expectation for use in higher education, fun, etc.), disinhibiting the potential effect of any negative beliefs students have in the long-run. This leads students to maintain misleading expectations as for to the effects of drug use in the long-term. Therefore, increasing students’ awareness of the perceived long-term consequences versus the perceived short-term benefits may lead to an increase in students’ harm reduction practices and possibly reduces levels of drug-use.

**Intention:** the current users’ intention to abstain from drug-use, in contrast to the non-users, was found to be low. Students report confidence to use harm-reduction practices, selecting the ones that they consider as the most effective ones (e.g. avoid certain environments or people who frequently use). However, the use of harm-reduction practices is buffered under the presence of contextual influences that prevail. Low behavioural awareness, lack of planned alternatives, and long-term habitually established patterns of drug-use behaviours are theorized to lower students’ motivation to engage in preventative or protective health behaviours. Harm reduction interventions should help students identify personally relevant valued-based activities and highlight the role of drugs in disrupting the completion of valued-based activities.

**Social/professional role and identity:** although students present with sufficient awareness of the negative effects of drug-use in their social identity (e.g. academic disruptions, risks in physical safety, reductions in popularity levels, etc.), contextual forces (e.g. acceptability of

**Table 4** Links between the COM-B components and intervention functions

COM-B Components	Intervention Functions								
	Coercion	Education	Enablement	Environmental Restructuring	Incentivisation	Modelling	Persuasion	Restriction	Training
Reflective motivation									
Automatic motivation									
Physical Capability									
Psychological Capability									
Physical Opportunity									
Social Opportunity									

Note: The shaded squares highlight evidence or consensus agreement among the members of the advisory group and show that the identified clusters of target behaviours (8) can be targeted with a particular intervention function or a combination of them

drug-use, peer pressure, fear of not fitting in, etc.) undermine the effect of this awareness on students’ motivation to change (e.g. protect themselves, abstain or reduce the use). Increasing awareness of the negative effects of drugs on students’ identities (e.g. valued self-identity) can support harm-reduction interventions in higher education.

**Goals:** The goal of students who take drug is to fulfil some personal desires (e.g. such as induced fun and excitement while in college). This goal can become habitual, forcing students to either plan specific actions to get drugs or prioritize activities around the drug-use. Findings showed that 50% of students plan to use drugs in time proximity (hours shortly before using), and 38% have a conscious plan (goal) several days in advance. Students report a willingness to abstain or reduce their drug-use if alternative activities will help them to achieve certain value-based outcomes (e.g. academic progress, secure physical safety, etc.). Therefore, altering the means via which students reach desired effects in personally relevant areas of interest (e.g. fun) can support harm-reduction interventions.

**Automatic motivation**

**Reinforcement:** Students’ drug-use (a response) is contingency related with some positive effects (a stimulus) in certain college’s areas of interest (e.g. athletic performance, concentration enhancement, academic achievements). Changing the contingency from having a specific stimulus (desire to achieve positive effects in certain areas of college’s life; e.g. have fun) and a response (take drugs) to acquire a new stimulus (drugs = risky) and a respond (protect myself through harm reduction practices or cease/reduce drug use), could support harm reduction interventions.


**Opportunity**

**Environmental context and resources (Physical):** Each contextual factor (e.g. perceived normalization/acceptability of drug-use within university settings, peer influence, venue, etc.) has a linear effect on students’ decision making; as the contextual factors increase, so does the degree of influence for students’ decision to take drugs. A harm reduction intervention should enhance students’ awareness about the “synergetic” effects of environmental antecedents (personal and interpersonal cues) and their role in increasing social opportunities for drug-use.

In the final step (paradigm coding), we organized the information originating from the previous coding step, into eight narrative statements (clusters of drug-use behaviours), coded around the central phenomena of the COM-B components (see Additional file 1: Table A3; step 4). As presented in Table 3, five clusters target reflective motivation, one automatic motivation, one physical capability, and one social opportunity.

**Step 5—Identify intervention functions**

Five intervention functions were identified from the eight clusters of target behaviours, using the APEASE criteria. The overall reliability of agreement between the raters was moderate  $k = 0.47$  (0.95% CI 33 to 0.60),  $p < 0.001$ . As seen in Table 4, we selected education, modelling, and persuasion as the predominant intervention functions. To address the possible low engagement with the new behavioural repertoires (skills), we included incentivisation, considering that the expectation of rewards in personally relevant behaviours may have reinforcing effects on the target behaviours. Finally, in response to students’ lack of knowledge about implementing harm reduction practices, training was recognized as an important intervention function, mostly because it promotes procedural



knowledge and practical skills (e.g. how to implement harm reduction practices).

**Step 6—Identify policy categories**

We also decided that five policy categories could serve the five selected intervention functions: (a) communication/marketing, (b) guidelines, (c) regulation, (d) service provision, and (e) environmental/social planning. The first three policies were shared across at least four of the five intervention functions. Environmental/social planning was considered a supporting policy for the incentivisation as an intervention function. Both communication/marketing and service provision policies were selected to support post-design promotional and delivery activities of the MyUSE digital intervention, rather than to update its content.

**Step 7—Identify behaviour change techniques**

We created a long list of potential BCTs (see Additional file 1: Table A4). Using the APEASE criteria, we identified 29 BCTs matched with the eight clusters of behaviour to change and COM-B components (see Table 3).

In Fig. 3, we illustrate the combination of the selected BCTs (BCCTv1) matched with the eight clusters of target behaviours.

**The MyUSE digital intervention**

The identified BCTs have led us to develop a multi-component personalized web-based digital intervention. The intervention consists of 11 modules, distributed in three phases: the Allocation phase I (module 1), the Profile building phase II (modules 2 to 4), and the Skill-building phase III (modules 5 to 11). During phase I, the intervention assesses individuals’ drug use history and drug type and allocates users onto one of the following three strands: non-users, low/ moderate users, or severe users. In phase II, individuals build their profile with the risk of harms from drug use and identify areas of skills lacking in relation to harm-reduction practices (see Table 3). During phase III, participants receive a series of personalized skill-building modules, attempting to address their needs, either for harm reduction practices (in case of low/moderate or severe users) or prevention skills (in case of non-using students). For example, individuals may enhance



skills attempting to increase mindful decision making in relation to drug use, help them identify their personalized triggers, behaviours, and consequences of their drug use, and motivate them towards developing their own harm-reduction plan (either drug or non-drug related). A new clinical algorithm that is embedded in the system's interface decision tree logic (see [84]) harnesses individuals' anonymous personal data to present personalized suggestions, and provide modularized intervention's components based on their needs.

## Discussion

To the best of our knowledge, this is the first study to systematically apply the BCW and the TDF frameworks to understand the dynamic and complex determinants supporting harm reduction practices in the context of higher education students' illicit drug-use. The findings have guided the design of the MyUSE intervention through the identification of the contextual, cognitive, and emotional determinants that support students' decision making to use drugs. The findings also generated a novel comprehensive conceptual map of the influences on drug-use behaviours in higher education students. This conceptual map indicates foci for harm reduction implementation practices and new paradigms in addressing drug-use among higher education students.

Increasing reflective and automatic motivation, physical capability, and social opportunity are important determinants to consider in supporting harm-reduction practices for higher education students. Harm reduction practices that respond to these determinants can be translated into selected BCTs which based on the study's theoretically grounded hypothesis, may work synergistically to increase students' mindful decision-making to drug-use, and enhance their motivation that lowers the risk of harm (either drug or non-drug related). The analysis showed that reflective motivation prevails, indicating that any behavioural change intervention should focus on increasing students' reflective motivation. Yet, several other determinants should be also considered. Correcting students' expectancies about the benefits of drug-use in the long-run, increasing insight of finding alternatives to drug-use activities as means for fulfilling students' desires (mostly to have fun and enjoy activities), and enhancing awareness of their personal (e.g. personality, sensation-seeking, emotional dysregulation) and contextual (e.g. peer influences, norm perceptions, etc.) factors that influence students' drug-use decision are all novel implementation practices that can tackle the harm drugs can cause in higher education students populations.

Findings from the discourse pattern-based analysis identified eight patterns of drug-use behaviours. The

eight patterns make use of education, modelling, persuasion, incentivisation, and training as the predominant intervention functions. These functions are supported by communication/marketing, guidelines, regulation, service provision, and environmental/social planning policy categories. Any activity that focuses on cultivating one of the eight clusters of behaviours relevant to drug use should be implemented, using a combination of the 29 identified theory-driven BCTs which fulfil the criteria for interventions being implementable in an affordable, practical, and acceptable way [63]. Notably, BCTs can be used in different modes of delivery, populations, contexts, and relevant types of behaviours [64], and as expected from the TDF framework, can increase clarity as to the mechanisms of action through which behaviour changes occur.

Drawing on the findings from the BCW analysis, this study provides a clear theoretical map for researchers and implementation scientists, highlighting novel context-specific components that can be translated into effective modularized, personalized harm-reduction practices. To achieve this objective, any effort to mitigate the harms drugs can cause in students' lives, require a multicomponent intervention that takes account of the specific developmental context of higher education and the life stage of students [12, 65]. This study illustrates how this can be achieved through the use of the eight clusters of target behaviours. In combination, these behavioural changes may enhance opportunities for creating positive life trajectories via teaching mindful decision-making and value-based actions. They also focus on increasing motivations for change and enhancing sensitivity to contextual influences and opportunities, including drug availability, environmental triggers, and the most salient features of the educational context [21, 46–67]. To date, previous interventional efforts have received criticisms as being too narrow (e.g. misperceived norms, lack of knowledge about harm-reduction practices, low motivation for change, etc.) or adopting a one-size-fits-all approach (e.g. individuals presenting with different levels of use, non-using students, etc.) [27]. Though these interventions are promising [68], they address context variation and personalization for students, poorly [66, 69]. What is missing are innovations in delivering tailored-made harm-reduction supports to students' in higher education.

There is a need for use of contextually driven approaches that can deliver greater behavioural regulation by harnessing social, psychological, and situational forces [38, 70]. Of equal importance is the need for multidisciplinary collaboration in developing and delivering such preventive and intervention programmes. While COM-B and TDF frameworks provide a more granular understanding of psychological capability and reflective

motivational processes [42], this knowledge can also guide researchers and implementation scientists to other relevant theories and approaches [31]. The findings from the BCW approach identified the key role of students' mindful decision-making in relation to drug-use and the promotion of valued-based activities; two components originating from the positive psychology strand of the third-wave cognitive behavioural interventions [72, 73].

Contemporary behavioural accounts of psychological health indicate Psychological Flexibility (PF) [35] as a potentially effective construct to support students' mindful decision-making. PF encourages the disinhibition of immediate habitual gratification (e.g. taking drugs for having fun) over individuals' long-term goals [38, 72]. It does this by teaching individuals behavioural awareness or the ability to be present and take decisions, considering all the possible influencing factors [74]. Several combinations of BCTs, as presented in clusters #4 and #5, teach individuals how to practice mindful decision making. For example, for those declaring previous use, the skill attempts to teach awareness of the triggering influences of behaviours (e.g. contextual and interpersonal) and the consequences (effects) of them prior to decision making. For those with no-previous use, the skill attempts to cultivate awareness of the potential factors that can influence the decision to abstain. The PF model approach also reinforces the recognition of personally desired life directions in domains that are congruent with students' values, such as academic achievement and attenuation of personal career goals [38]. Given that targeted behaviour change can be effective, if they adopt specific approaches [42], PF purports to enact behaviours (e.g. coping strategies) that maintain positive trajectories in youths' lives [72].

Research shows that PF can achieve these goals at higher education [32, 75, 76] and in drug-related behaviours and disorders [77]. This can be achieved by capitalizing on the clinical application of the PF model, coined Acceptance and Commitment Therapy (ACT; [35]) that indicates six therapeutic facets, three of them align to the MyUSE goals: to increase mindfulness, promote the identification of values, and cultivate committed actions [32, 78, 79]. Employing specific evidence-based therapeutic facets, such as the ones the ACT approach proposes [78–80], can increase sensitivity in treatment outcomes and knowledge of the mechanisms via which behavioural changes occur [35]. Finally, conceptualizing harm-reduction practices from a PF perspective can enhance the idea of personal responsibility from a positive psychology perspective [72], as opposed to existing medico-legal

perspectives that highlight users' passivity and risk-taking behaviours [81, 82].

On the ground of these findings, we developed a digital intervention that will "start where the users are at" [83], motivating them to either reduce the harms, lower the use or prevent non-users from potential harms by reinforcing targeted skills. MyUSE approach of harm reduction provides a pragmatic yet compassionate set of skills developed to reduce the harmful consequences and the use itself [40]. This necessitates the development of a multi-component intervention that will equip individuals with tailored skills in identifying, recognizing, and preventing harms before they occur. From a developmental perspective, we believe that these skills will not only help young adults to cope with the effects of drug use, but also, shape behaviours towards value-based and committed actions [72, 73]. Therefore, this intervention should be considered as both a primary prevention intervention for those with no previous experience and a secondary intervention for those with previous or current drug use.

Findings should be interpreted in the light of some limitations. Despite the systematic approach of the BCW approach, the triangulation of the present study data analyses occurred without specifying the primary drug-use. Though most higher education students reporting using cannabis, a sensitivity analysis per drug-use type may have revealed influences on behaviours relevant to specific types of drug-use (e.g. different clusters of behaviours from students using stimulants). This analysis could highlight aspects of drug use types and behavioural responding that are perceived as both having positive and negative effects to individuals. Likewise, the findings from the behavioural analysis focus mainly on understanding the drug-use of experimental/occasional-use students which is the most frequent group of using students [2], with limited references to those students who do not use drugs or those who describe heavy drug use. This limits interpretation of the present study findings to non-using or severe using populations, warranting further research. Further, the interrater reliability for the selection of the intervention's functions was moderate, limiting interpretation of the selected intervention function. Finally, given that no well-validated scales were employed to capture the key indicators of the COM-B components in the survey, the findings may include measurement biases and so replication may be worthwhile.

Future research should assess which identified clusters of behaviours may improve efficacious outcomes if a change is needed. This should be conducted prior to pilot feasibility or pragmatic implementation trials to avoid premature development of a full service which may later need significant modifications. In addition, future research should examine how selected clusters

of behaviours could be effective when applied in different contexts (e.g. digital, public health policy practices, etc.). Furthermore, validation of the eight clusters of behaviours through an in-depth qualitative inquiry of student users and stakeholders (university policymakers) can increase insights as to the implementation barriers which otherwise may not be captured within quantitative data collection, possibly due to the sensitive nature of the topic of drug-use as a research area. Finally, reporting of any innovations (e.g. personalization algorithms, computational models; e.g. [84]), resulting from turning the identified BCTs concepts into digital and interactive modules, can increase knowledge base for behavioural change interventions.

## Conclusions

In conclusion, this research provides an approach to applying the BCW approach to intervention development that draws upon primary and secondary data sources. The findings from the synthesis analysis enabled the identification of targeted problematic behaviours related to drug use. Increasing students' mindful awareness in relation to drug-use decision making and promoting alternatives to drug-use activities indicate foci for the implementation of harm reduction strategies for higher education students' drug-use. These can be delivered through a combination of the eight identified sources of drug-use behaviours. Researchers and implementation scientists can use the presented conceptual map to develop and design interventions and public policy strategies that can be sensitive to mitigate the harms resulting from the use of drugs within the context of higher education.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12954-021-00491-7>.

**Additional file 1.** Main results from the Delphi type exercise, samples from the Pattern-based Grounded Discourse Analysis, and identification of MyUSE BCTs using the APEASE criteria.

## Acknowledgements

The authors gratefully acknowledge the Student Advisory Group (SAG) who assisted in the development of MyUSE and the on-line survey.

## Authors' contributions

All authors conceived the manuscript and participated in the research activities to complete the BCW approach. VSV and SD2 led the writing of the manuscript and revised the manuscript. SD1 conducted the literature search and assisted in the Delphi-type exercise and with the writing of the background and edited the manuscript. MD edited the manuscript and assisted in the ranking exercise of the policy categories. CL edited the manuscript and assisted in the intervention function and BCTs identification exercises. CH edited the manuscript. MB edited the manuscript and provided overall supervision of the project. Also, MB was guarantor of the study. All authors read and approved the final manuscript.

## Funding

The research leading to these results has received funding from UCC. The funding body was not involved in the study design, collection, analysis or interpretation of the data, or in writing this manuscript.

## Availability of data and materials

Not applicable.

## Declarations

### Ethics approval and consent to participate

Ethical approval for this project was granted by the Social Research Ethics Committee at the University College Cork (SREC reference number no: 2018-072A).

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### Author details

<sup>1</sup>School of Applied Psychology, University College Cork, N Mall, Kilbarry Enterprise Centre, Cork Enterprise Centre, Cork, Ireland. <sup>2</sup>School of Public Health, University College Cork, Fourth Floor, Western Gateway Building, Cork, Ireland. <sup>3</sup>Cork Sexual Health Centre, 16 Peter's Street, Centre, Cork, Ireland. <sup>4</sup>Cork University Business School, University College Cork, West Wing, Main Quadrangle, Cork, Ireland. <sup>5</sup>Student Health Department, University College Cork, Ardpark College Road, Cork, Ireland.

Received: 4 December 2020 Accepted: 31 March 2021

Published online: 20 May 2021

## References

- Smyth BP, Cannon M. Cannabis and public health—a need to reclaim the narrative. *Ir J Med Sci*. 2021. <https://doi.org/10.1007/s11845-021-02570-x>.
- Skidmore CR, Kaufman EA, Crowell SE. Substance use among college students. *Child Adolesc Psychiatr Clin N Am*. 2016;25(4):735–53.
- European Union, European Monitoring Centre for Drugs and Drug Addiction. European drug report: trends and developments 2019. Luxembourg: Publications Office of the European Union; 2019.
- Silvestri MM, Knight H, Britt J, Correia CJ. Beyond risky alcohol use: screening non-medical use of prescription drugs at National Alcohol Screening Day. *Addict Behav*. 2015;43:25–7.
- Arria AM, Caldeira KM, Allen HK, Bugbee BA, Vincent KB, O'Grady KE. Prevalence and incidence of drug use among college students: an 8-year longitudinal analysis. *Am J Drug Alcohol Abuse*. 2017;43(6):711–8.
- Bjork JM, Pardini DA. Who are those “risk-taking adolescents”? Individual differences in developmental neuroimaging research. *Dev Cogn Neurosci*. 2015;11:56–64.
- Dalton ED, Hammen CL. Independent and relative effects of stress, depressive symptoms, and affect on college students' daily health behaviors. *J Behav Med*. 2018;41(6):863–74.
- Acharya L, Jin L, Collins W. College life is stressful today – Emerging stressors and depressive symptoms in college students. *J Am Coll Health*. 2018;66(7):655–64.
- Awang MM, Kutty FM, Ahmad AR. Perceived social support and well being: first-year student experience in university. *IES*. 2014;7(13):261.
- Caldeira KM, Kasperski SJ, Sharma E, Vincent KB, O'Grady KE, Wish ED, et al. College students rarely seek help despite serious substance use problems. *J Subst Abuse Treat*. 2009;37(4):368–78.
- White HR, Labouvie EW, Papadaratsakis V. Changes in substance use during the transition to adulthood: a comparison of college students and their noncollege age peers. *J Drug Issues*. 2005;35(2):281–306.
- Schwartz SJ, Petrova M. Prevention science in emerging adulthood: a field coming of age. *Prev Sci*. 2019;20(3):305–9.
- Lenton S. The essence of prevention. In: *Perspectives on Addiction: Making Sense of the Issues*. Perth: William Montgomery; 1996. p. 73–80.

14. Arria AM, Caldeira KM, Bugbee BA, Vincent KB, O'Grady KE. The academic consequences of marijuana use during college. *Psychol Addict Behav*. 2015;29(3):564–75.
15. O'Grady KE, Arria AM, Fizzle DMB, Wish ED. Heavy drinking and polydrug use among college students. *J Drug Issues*. 2008;38(2):445–65.
16. Meshesha L, Dennhardt A, Murphy GJ. Polysubstance use is associated with deficits in substance-free reinforcement in college students. *J Stud Alcohol Drugs*. 2015;76(1):106–16.
17. McCabe SE, Teter CJ. Drug use related problems among nonmedical users of prescription stimulants: a web-based survey of college students from a Midwestern university. *Drug Alcohol Depend*. 2007;91(1):69–76.
18. Pedrelli P, Nyer M, Yeung A, Zulauf C, Wilens T. College students: mental health problems and treatment considerations. *Acad Psychiatry*. 2015;39(5):503–11.
19. Schuster C, O'Malley PM, Bachman JG, Johnston LD, Schulenberg J. Adolescent marijuana use and adult occupational attainment: a longitudinal study from age 18 to 28. *Subst Use Misuse*. 2001;36(8):997–1014.
20. Taylor DR, Poulton R, Moffitt TE, Ramankutty P, Sears MR. The respiratory effects of cannabis dependence in young adults. *Addiction*. 2000;95(11):1669–77.
21. Stockings E, Hall WD, Lynskey M, Morley KI, Reavley N, Strang J, et al. Prevention, early intervention, harm reduction, and treatment of substance use in young people. *Lancet Psychiatry*. 2016;3(3):280–96.
22. Lee CM, Neighbors C, Kilmer JR, Larimer ME. A brief, web-based personalized feedback selective intervention for college student marijuana use: a randomized clinical trial. *Psychol Addict Behav*. 2010;24(2):265–73.
23. Elliott JC, Carey KB, Venable PA. A preliminary evaluation of a web-based intervention for college marijuana use. *Psychol Addict Behav*. 2014;28(1):288–93.
24. Palfai TP, Saitz R, Winter M, Brown TA, Kypri K, Goodness TM, et al. Web-based screening and brief intervention for student marijuana use in a university health center: pilot study to examine the implementation of eCHECKUP TO GO in different contexts. *Addict Behav*. 2014;39(9):1346–52.
25. Boumparis N, Karyotaki E, Schaub MP, Cuijpers P, Riper H. Internet interventions for adult illicit substance users: a meta-analysis: Internet interventions for illicit substances. *Addiction*. 2017;112(9):1521–32.
26. Gulliver A, Farrer L, Chan JK, Tait RJ, Bennett K, Calear AL, et al. Technology-based interventions for tobacco and other drug use in university and college students: a systematic review and meta-analysis. *Addict Sci Clin Pract*. 2015;10(1):5.
27. Dick S, Whelan E, Davoren MP, Dockray S, Heavin C, Linehan C, et al. A systematic review of the effectiveness of digital interventions for illicit substance misuse harm reduction in third-level students. *BMC Public Health*. 2019;19(1):1244.
28. Organ DD, Dick S, Hurley C, Heavin C, Linehan C, Dockray S, et al. A systematic review of user-centred design practices in illicit substance use interventions for higher education students. 2018;18.
29. Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci*. 2011;6(1):42.
30. Ovretveit J. Understanding the conditions for improvement: research to discover which context influences affect improvement success. *BMJ Qual Saf*. 2011;20(Suppl 1):i18–23.
31. Atkins L, Francis J, Islam R, O'Connor D, Patey A, Ivers N, et al. A guide to using the Theoretical Domains Framework of behaviour change to investigate implementation problems. *Implement Sci*. 2017;12(1):77.
32. Levin ME, Krafft J, Twohig MP. Examining processes of change in an online acceptance and commitment therapy dismantling trial with distressed college students. *J Context Behav Sci*. 2020;17:10–6.
33. Hayes SC, Hofmann SG, Stanton CE, Carpenter JK, Sanford BT, Curtiss JE, et al. The role of the individual in the coming era of process-based therapy. *Behav Res Ther*. 2019;117:40–53.
34. Hofmann SG, Hayes SC. The future of intervention science: process-based therapy. *Clin Psychol Sci*. 2019;7(1):37–50.
35. Hayes SC, Levin ME, Plumb-Villardaga J, Villatte JL, Pistorello J. Acceptance and commitment therapy and contextual behavioral science: examining the progress of a distinctive model of behavioral and cognitive therapy. *Behav Ther*. 2013;44(2):180–98.
36. Michie S, Johnston M. Theories and techniques of behaviour change: developing a cumulative science of behaviour change. *Health Psychol Rev*. 2012;6(1):1–6.
37. Simons JS, Joseph Clarke C, Simons RM, Spelman PJ. Marijuana consequences in a motivational context: goal congruence reduces likelihood of taking steps toward change. *Addict Behav*. 2016;52:83–90.
38. Zhang C-Q, Leeming E, Smith P, Chung P-K, Hagger MS, Hayes SC. Acceptance and commitment therapy for health behavior change: a contextually-driven approach. *Front Psychol*. 2018;8:2350.
39. Prochaska JO, Redding CA, Evers KA. The transtheoretical model and stages of change. In: *Health behavior: Theory, research, and practice* [Internet]. 4th ed. John Wiley & Sons, Inc.; p. 125–48. (2008). Available from: file:///C:/Users/v.vasilioi/OneDrive%20-%20University%20College%20Cork/Desktop/Theory\_of\_reasoned\_action\_theory\_of\_planned\_behavi.pdf
40. Leslie K. Harm reduction: an approach to reducing risky health behaviours in adolescents. *Paediatr Child Health*. 2008;13(1):53–6.
41. Michie S, Atkins L, West R. The behaviour change wheel: a guide to designing interventions. Silverback; 2014. 329 p.
42. Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W, et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med*. 2013;46(1):81–95.
43. Grol R, Grimshaw J. From best evidence to best practice: effective implementation of change in patients' care. *Lancet*. 2003;362(9391):1225–30.
44. Niilen P. Making sense of implementation theories, models and frameworks. *Implement Sci*. 2015;10(1):53.
45. Dugdale S, Ward J, Hernen J, Elison S, Davies G, Donkor D. Using the Behavior Change Technique Taxonomy v1 to conceptualize the clinical content of Breaking Free Online: a computer-assisted therapy program for substance use disorders. *Subst Abuse Treat Prev Policy*. 2016;11(1):26.
46. Dick S, Vasilioi SV, Davoren MP, Dockray S, Heavin C, Linehan C, et al. My Understanding of Drug Use Experiences (MIUSE): a protocol for the development of a digitally delivered harm reduction intervention for students in higher education. *JMIR Research Protocols*, 2020;9(8):e17829. Available from: <https://www.researchprotocols.org/2020/8/e17829>.
47. Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey technique. *J Adv Nurs*. 2000;32(4):1008–15.
48. Whelan E, Dockray S, Dick S, Davoren M, Heavin C, Linehan C, Byrne M. Motivations to decrease and cease substance use in third-level students: A scoping review. 2020. Available at <https://psyarxiv.com/5u7bs/download?format=pdf>.
49. Sumson T. The Delphi technique: an adaptive research tool. *Br J Occup Ther*. 1998;61(4):153–6.
50. Creswell JW, Creswell JD. *Research design: Qualitative, quantitative, and mixed methods approaches*. 6th ed. London: Sage Publications; 2017. p. 304.
51. Charmaz K. *Constructing grounded theory*. London: Sage Publications; 2006. p. 208.
52. Birks M, Mills J. *Grounded theory: a practical guide*. Los Angeles: Sage; 2011. p. 210.
53. Strauss AL, Corbin JM. *Basics of qualitative research: techniques and procedures for developing grounded theory*. 2nd ed. Thousand Oaks: Sage Publications; 1998. p. 312.
54. Clarke AE. Situating grounded theory and situational analysis in interpretive qualitative inquiry. In: *The SAGE handbook of current developments in grounded theory* [Internet]. 1st ed. Sage Publications; 2019. p. 3–48. Available from: <https://uk.sagepub.com/en-gb/eur/the-sage-handbook-of-current-developments-in-grounded-theory/book249299>
55. Coyle A. Discourse Analysis. In: *Analysing Qualitative Data in Psychology* [Internet]. London: SAGE Publications, Ltd; 2007. p. 98–116. (SAGE Research methods). Available from: <https://methods.sagepub.com/base/download/BookChapter/analysing-qualitative-data-in-psychology/d14.xml>
56. Meissner H, Creswell J, Klassen AC, Plano V, Smith KC. *Best practices for mixed methods research in the health sciences*. Bethesda: National Institutes of Health; 2013. p. 541–5.
57. Bishop BJ, Sonn CC, Drew NM, Contos NE. The evolution of epistemology and concepts in an iterative-generative reflective practice: the importance of small differences. *Am J Community Psychol*. 2002;30(4):493–510.



58. Braun V, Clarke V. *Successful Qualitative Research: A Practical Guide for Beginners* [Internet]. illustrated, reprint ed. SAGE; 2014. 400 p. Available from: [https://books.google.ie/books/about/Successful\\_Qualitative\\_Research.html?id=PVBGngEACAAJ&redir\\_esc=y](https://books.google.ie/books/about/Successful_Qualitative_Research.html?id=PVBGngEACAAJ&redir_esc=y)
59. Fine M, Weis L. Writing the 'Wrongs' of Fieldwork: Confronting Our Own Research/ Writing Dilemmas in Urban Ethnographies. :24.
60. Vasilioi VS, Dick S, Heavin C, Davoren MP, Dockray S, Linehan C, Byrne M. "One second now, I am going to check what is going to happen me from this": Decomposing Perspectives of Harm-reduction at Student University Level Using the Theoretical Domain Framework (TDF) (in preparation).
61. Fleiss JL, Bruce L, Paik MC. *Statistical Methods for Rates and Proportions* [Internet]. 3rd ed. Wiley; 800 p. Available from: <https://www.wiley.com/en-ie/Statistical+Methods+for+Rates+and+Proportions%2C+3rd+Edition-p-9780471526292>
62. Davoren MP, Calnan S, Mulcahy J, Lynch E, Perry IJ, Byrne M. Responding to excessive alcohol consumption in third-level (REACT): a study protocol. *BMC Health Serv Res*. 2018;18(1):364.
63. Presseau J, McCleary N, Lorencatto F, Patey AM, Grimshaw JM, Francis JJ. Action, actor, context, target, time (AACTT): a framework for specifying behaviour. *Implement Sci*. 2019;14(1):102.
64. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ*. 2008;337.
65. Arnett JJ. The developmental context of substance use in emerging adulthood. *J Drug Issues*. 2005;35(2):235–54.
66. Jenkins EK, Slemmon A, Haines-Saah RJ. Developing harm reduction in the context of youth substance use: insights from a multi-site qualitative analysis of young people's harm minimization strategies. *Harm Reduct J*. 2017;14(1):53.
67. Dick S, Davoren MP, Byrne M, Heavin C, Dockray S, Vasilioi V, et al. The Design of a Digital Behaviour Change Intervention for Third-Level Student Illicit Substance Use: A Persona Building Approach. In: *Proceedings of the 53rd Hawaii International Conference on System Sciences* [Internet]. Hawaii; p. 9. Available from: <http://hdl.handle.net/10125/64157>.
68. Dennhardt AA, Murphy JG. Prevention and treatment of college student drug use: a review of the literature. *Addict Behav*. 2013;38(10):2607–18.
69. Farrugia A. Assembling the dominant accounts of youth drug use in Australian harm reduction drug education. *International Journal of Drug Policy*. 2014;25(4):663–72.
70. Hansen S, Kanning M, Lauer R, Steinacker JM, Schlicht W. MAP-IT: a practical tool for planning complex behavior modification interventions. *Health Promot Pract*. 2017;18(5):696–705.
71. Connell Bohlen L, Michie S, de Bruin M, Rothman A, Kelly MP, Groarke H, et al. Do combinations of behaviour change techniques that occur frequently in interventions reflect underlying theory? [Internet]. *PsyArXiv*; 2019 Dec [cited 2020 Jul 24]. Available from: <https://osf.io/49djm>
72. Ciarrochi J, Atkins PWB, Hayes LL, Sahdra BK, Parker P. Contextual positive psychology: policy recommendations for implementing positive psychology into schools. *Front Psychol*. 2016. <https://doi.org/10.3389/fpsyg.2016.01561/full>.
73. Garland EL, Farb NA, Goldin P, Fredrickson BL. Mindfulness broadens awareness and builds eudaimonic meaning: a process model of mindful positive emotion regulation. *Psychol Inq*. 2015;26(4):293–314.
74. Hayes SC, Villatte M, Levin M, Hildebrandt M. Open, aware, and active: contextual approaches as an emerging trend in the behavioral and cognitive therapies. *Annu Rev Clin Psychol*. 2011;7(1):141–68.
75. Levin ME, Hicks ET, Krafft J. Pilot evaluation of the stop, breathe & think mindfulness app for student clients on a college counseling center waitlist. *J Am Coll Health*. 2020:1–9.
76. Räsänen P, Lappalainen P, Muotka J, Tolvanen A, Lappalainen R. An online guided ACT intervention for enhancing the psychological wellbeing of university students: a randomized controlled clinical trial. *Behav Res Ther*. 2016;78:30–42.
77. Li T, Sato H, Watanabe N, Kondo M, Masuda A, Hayes SC, et al. Psychological flexibility-based interventions versus first-line psychosocial interventions for substance use disorders: systematic review and meta-analyses of randomized controlled trials. *J Context Behav Sci*. 2019;13:109–20.
78. Fledderus M, Bohlmeijer ET, Fox J-P, Schreurs KMG, Spinhoven P. The role of psychological flexibility in a self-help acceptance and commitment therapy intervention for psychological distress in a randomized controlled trial. *Behav Res Ther*. 2013;51(3):142–51.
79. Gloster AT, Meyer AH, Lieb R. Psychological flexibility as a malleable public health target: Evidence from a representative sample. *J Context Behav Sci*. 2017;6(2):166–71.
80. Levin ME, Hildebrandt MJ, Lillis J, Hayes SC. The impact of treatment components suggested by the psychological flexibility model: a meta-analysis of laboratory-based component studies. *Behav Ther*. 2012;43(4):741–56.
81. Engel LB, Bright SJ, Barratt MJ, Allen MM. Positive drug stories: possibilities for agency and positive subjectivity for harm reduction. *Addict Res Theory*. 2020:1–9.
82. Palamar JJ, Halkitis PN, Kiang MV. Perceived public stigma and stigmatization in explaining lifetime illicit drug use among emerging adults. *Addict Res Theory*. 2013;21(6):516–25.
83. Marlatt GA. Harm reduction: come as you are. *Addict Behav*. 1996;21(6):779–88.
84. Vasilioi SV, Dillan B, Dick S, Davoren MP, Dockray S, Heavin C, Linehan C, Byrne M. A new clinical algorithm embedded in a contextual behavior change intervention for higher education student drug use. In *Proceedings of the International Conference on Information Systems, Virtual Pre-ICIS (International Conference on Information Systems)*. 2021.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

